arXiv:1111.0632v2 [math.SG] 21 Apr 2012

HOMOLOGICAL MIRROR SYMMETRY FOR CALABI-YAU HYPERSURFACES IN PROJECTIVE SPACE

NICK SHERIDAN

Abstract. We prove Homological Mirror Symmetry for a smooth d-dimensional Calabi-Yau hypersurface in projective space, for any $d \geq 3$ (for example, $d = 3$ is the quintic three-fold). The main techniques involved in the proof are: the construction of an immersed Lagrangian sphere in the 'd-dimensional pair of pants'; the introduction of the 'relative Fukaya category', and an understanding of its grading structure; a description of the behaviour of this category with respect to branched covers (via an 'orbifold' Fukaya category); a Morse-Bott model for the relative Fukaya category that allows one to make explicit computations; and the introduction of certain graded categories of matrix factorizations mirror to the relative Fukaya category.

Contents

1. Introduction 3

1.1. Mirror symmetry context 3

1.2. Statement of the main result 4

1.3. Affine and relative Fukaya categories 6

1.4. The B-model mirror to the affine, relative and full Fukaya categories 9

1.5. Behaviour of the Fukaya category under branched covers 14

1.6. The Fukaya category of M^n 15

2. Graded and equivariant categories 20

2.1. Grading data 20

2.2. Graded vector spaces 23

2.3. $\boldsymbol{G}$-graded A_∞ algebras and Hochschild cohomology 26

2.4. Deformations of A_∞ algebras 34

This research was partially supported by NSF Grant DMS-0652620.

2.5. Computations 38

2.6. First-order deformations 53

3. The affine Fukaya category 53

3.1. Grading data from the Lagrangian Grassmannian 54

3.2. Anchored branes 56

3.3. The affine Fukaya category 57

3.4. Covers 60

3.5. The relative case 61

4. Moduli Spaces of Disks 65

4.1. Moduli spaces of holomorphic spheres and disks 65

4.2. Deligne-Mumford compactifications 66

4.3. Moduli spaces of pseudoholomorphic disks 70

4.4. Indices and orientations 73

4.5. Gromov compactness 76

4.6. Branched covers 84

5. The relative Fukaya category 86

5.1. The definition 86

5.2. Behaviour with respect to ramified covers 88

6. Morse-Bott computations in the Fukaya category 93

6.1. The Lagrangian immersion $L^n : S^{n-2} \to M \setminus \boldsymbol{D}$ 94

6.2. Flipping Pearly trees 97

6.3. Floer and perturbation data 102

6.4. Transversality and compactness 108

6.5. Morse-Bott model for the first-order Fukaya category 110

7. The B-model 115

7.1. Homological perturbation lemma 115

7.2. Matrix factorization computations 116

7.3. $\boldsymbol{G}$-Graded matrix factorizations 119

7.4. Equivariant matrix factorizations 121

7.5. Coherent sheaves 122

8. The full Fukaya category 125

8.1. Relating the full and relative Fukaya categories 125

8.2. Split-generation 127

Appendix A. Strict group actions on Fukaya categories 129

References 130

Acknowledgments. I would like to thank my advisor, Paul Seidel, for all of his help with this work. These results owe a great deal to him, both because his previous work [1, 2, 3, 4] laid the foundations for them, and because he gave me a lot of guidance and made many useful suggestions along the way. I would also like to thank Mohammed Abouzaid for many helpful discussions and suggestions, and for showing me a preliminary version of [5]. I would also like to thank Grisha Mikhalkin for helping me to find the construction of the immersed Lagrangian sphere in the pair of pants, on which this work is based.

1. Introduction

1.1. Mirror symmetry context.

The mirror symmetry phenomenon was discovered by string theorists. In its original form, it dealt with Calabi-Yau Kähler manifolds. On such a manifold, one can define symplectic invariants (the 'A-model') and complex invariants (the 'B-model'). Broadly, mirror symmetry says that there exist pairs of manifolds (M, N) such that the A-model on M is equivalent to the B-model on N, and vice versa.

The closed-string version of the A-model is encoded in Gromov-Witten invariants (the quantum cohomology ring), and the closed-string version of the B-model is encoded in period integrals of the holomorphic volume form. Mirror symmetry first came to the attention of mathematicians in 1991, when Candelas, de la Ossa, Green and Parkes [6] applied it to make predictions about rational curve counts on the quintic three-fold M. They constructed a manifold N which ought to be mirror to M, and computed the closed-string B-model on N. Assuming mirror symmetry to hold, this allowed them to predict the closed-string A-model on M, from which they extracted rational curve counts. This closed-string version of mirror symmetry was proven by Givental for Calabi-Yau (and Fano) complete intersections in toric varieties in 1996 [7, 8].

In the meantime, Kontsevich had introduced the open-string version of mirror symmetry, called the Homological Mirror Symmetry conjecture [9] in 1994. He proposed that the A-model should be encoded in the Fukaya category, and the B-model should be encoded in the category of coherent sheaves. Then if M and N are mirror Calabi-Yaus, there should be an equivalence of the Fukaya category of M with the category of coherent sheaves on Y (on the derived level). Complete or partial proofs of homological mirror symmetry for Calabi-Yau varieties are known for elliptic curves [10, 11], abelian varieties [12] (see [13] for the case of the four-torus), Strominger-Yau-Zaslow dual torus fibrations [14], and the quartic K3 surface [2].

In this paper we consider a smooth Calabi-Yau hypersurface $M^n \subset \mathbb{CP}^{n-1}$, and its (predicted) mirror N^n. Our main result is to prove one direction of homological mirror symmetry: we prove that there is an equivalence of triangulated categories between the split-closed derived Fukaya category of M^n and the bounded derived category of coherent sheaves on N^n (see Theorem 1 for the precise statement). M^3 is the elliptic curve considered by Polishchuk and Zaslow, M^4 is the quartic surface considered by Seidel, and M^5 is the quintic three-fold. We remark that Nohara and Ueda have also considered the case of the quintic three-fold [15], using the results of Seidel [2] and our earlier paper [16].

In future work [17], we will extend this result to the case of Fano hypersurfaces in projective space, and we plan also to consider the case of hypersurfaces of general type.

1.2. Statement of the main result.

First let us introduce the coefficient rings of the categories we will be considering.

Definition 1.2.1. We define the **universal Novikov ring** Λ_0, whose elements are formal sums

$$\psi(r) = \sum_{j=1}^{\infty} c_j r^{\lambda_j},$$

where $c_j \in \mathbb{C}$, and $\lambda_j \in \mathbb{R}_{\geq 0}$ is an increasing sequence of non-negative reals such that

$$\lim_{j\to\infty} \lambda_j = \infty.$$

It is a valuation ring whose value group is $\mathbb{R}$. Its field of fractions is the **universal Novikov field**

$$\Lambda := \Lambda_0[r^{-1}].$$

We denote the maximal ideal by $\mathfrak{m} \subset \Lambda_0$.

Definition 1.2.2. Given an element $\psi \in \mathbb{C}[[r]]$ such that $\psi(0) \neq 0$, we obtain an automorphism

$$\begin{aligned} f_\psi : \Lambda &\rightarrow \Lambda, \text{ such that} \\ f_\psi(r) &= r\psi(r). \end{aligned}$$

If $\mathcal{C}$ is a Λ-linear category, and ψ such an element, we define a new Λ-linear category $\psi \cdot \mathcal{C}$: it is the same category as $\mathcal{C}$, but the Λ-vector space structure of the morphism spaces is changed via f_ψ. We say that $\psi \cdot \mathcal{C}$ is obtained from $\mathcal{C}$ via a **change of variables** in the Novikov field.

Now we introduce the relevant categories.

Definition 1.2.3. On the symplectic side, let $M^n \subset \mathbb{CP}^{n-1}$ be a smooth hypersurface of degree n. M^n is an $(n-2)$-dimensional Calabi-Yau. The Fukaya category $\mathcal{F}(M^n)$ (as defined in [5]) is a $\mathbb{Z}$-graded Λ-linear A_∞ category. The split-closed derived Fukaya category $D^\pi\mathcal{F}(M^n)$ (see [3, Section I.4]) is a Λ-linear triangulated category. We remark that the Fukaya category is a symplectic invariant (up to A_∞ quasi-isomorphism), so it does not matter which smooth hypersurface M^n we choose.

Definition 1.2.4. On the algebraic side, we define

$$w_{nov} := u_1 \dots u_n + r\sum_{j=1}^{n} u_j^n \in \Lambda[u_1, \dots, u_n].$$

We set $\widetilde{N}^n_{nov} := \{w_{nov} = 0\} \subset \mathbb{P}^{n-1}_\Lambda$. We equip $\widetilde{N}^n_{nov}$ with the action of a finite group. Observe that the group

$$\tilde{\Gamma}^*_n := (\mathbb{Z}_n)^n/(1,1,\dots,1)$$

acts on $\mathbb{P}^{n-1}_\Lambda$ by multiplying the homogeneous coordinates u_j by nth roots of unity (we include the '$*$' for consistency with later notation). We have a homomorphism

$$\tilde{\Gamma}^*_n \to \mathbb{Z}_n,$$

given by summing the coordinates. We denote the kernel of this homomorphism by Γ^*_n. Note that the action of Γ^*_n preserves $\widetilde{N}^n_{nov}$, so Γ^*_n acts on $\widetilde{N}^n_{nov}$. We define

$$N^n_{nov} := \widetilde{N}^n_{nov}/\Gamma^*_n.$$

We consider the bounded derived category of coherent sheaves on N^n_{nov}, which is

$$D^bCoh(N^n_{nov}) := D^b\left(Coh^{\Gamma^*_n}\left(\widetilde{N}^n_{nov}\right)\right).$$

It is a triangulated category over Λ.

Theorem 1. *If $n \ge 5$, then there exists a power series $\psi \in \mathbb{C}[[r]]$, $\psi(0) = \pm 1$, and an equivalence of Λ-linear triangulated categories*

$$D^\pi\mathcal{F}(M^n) \cong \psi \cdot D^bCoh(N^n_{nov}).$$

In fact, we will see that $\psi \in \mathbb{C}[[r^n]] \subset \mathbb{C}[[r]]$.

Remark 1.2.5. The requirement that $n \ge 5$ can be removed without difficulty, but would require some ad-hoc arguments in the cases $n = 3$ and $n = 4$. We prefer to leave these to the reader, as they distract from the main flow of the argument.

We remark that the reference [5], in which the full Fukaya category $\mathcal{F}(M^n)$ is defined, and one of the results we use in the proof of Theorem 1 is proven (we have stated it as Theorem 5), is still in preparation. This is not an ideal situation, but we reassure the reader that this paper is written with minimal reliance on [5]. Let us explain what we mean.

We rigorously define a version of the Fukaya category called the 'relative' Fukaya category $\mathcal{F}(M^n, \boldsymbol{D})$. It is an A_∞ category defined over some coefficient ring R, and there are natural homomorphisms of $\mathbb{C}$-algebras

$$R \to \mathbb{C}[[r]] \hookrightarrow \Lambda.$$

We also define a certain category of matrix factorizations, denoted by $\boldsymbol{p}_1^* MF^{\boldsymbol{G}}(S,w)$, which is a differential graded category over the same coefficient ring R. We introduce full subcategories

$$\widetilde{\mathcal{A}} \subset \mathcal{F}(M^n, \boldsymbol{D}), \quad \widetilde{\mathcal{B}} \subset \boldsymbol{p}_1^* MF^{\boldsymbol{G}}(S,w),$$

and prove (without reference to [5]) that there exists $\psi \in R$ and an A_∞ equivalence

$$\psi \cdot \widetilde{\mathcal{B}} \cong \widetilde{\mathcal{A}}.$$

For a more precise statement, see Theorem 2. Using a theorem of Orlov relating matrix factorizations to coherent sheaves, we show that there is an equivalence

$$D^b Coh(N^n_{nov}) \cong \mathrm{Ho}\left(\boldsymbol{p}_1^* MF^{\boldsymbol{G}}(S,w) \otimes_R \Lambda\right)$$

(the 'Ho' on the right-hand side denotes the homotopy category of a differential graded category), and the full subcategory $\widetilde{\mathcal{B}} \otimes_R \Lambda$ generates. It follows that we have a fully faithful embedding

$$\psi \cdot D^b Coh(N^n_{nov}) \hookrightarrow D^b\left(\mathcal{F}(M^n, \boldsymbol{D}) \otimes_R \Lambda\right).$$

We can come to this point without reference to [5], but no further.

In Section 8, we make the assumption (Assumption 8.1.1) that there is a fully faithful embedding

$$\mathcal{F}(M^n, \boldsymbol{D}) \otimes_R \Lambda \hookrightarrow \mathcal{F}(M^n),$$

where $\mathcal{F}(M^n)$ is defined as in [5]. We justify this assumption 'heuristically', and show that the full subcategory

$$\widetilde{\mathcal{A}} \otimes_R \Lambda \subset \mathcal{F}(M^n)$$

split-generates, by applying the split-generation criterion Theorem 5 (also due to [5]). This completes the proof of Theorem 1.

In the rest of this section, we give an overview of the main techniques introduced in the rest of the paper, and how they are used to prove Theorem 1. We will make a few imprecise statements and definitions, in the interest of giving the reader the correct intuitive picture.

1.3. **Affine and relative Fukaya categories.** Let M be a Kähler manifold, with a tuple $\boldsymbol{D} = (D_1, \ldots, D_k)$ of smooth, ample, irreducible divisors with normal crossings. In this paper, we consider three versions of the Fukaya category: the **affine Fukaya category** $\mathcal{F}(M \setminus \boldsymbol{D})$, the **relative Fukaya category** $\mathcal{F}(M, \boldsymbol{D})$, and the **full Fukaya category** $\mathcal{F}(M)$.

First let us describe the affine Fukaya category. It is closely related to the exact Fukaya category of the exact symplectic manifold $M \setminus \boldsymbol{D}$, as defined by Seidel in [3], but it has a more interesting grading structure.

First we explain the objects of the affine Fukaya category. If P is a symplectic manifold, we denote by $\mathcal{G}P \to P$ the Lagrangian Grassmannian, whose fibre over $p \in P$ is the space of Lagrangian subspaces of T_pP. Any smooth Lagrangian immersion $i : L \to P$ comes with an associated lift $i_* : L \to \mathcal{G}P$. We define an **anchored Lagrangian brane** $L^{\#}$ in P to be a

Lagrangian immersion $i : L \to P$, together with a lift

$$\begin{tikzcd} & \widetilde{\mathcal{G}}P \arrow[d] \\ L \arrow[ur, "i^{\#}"] \arrow[r, "i_*"] & \mathcal{G}P, \end{tikzcd}$$

where $\widetilde{\mathcal{G}}P$ is the universal cover of the total space of $\mathcal{G}P$, together with a choice of Pin structure on L. The terminology comes from [18], where a closely-related concept is studied.

We define the objects of the affine Fukaya category $\mathcal{F}(M \setminus \boldsymbol{D})$ to be compact, exact, embedded, anchored Lagrangian branes in $M \setminus \boldsymbol{D}$. The category is $\mathbb{C}$-linear, and morphism spaces and A_∞ structure maps are defined in exactly the same way as in the exact Fukaya category of the exact symplectic manifold $M \setminus \boldsymbol{D}$. Namely, the morphism spaces are

$$CF^*_{\mathcal{F}(M\setminus\boldsymbol{D})}(L_0, L_1) := \bigoplus_{p \in L_0 \cap L_1} \mathbb{C} \cdot p$$

(with appropriate modifications to allow for non-transverse intersection of L_0 with L_1). The A_∞ structure map

$$\mu^s : CF^*(L_{s-1}, L_s) \otimes \ldots \otimes CF^*(L_0, L_1) \to CF^*(L_0, L_s)$$

is defined by counting rigid holomorphic disks with $s+1$ boundary punctures in $M\setminus\boldsymbol{D}$. Namely, the coefficient of p_0 in $\mu^s(p_s, \ldots, p_1)$ is given by the signed count of rigid holomorphic disks

$$u : \mathbb{D} \setminus \{\zeta_0, \ldots, \zeta_s\} \to M \setminus \boldsymbol{D},$$

sending the jth boundary component to Lagrangian L_j, and asymptotic at puncture ζ_j to intersection point p_j. However, we treat the grading differently.

To start with, we equip each morphism space with a grading in the abelian group $H_1(M\setminus\boldsymbol{D})$. The A_∞ structure maps respect this grading, essentially because if there is a holomorphic disk contributing to some A_∞ product in $M \setminus \boldsymbol{D}$, then its boundary is nullhomologous in $M \setminus \boldsymbol{D}$. Gradings of this type have appeared in, for example, [4, 16].

Remark 1.3.1. Another way of defining this grading (following [18]) would be to equip M with a basepoint $q \in M \setminus \boldsymbol{D}$, and define an object of the Fukaya category to be a Lagrangian $L \subset M \setminus \boldsymbol{D}$, equipped with a path from q to a point $q_L \in L$, inside $M \setminus \boldsymbol{D}$. Then, given an intersection point $p \in L_0 \cap L_1$, which by definition is a generator of $CF^*(L_0, L_1)$, we can define a class in $H_1(M \setminus \boldsymbol{D})$: start from q and follow the path from q to q_{L_0}, then follow a path in L_0 to the intersection point p, then follow a path in L_1 to q_{L_1}, then follow the path back to q. The class of this path in $H_1(M \setminus \boldsymbol{D})$ defines the grading of p.

Secondly, if we were given a quadratic complex volume form on $M \setminus \boldsymbol{D}$, we could define a $\mathbb{Z}$-grading of the morphism spaces (as in [3]). This defines a $\mathbb{Z} \oplus H_1(M \setminus \boldsymbol{D})$-graded category. However, this formulation is unsatisfactory: the $\mathbb{Z}$-grading and H_1-grading are related. For example, changing the quadratic volume form has the effect of changing the $\mathbb{Z} \oplus H_1$ grading by an automorphism preserving the H_1 factor. We really want a new notion of grading.

In Section 2, we define a **grading datum $\boldsymbol{G}$** to be an abelian group Y together with a morphism $f : \mathbb{Z} \to Y$. We say that an A_∞ category is **$\boldsymbol{G}$-graded** if its morphism spaces are Y-graded, and the A_∞ structure map μ^s has degree $f(2-s)$. We also study the deformation theory of $\boldsymbol{G}$-graded A_∞ algebras and categories, and prove various classification results about them.

In Section 3.1, we introduce a grading datum $\boldsymbol{G}(M,\boldsymbol{D})$ associated to $M \setminus \boldsymbol{D}$, as follows: we consider the fibre bundle $\mathcal{G}(M \setminus \boldsymbol{D})$, with the associated fibration

$$\mathcal{G}_p(M \setminus \boldsymbol{D}) \hookrightarrow \mathcal{G}(M \setminus \boldsymbol{D}) \to M \setminus \boldsymbol{D}.$$

Taking the abelianization of the associated exact sequence of homotopy groups, we obtain an exact sequence

$$H_1\left(\mathcal{G}_p(M \setminus \boldsymbol{D})\right) \to H_1\left(\mathcal{G}(M \setminus \boldsymbol{D})\right) \to H_1(M \setminus \boldsymbol{D}).$$

We observe that $\mathcal{G}_p(M \setminus \boldsymbol{D})$ is the Lagrangian Grassmannian of the symplectic vector space T_pM, so $H_1(\mathcal{G}_p(M \setminus \boldsymbol{D})) \cong \mathbb{Z}$ (see [19]). Thus we can define the grading datum $\boldsymbol{G}(M,\boldsymbol{D})$ to be given by the first morphism in this exact sequence. We show that the affine Fukaya category is naturally $\boldsymbol{G}(M,\boldsymbol{D})$-graded. Observe that the second map in the exact sequence gives the $H_1(M \setminus \boldsymbol{D})$-grading mentioned earlier.

Next, we introduce the relative Fukaya category, denoted $\mathcal{F}(M,\boldsymbol{D})$ (following [1, 2]). Its objects are exactly the same as those of $\mathcal{F}(M \setminus \boldsymbol{D})$: compact, exact, embedded, anchored Lagrangian branes in $M \setminus \boldsymbol{D}$. It is defined over the coefficient ring

$$R(M,\boldsymbol{D}) := \mathbb{C}[[r_1, \ldots, r_k]],$$

a power series ring with one generator for each divisor. We will often write R instead of $R(M,\boldsymbol{D})$ when no confusion is possible. We define morphism spaces by

$$CF^*_{\mathcal{F}(M,\boldsymbol{D})}(L_0, L_1) := \bigoplus_{p \in L_0 \cap L_1} R \cdot p.$$

The A_∞ structure maps μ^s count rigid boundary-punctured holomorphic disks in M. Namely, the coefficient of p_0 in $\mu^s(p_s, \ldots, p_1)$ is given by a signed count of rigid holomorphic disks

$$u : \mathbb{D} \setminus \{\zeta_0, \ldots, \zeta_s\} \to M$$

with boundary and asymptotic conditions as before. Each such disk u contributes a term

$$r^{u \cdot \boldsymbol{D}} := r_1^{u \cdot D_1} \ldots r_k^{u \cdot D_k} \in R,$$

where $u \cdot D_j$ denotes the topological intersection number of u with D_j. We observe that $u \cdot D_j \geq 0$ by positivity of intersection, so indeed the coefficients lie in R.

$\mathcal{F}(M,\boldsymbol{D})$ is still a $\boldsymbol{G}(M,\boldsymbol{D})$-graded category, but we remark that the coefficient ring R must have a non-trivial grading for this to be true. For example, consider the $H_1(M \setminus \boldsymbol{D})$-grading that we mentioned earlier. It is no longer true that a holomorphic disk u contributing to an A_∞ product μ^s has nullhomologous boundary in $M \setminus \boldsymbol{D}$, because the disk now maps into M, not $M \setminus \boldsymbol{D}$ as it did for the affine Fukaya category. However, if we remove small balls surrounding each intersection point of u with a divisor D_j, then the resulting surface defines a homology in $M \setminus \boldsymbol{D}$ between the boundary of u and a collection of meridian loops around the divisors. Thus, if we define the $H_1(M \setminus \boldsymbol{D})$-grading of the generator $r_j \in R$ to be the class of a meridian loop around divisor D_j, then the A_∞ structure respects the H_1-grading. Of course there remains more to check to show that $\mathcal{F}(M,\boldsymbol{D})$ is $\boldsymbol{G}(M,\boldsymbol{D})$-graded, but this is the basic idea – the details can be found in Section 5.

We observe that the zeroth-order part of $\mathcal{F}(M,\boldsymbol{D})$ defines an A_∞ category over $\mathbb{C}$, in which a holomorphic disk $u:\mathbb{D}\to M$ contributes to μ^s only if $u\cdot D_j=0$ for all j. This corresponds to counting only holomorphic disks which avoid the divisors $\boldsymbol{D}$, i.e., which lie in $M\setminus\boldsymbol{D}$. By definition, this corresponds to the affine Fukaya category $\mathcal{F}(M\setminus\boldsymbol{D})$. We therefore say that $\mathcal{F}(M,\boldsymbol{D})$ is a **$\boldsymbol{G}(M,\boldsymbol{D})$-graded deformation** of $\mathcal{F}(M\setminus\boldsymbol{D})$ over $R(M,\boldsymbol{D})$.

Finally, we recall the definition of the full Fukaya category $\mathcal{F}(M)$, as in [20, 5], where M is a Calabi-Yau symplectic manifold. Its objects are graded spin Lagrangians $L\subset M$. It is linear over the Novikov field Λ. Its morphism spaces are $\mathbb{Z}$-graded Λ-vector spaces (where Λ has degree $0\in\mathbb{Z}$), defined by

$$CF^*_{\mathcal{F}(M)}(L_0,L_1):=\bigoplus_{p\in L_0\cap L_1}\Lambda\cdot p.$$

The A_∞ structure maps are defined by signed counts of rigid boundary-punctured holomorphic disks $u:\mathbb{D}\to M$ as before. Each such disk u contributes a term $r^{\omega(u)}\in\Lambda$ to the corresponding coefficient of μ^s.

If we define the ring homomorphism

$$\begin{array}{rcl} R & \to & \Lambda \\ r_j & \mapsto & r \text{ for all } j, \end{array}$$

then Λ becomes an R-algebra. We expect that there is a fully faithful embedding of $\mathbb{Z}$-graded Λ-linear A_∞ categories,

$$\mathcal{F}(M,\boldsymbol{D})\otimes_R\Lambda\to\mathcal{F}(M).$$

We do not rigorously establish that such an embedding exists, so we state this as Assumption 8.1.1, and give some justification in Remark 8.1.2.

We summarize the properties of the three versions of the Fukaya category in the following table:

Notation	Affine: $\mathcal{F}(M\setminus\boldsymbol{D})$	Relative: $\mathcal{F}(M,\boldsymbol{D})$	Full: $\mathcal{F}(M)$
Objects	Eq. branes $L^{\#}$ in $M\setminus\boldsymbol{D}$	Same as for $\mathcal{F}(M\setminus\boldsymbol{D})$	Graded, Spin $L\subset M$
Coefficients	$\mathbb{C}$	$R(M,\boldsymbol{D})$	Λ
Morphisms	$\mathbb{C}\langle L_0\cap L_1\rangle$	$R\langle L_0\cap L_1\rangle$	$\Lambda\langle L_0\cap L_1\rangle$
Grading	$\boldsymbol{G}(M,\boldsymbol{D})$-graded	$\boldsymbol{G}(M,\boldsymbol{D})$-graded	$\mathbb{Z}$-graded (if M is C-Y)
A_∞ maps μ^s	$\#\{u:\mathbb{D}\to M\setminus\boldsymbol{D}\text{ hol.}\}$	$\#\{u:\mathbb{D}\to M\text{ hol.}\}$, with $r^{u\cdot\boldsymbol{D}}\in R$	$\#\{u:\mathbb{D}\to M\text{ hol.}\}$, with $r^{\omega(u)}\in\Lambda$

$$\mathcal{F}(M\setminus\boldsymbol{D})\xrightarrow{\boldsymbol{G}(M,\boldsymbol{D})\text{-graded deformation}}\mathcal{F}(M,\boldsymbol{D})\xrightarrow{\otimes_R\Lambda}\mathcal{F}(M).$$

1.4. The B-model mirror to the affine, relative and full Fukaya categories.

We define the smooth Calabi-Yau Fermat hypersurface

$$M^n:=\left\{\sum_{j=1}^n z_j^n=0\right\}\subset\mathbb{CP}^{n-1},$$

with ample divisors $D_j := \{z_j = 0\}$ for $j = 1, \ldots, n$. In this section, we will introduce the B-models which ought to be mirror to $\mathcal{F}(M^n \setminus \boldsymbol{D})$, $\mathcal{F}(M^n, \boldsymbol{D})$, and $\mathcal{F}(M^n)$.

We define the power series ring

$$R := \mathbb{C}[[r_1, \ldots, r_n]],$$

which is the coefficient ring of $\mathcal{F}(M^n, \boldsymbol{D})$. We define the R-algebra

$$S := R[u_1, \ldots, u_n],$$

and equip it with the $\mathbb{Z}$-grading so that R is concentrated in degree 0, and each u_j has degree 1. We define the element

$$w = u_1 \ldots u_n + \sum_{j=1}^{n} r_j u_j^n \in S,$$

of degree n.

Now note that $\mathrm{Proj}(S) = \mathbb{P}^{n-1}_R$. We consider the variety

$$\widetilde{N}^n := \{w = 0\} \subset \mathbb{P}^{n-1}_R,$$

and equip it with the action of Γ^*_n, exactly as we did for $\widetilde{N}^n_{nov}$ in Definition 1.2.4, then define

$$N^n := \widetilde{N}^n / \Gamma^*_n.$$

We note that the algebraic torus

$$\mathbb{T} := \{(\lambda_1, \ldots, \lambda_n) \in (\mathbb{C}^*)^n : \lambda_1 \ldots \lambda_n = 1\}$$

acts on S, by sending

$$\begin{array}{rcl} u_j & \mapsto & \lambda_j u_j, \\ r_j & \mapsto & \lambda_j^{-n} r_j. \end{array}$$

This action preserves w, and commutes with the action of Γ^*_n, and therefore defines an action of $\mathbb{T}$ on N^n.

We note that there are homomorphisms

$$\mathbb{C} \leftarrow R \rightarrow \Lambda,$$

given by

$$0 \leftarrow\!\shortmid r_j \mapsto r$$

for all j. Hence, by base change, we obtain

$$\begin{tikzcd} N^n_0 \arrow[r] \arrow[d] & N^n \arrow[d] & N^n_{nov} \arrow[l] \arrow[d] \\ \mathrm{Spec}(\mathbb{C}) \arrow[r] & \mathrm{Spec}(R) & \mathrm{Spec}(\Lambda). \arrow[l] \end{tikzcd}$$

We call N^n the **total space** of the family over $\mathrm{Spec}(R)$, we call N^n_0 the **special fibre**, and N^n_{nov} the **generic fibre**. We observe that this N^n_{nov} coincides with the definition given in Definition 1.2.4.

We expect mirror relationships as follows:

Coefficients	$\mathbb{C}$	R	Λ
B-model	$\mathrm{Perf}(N_0^n)$	$\mathrm{Perf}(N^n)$	$Coh(N^n_{nov})$
A-model	$\mathcal{F}(M^n \setminus \boldsymbol{D})$	$\mathcal{F}(M^n, \boldsymbol{D})$	$\mathcal{F}(M^n)$

(however note that we do **not** necessarily claim to prove all of these equivalences – this table is included to assist the reader in seeing the 'big picture').

Note that the map from R to $\mathbb{C}$ is naturally $\mathbb{T}$-equivariant. Thus, we can define $\mathbb{T}$-equivariant sheaves on N_0^n and N^n. However it is impossible to equip Λ with a $\mathbb{T}$-action so that the map $R \to \Lambda$ is $\mathbb{T}$-equivariant, so we can not talk about $\mathbb{T}$-equivariant sheaves on N^n_{nov}.

We expect that $\mathbb{T}$-equivariant sheaves correspond to anchored Lagrangian branes. Namely, we have seen that the morphism spaces between anchored Lagrangian branes admit a grading in the group $H_1(M^n \setminus \boldsymbol{D})$, and hence an action of its character group. In this case, there is a natural isomorphism

$$\mathrm{Hom}\,(H_1(M^n \setminus \boldsymbol{D}), \mathbb{C}^*) \cong \mathbb{T},$$

and we expect the mirror correspondences in the above table to be $\mathbb{T}$-equivariant (excluding the last column). In fact, we expect something stronger: they should be equivalences of $\boldsymbol{G}$-graded categories.

In [16, Theorem 7.4], we proved that there is a fully faithful, $\mathbb{T}$-equivariant embedding

$$\mathrm{Perf}(N_0^n) \hookrightarrow D^b\mathcal{F}(M^n \setminus \boldsymbol{D}).$$

In this paper, we extend this to prove results about the other columns.

Our ultimate aim is to understand the final column, and give a proof of Theorem 1. Thus we need a method of making computations in $D^bCoh(N^n_{nov})$. For this purpose, we use the category of **graded matrix factorizations**. Let us denote

$$S_{nov} := S \otimes_R \Lambda = \Lambda[u_1, \ldots, u_n],$$

with

$$w_{nov} := w \otimes 1 \in S_{nov}.$$

Then S_{nov} is $\mathbb{Z}$-graded, and w_{nov} is homogeneous of degree n. In [21], Orlov introduced the differential graded category of graded matrix factorizations of a homogeneous superpotential, $\mathrm{GrMF}(S_{nov}, w_{nov})$, and proved that there is an equivalence

$$\mathrm{Ho}\,(\mathrm{GrMF}(S_{nov}, w_{nov})) \cong D^bCoh\left(\widetilde{N}^n_{nov}\right),$$

where

$$\widetilde{N}^n_{nov} := \{w_{nov} = 0\} \subset \mathbb{P}^{n-1}_{\Lambda}$$

(see [21, Theorem 3.11]). Similarly, there is an equivalence of Γ_n^*-equivariant categories,

$$\mathrm{Ho}\left(\mathrm{GrMF}(S_{nov}, w_{nov})^{\Gamma_n^*}\right) \cong D^bCoh\left(N^n_{nov}\right).$$

Orlov's theorem applies because we work over the field Λ, and $\widetilde{N}^n_{nov} = \{w_{nov} = 0\}$ is smooth and Calabi-Yau. However, recall that by passing from the variety N^n, defined over R, to the variety N^n_{nov}, defined over Λ, we lose the $\mathbb{T}$-action. This is a disadvantage, because

$\mathbb{T}$-equivariance constrains the algebraic structures we consider significantly, and makes our classification problems tractable.

Therefore, we introduce (in Section 7) the category $\mathrm{GrMF}(S,w)$ of graded matrix factorizations of $w \in S$, over the coefficient ring R (Orlov mainly considered graded matrix factorizations over a field in [21], but the definition works over any ring). The coefficient ring R and the graded R-algebra S admit an action of $\mathbb{T}$, which preserves w. Therefore we can talk about $\mathbb{T}$-equivariant objects in $\mathrm{GrMF}(S,w)$, and furthermore there is a fully faithful embedding

$$\mathrm{GrMF}(S,w) \otimes_R \Lambda \hookrightarrow \mathrm{GrMF}(S_{nov}, w_{nov}).$$

However, because R is not a field, Orlov's theorem does **not** apply to give a relationship between the categories $\mathrm{GrMF}(S,w)$ and $\mathrm{Perf}(\widetilde{N}^n)$ (although it seems reasonable to hope that some sort of relationship might hold).

In fact, we first introduce a differential $\boldsymbol{G}$-graded category $MF^{\boldsymbol{G}}(S,w)$ of $\boldsymbol{G}$-graded matrix factorizations of $w \in S$ (these combine the $\mathbb{T}$-action with the $\mathbb{Z}$-grading, in the same way that anchored Lagrangian branes combine the $H_1(M^n \setminus \boldsymbol{D})$-grading with the $\mathbb{Z}$-grading in the Fukaya category). We show that $\mathrm{GrMF}(S,w)^{\Gamma_n^*}$ is some 'orbifolding' of it.

The starting point for this is the observation that $\mathrm{GrMF}(S,w)$ is a $\mathbb{Z}_n$-equivariant version of $MF^{\boldsymbol{G}}(S,w)$ (compare [22, 23]). In fact there is an action of $\tilde{\Gamma}_n^*$ on $MF^{\boldsymbol{G}}(S,w)$, and we show that there is a fully faithful embedding

$$MF^{\boldsymbol{G}}(S,w)^{\tilde{\Gamma}_n^*} \hookrightarrow \mathrm{GrMF}(S,w)^{\Gamma_n^*}$$

(recall that $\tilde{\Gamma}_n^*$ is an extension of $\mathbb{Z}_n$ by Γ_n^*; the $\mathbb{Z}_n$ got eaten up turning MF into GrMF). In fact, our notation for the $\tilde{\Gamma}_n^*$-equivariant category is different – we will write it as

$$\boldsymbol{p}_1^* MF^{\boldsymbol{G}}(S,w) \equiv MF^{\boldsymbol{G}}(S,w)^{\tilde{\Gamma}_n^*}.$$

We will not give the precise meaning of '$\boldsymbol{p}_1^*$' in this introduction, but will continue to use it for consistency with our later notation.

We consider the object $\mathcal{O}_0$ of $MF^{\boldsymbol{G}}(S,w)$, corresponding to the ideal $(u_1,\ldots,u_n) \subset S$. We denote its endomorphism algebra by $\mathcal{B}$. It is a deformation of the exterior algebra

$$\mathrm{Ext}^*_{Coh(\mathbb{C}^n)}(\mathcal{O}_0,\mathcal{O}_0) \cong \Lambda^*\mathbb{C}^n$$

over the power series ring R. Deformations of $\Lambda^*\mathbb{C}^n$ are governed by the Hochschild cohomology, which is given by polyvector fields, by the Hochschild-Kostant-Rosenberg isomorphism:

$$HH^*(\Lambda^*\mathbb{C}^n) \cong \mathbb{C}[[u_1,\ldots,u_n]][\theta_1,\ldots,\theta_n],$$

where the variables u_j commute and the variables θ_j anti-commute. We construct a minimal A_∞ model for this A_∞ deformation of $\Lambda^*\mathbb{C}^n$, and prove that its deformation classes are given exactly by the coefficients of w (following [24]). We prove a classification theorem (Theorem 3), which shows that these deformation classes, together with the $\boldsymbol{G}$-grading, are enough to determine the deformation up to A_∞ quasi-isomorphism and formal change of variables.

We then consider the full subcategory of $\boldsymbol{p}_1^* MF^{\boldsymbol{G}}(S,w)$ whose objects are the equivariant twists of $\mathcal{O}_0$. We denote it by $\widetilde{\mathcal{B}}$. It can be determined completely from $\mathcal{B}$. We also introduce a full subcategory

$$\widetilde{\mathcal{A}} \subset \mathcal{F}(M^n, \boldsymbol{D})$$

(the remaining sections of this introduction will consist of an explanation of how to compute $\widetilde{\mathcal{A}}$). We prove the following generalization of [16, Theorem 7.4]:

Theorem 2. *There exists a* $\psi \in \mathbb{C}[[T]] \subset R$*, where* $T = r_1 \dots r_n$*, with* $\psi(0) = \pm 1$*, and a quasi-isomorphism of* $\boldsymbol{G}$*-graded* R*-linear* A_∞ *categories*

$$\psi \cdot \widetilde{\mathcal{B}} \cong \widetilde{\mathcal{A}}.$$

By our previous discussion, there is a fully faithful embedding

$$\mathrm{Ho}\left(\boldsymbol{p}_1^* MF^{\boldsymbol{G}}(S, w) \otimes_R \Lambda\right) \hookrightarrow D^b Coh(N^n_{nov}) \cong D^b Coh^{\Gamma^*_n}\left(\widetilde{N}^n_{nov}\right).$$

The images of the equivariant twists of $\mathcal{O}_0$ correspond to equivariant twists of the restrictions of the Beilinson exceptional collection $\Omega^j(j)$ restricted to $\widetilde{N}^n_{nov}$ (for $j = 0, 1, \dots, n-1$) by characters of Γ^*_n.

The equivariant twists of the restrictions of the Beilinson exceptional collection generate $D^b Coh(N^n_{nov})$, so it follows immediately from Theorem 2, by tensoring with Λ, that there is an equivalence of triangulated categories

$$\psi \cdot D^b Coh(N^n_{nov}) \cong \mathrm{Ho}\left(Tw\left(\widetilde{\mathcal{A}} \otimes_R \Lambda\right)\right) \subset D^\pi \mathcal{F}(M^n)$$

(under our assumption (Assumption 8.1.1) that the second embedding above exists).

Finally, to complete the proof, we wish to show that $\widetilde{\mathcal{A}} \otimes_R \Lambda$ split-generates the Fukaya category. We do this by applying the split-generation result of [5], which says that, if the closed-open string map

$$\mathcal{CO} : QH^*(M^n) \to HH^*\left(\widetilde{\mathcal{A}} \otimes_R \Lambda\right)$$

is non-zero in the top degree $2(n-2)$, then $\widetilde{\mathcal{A}} \otimes_R \Lambda$ split-generates $D^\pi \mathcal{F}(M^n)$.

We observe that

$$\mathcal{CO}([\omega]) = r\frac{\partial \mu^*}{\partial r} \in HH^2\left(\widetilde{\mathcal{A}} \otimes_R \Lambda\right)$$

(in words, the image of the class of the symplectic form under the closed-open string map is the class in HH^2 corresponding to deforming the Fukaya category by scaling the symplectic form). We now observe that $\mathcal{CO}$ is a Λ-algebra homomorphism, so

$$\mathcal{CO}([\omega]^{n-2}) = \left(r\frac{\partial \mu^*}{\partial r}\right)^{n-2}.$$

We then compute that this class is non-zero in the Hochschild cohomology. It follows from the split-generation criterion that $\widetilde{\mathcal{A}} \otimes_R \Lambda$ split-generates $D^\pi \mathcal{F}(M^n)$. This completes the proof of Theorem 1.

In the rest of this introduction, we explain how we make computations in the relative Fukaya category $\mathcal{F}(M^n, \boldsymbol{D})$, which are sufficient to prove Theorem 2.

1.5. **Behaviour of the Fukaya category under branched covers.** Suppose that N and M are compact Kähler manifolds with ample normal-crossings divisors $\boldsymbol{E} \subset N$ and $\boldsymbol{D} \subset M$ as before, and that

$$\phi : (N, \boldsymbol{E}) \to (M, \boldsymbol{D})$$

is a branched cover ramified about the divisors $\boldsymbol{E}$, sending divisor E_j to divisor D_j, and with ramification of degree a_j about divisor E_j. We aim to understand how the affine and relative Fukaya categories of $(N, \boldsymbol{E})$ and $(M, \boldsymbol{D})$ are related.

First, we observe that the map

$$\phi : N \setminus \boldsymbol{E} \to M \setminus \boldsymbol{D}$$

is an unbranched cover. Therefore, any holomorphic disk in $M \setminus \boldsymbol{D}$ lifts to $N \setminus \boldsymbol{E}$, because it is contractible. It follows that the problem of relating $\mathcal{F}(N \setminus \boldsymbol{E})$ to $\mathcal{F}(M \setminus \boldsymbol{D})$ is essentially one of algebraic bookkeeping: we need to keep track of how the holomorphic disks lift, but do not need to compute any new moduli spaces of disks. This leads one to the statement that $\mathcal{F}(N \setminus \boldsymbol{E})$ is a 'semi-direct product of $\mathcal{F}(M \setminus \boldsymbol{D})$ with the character group of the covering group of ϕ' (see [2, Section 8b] and [4, Section 9]). We rephrase this in Section 3.4 using the language of $\boldsymbol{G}$-graded categories, in which we write

$$\mathcal{F}(N \setminus \boldsymbol{D}) \cong \boldsymbol{p}^* \mathcal{F}(M \setminus \boldsymbol{D})$$

(we won't explain this notation in the introduction).

Now we try to understand the behaviour of the relative Fukaya category with respect to branched covers. This is not as simple as the unramified cover case, because holomorphic disks may pass through the branching locus, and then they do not lift to the cover. In order to relate $\mathcal{F}(N, \boldsymbol{E})$ to $\mathcal{F}(M, \boldsymbol{D})$, we introduce a 'smooth orbifold relative Fukaya category' $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$, where $\boldsymbol{a} = (a_1, \ldots, a_k)$ denotes the degrees of ramification of the cover about the divisors (but we could define the orbifold Fukaya category for any tuple $\boldsymbol{a}$ of k positive integers).

The objects, morphism spaces and coefficient ring of $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$ are the same as for $\mathcal{F}(M, \boldsymbol{D})$. The A_∞ structure maps μ^s, however, count holomorphic disks $u : \mathbb{D} \to M$ that have ramification of degree a_j about divisor D_j wherever they intersect it. Each such disk contributes

$$r_1^{\#(\text{intersection points with } D_1)} \ldots r_k^{\#(\text{intersection points with } D_k)} \in R(M, \boldsymbol{D}).$$

In particular, if $\boldsymbol{a} = (1, 1, \ldots, 1)$ then we recover the relative Fukaya category. The category is still $\boldsymbol{G}(M, \boldsymbol{D})$-graded, but the coefficient ring $R(M, \boldsymbol{D})$ is equipped with a different $\boldsymbol{G}(M, \boldsymbol{D})$-grading depending on $\boldsymbol{a}$.

The holomorphic disks $u : \mathbb{D} \to M$ contributing to the orbifold relative Fukaya category, now do lift to holomorphic disks $u : \mathbb{D} \to N$ (by the homotopy lifting criterion). Thus, the relative Fukaya category $\mathcal{F}(N, \boldsymbol{E})$ is related to $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$ in exactly the same way that the affine Fukaya category $\mathcal{F}(N \setminus \boldsymbol{E})$ is related to $\mathcal{F}(M \setminus \boldsymbol{D})$: in the language of $\boldsymbol{G}$-graded categories,

$$\mathcal{F}(N, \boldsymbol{D}) \cong \boldsymbol{p}^* \mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a}).$$

It now remains to relate $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$ to $\mathcal{F}(M, \boldsymbol{D})$. In fact, we are only able to relate the 'first-order' parts of the categories (but this turns out to be enough for our purposes). The first-order relative Fukaya category is defined to be

$$\mathcal{F}(M,\boldsymbol{D})/\mathfrak{m}^2 := \mathcal{F}(M,\boldsymbol{D}) \otimes_R R/\mathfrak{m}^2,$$

where $\mathfrak{m} \subset R$ is the maximal ideal. It is linear over $R/\mathfrak{m}^2$. It retains only the information about rigid holomorphic disks $u : \mathbb{D} \to M$ passing through a **single** divisor D_j (with multiplicity 1).

Let us write

$$\mu^* = \mu_0^* + \mu_1^*$$

for the A_∞ structure maps μ^* in $\mathcal{F}(M,\boldsymbol{D})/\mathfrak{m}^2$, where μ_0^* gives the affine Fukaya category and μ_1^* gives the first-order terms. Then the A_∞ relations tell us that μ_1^* is a Hochschild cocycle, hence defines an element

$$\sum_{j=1}^{k} r_j \alpha_j \in HH^*(\mathcal{F}(M \setminus \boldsymbol{D})) \otimes \mathfrak{m}/\mathfrak{m}^2.$$

We call α_j the **first-order deformation classes** of $\mathcal{F}(M,\boldsymbol{D})$.

We prove (Theorem 4) that, if $\mathcal{F}(M,\boldsymbol{D})$ has first-order deformation classes α_j, then $\mathcal{F}(M,\boldsymbol{D},\boldsymbol{a})$ has first-order deformation classes $\alpha_j^{a_j}$, where the power is taken with respect to the Yoneda product on Hochschild cohomology. The proof looks very similar to the proof that the map $QH^*(M) \to HH^*(\mathcal{F}(M))$ is a ring homomorphism.

Remark 1.5.1. At first sight, this may seem a strange result: first-order deformation classes of a category live in HH^2, and the Yoneda product respects the $\mathbb{Z}$-grading, so one would expect the class $\alpha_j^{a_j}$ to no longer live in HH^2 and therefore not be an appropriate first-order deformation class. The solution lies in the fact that the coefficient rings R have non-trivial gradings, and in fact the coefficient rings for $\mathcal{F}(M,\boldsymbol{D})$ and $\mathcal{F}(M,\boldsymbol{D},\boldsymbol{a})$ have **different** gradings: thus, both $r_j\alpha_j$ and $r_j\alpha_j^{a_j}$ have degree 2 in the respective Hochschild cohomology groups in which they live.

Combining this result with our previous observations, if we have a branched cover $\phi : (N,\boldsymbol{E}) \to (M,\boldsymbol{D})$, then we can compute $\mathcal{F}(N,\boldsymbol{E})$ to first order if we know $\mathcal{F}(M,\boldsymbol{D})$ to first order.

1.6. The Fukaya category of M^n.

We will now explain how to compute the Fukaya category of M^n. We will keep the one-dimensional case ($n = 3$) as a running example throughout, despite the fact that we do not prove this case of Theorem 1 completely in this paper. We do this because one can see all of the holomorphic disks in the Fukaya category in this case, and gain intuition for the various versions of the Fukaya category that we introduce, and results that we prove about it.

We consider the Fermat hypersurfaces

$$M_a^n := \left\{ \sum_{j=1}^{n} z_j^a = 0 \right\} \subset \mathbb{CP}^{n-1}$$



FIGURE 1.6.1. $\mathbb{CP}^1$ with its real locus $\mathbb{RP}^1$ is shown in black, with the divisors $\boldsymbol{D}$ indicated by black dots. The Lagrangian immersion $L : S^1 \to \mathbb{CP}^1 \setminus \boldsymbol{D}$ is shown in red, and its self-intersection points are marked in blue.

with the smooth ample normal-crossings divisors $D_j = \{z_j = 0\}$ for $j = 1, \ldots, n$. There is a branched cover

$$\begin{aligned} \phi : (M^n_a, \boldsymbol{D}) &\to (M^n_1, \boldsymbol{D}), \\ [z_1 : \ldots : z_n] &\mapsto [z_1^a : \ldots : z_n^a]. \end{aligned}$$

Thus, applying the results described in the previous section, if we can compute $\mathcal{F}(M^n_1, \boldsymbol{D})$ to first order, then we can compute $\mathcal{F}(M^n_a, \boldsymbol{D})$ to first order.

We observe that $M^n_1 \cong \mathbb{CP}^{n-2}$, and $\boldsymbol{D}$ consists of n hyperplanes in general position. $M^n_1 \setminus \boldsymbol{D}$ is called the $(n-2)$-dimensional **pair of pants**. In Section 6, we construct a Lagrangian immersion

$$L : S^{n-2} \to M^n_1 \setminus \boldsymbol{D}$$

which has an anchored Lagrangian brane structure. This Lagrangian was introduced in [16].

Example 1.6.1. $M^3_1 = \mathbb{CP}^1$, and $\boldsymbol{D}$ consists of three divisors (points). The Lagrangian immersion $L : S^1 \to \mathbb{CP}^1 \setminus \boldsymbol{D}$ is shown in Figure 1.6.1. The A_∞ algebra $CF^*(L, L)$ was described in [4]. It was introduced as a $\mathbb{Z} \oplus H_1(M^3_1 \setminus \boldsymbol{D})$-graded category, but it is not hard to see the underlying $\boldsymbol{G}(M^3_1, \boldsymbol{D})$-graded structure.

The reason this Lagrangian is important is because it can be regarded as a 'fibre' in a Strominger-Yau-Zaslow fibration. See Figure 1.6.2 for the picture in the one-dimensional case. More generally, as shown in [25], the pair of pants $\mathbb{CP}^{n-2} \setminus \boldsymbol{D}$ is a singular torus fibration over the 'tropical amoeba of the pair of pants', which is some space stratified by affine manifolds. The torus fibration is non-singular over the top-dimensional faces of the tropical pair of pants.

We suggest that one should think, not of an SYZ fibration of the pair of pants over the tropical pair of pants, with some singular fibres, but rather of an SYZ **family** of objects of the Fukaya category, parametrized by the tropical pair of pants. The immersed Lagrangian sphere L is the object corresponding to the central point in the tropical pair of pants in this picture. The objects corresponding to points of the top-dimensional strata are Lagrangian torus fibres (recall that the fibration is non-singular there). The objects corresponding to points on the in-between strata are lower-dimensional incarnations of L, crossed with tori. We provided some evidence for this philosophy in [16], where we showed that the endomorphism algebra of L in $\mathcal{F}(\mathbb{CP}^{n-2} \setminus \boldsymbol{D})$ is quasi-isomorphic to the endomorphism algebra of the structure sheaf of the origin in the mirror category of matrix factorizations. We plan to make this picture more precise in future work.

In Section 6, we compute $CF^*(L,L)$ to first order in $\mathcal{F}(\mathbb{CP}^{n-2}, \boldsymbol{D})$, using a Morse-Bott model for the relative Fukaya category, based on the 'cluster homology' of [26]. We compute that the underlying vector space is an exterior algebra:

$$CF^*(L,L) \cong \Lambda^* \mathbb{C}^n \cong \mathbb{C}[\theta_1, \ldots, \theta_n],$$

where the variables θ_j anti-commute. For example, when $n = 3$, $CF^*(L,L)$ is generated by $H^*(S^1)$ (whose two generators we identify as the bottom and top classes 1 and $\theta_1 \wedge \theta_2 \wedge \theta_3$), together with two generators for each self-intersection point, which we label as in Figure 1.6.3.

We next show that the zeroth-order algebra structure, μ^2_0, coincides with the exterior algebra. In the case $n = 3$, the corresponding holomorphic triangles are shown in Figure 3(a). The shaded triangle can be viewed as having inputs θ_1 and θ_2 and output $\theta_1 \wedge \theta_2$, while the corresponding triangle on the back of the figure can be viewed as having inputs θ_2 and θ_1 and output $-\theta_1 \wedge \theta_2$. The other products follow similarly.

A_∞ structures with underlying cohomology algebra $\Lambda^* \mathbb{C}^n$ are classified by the Hochschild cohomology, which is given by polyvector fields, by the Hochschild-Kostant-Rosenberg isomorphism:

$$HH(\Lambda^* \mathbb{C}^n) \cong \mathbb{C}[[u_1, \ldots, u_n]][\theta_1, \ldots, \theta_n],$$

where variables u_i commute and θ_i anti-commute. We show that the endomorphism algebra $CF^*(L,L)$ in the affine Fukaya category $\mathcal{F}(\mathbb{CP}^{n-2} \setminus \boldsymbol{D})$ is completely determined, up to A_∞ quasi-isomorphism, by a single higher-order product, having the form

$$\mu^n(\theta_1, \ldots, \theta_n) = 1,$$

corresponding to the Hochschild cohomology class

$$u_1 \ldots u_n \in \mathbb{C}[[u_1, \ldots, u_n]][\theta_1, \ldots, \theta_n].$$

In the case $n = 3$, we can see the corresponding holomorphic disk in Figure 3(a). It is the shaded triangle, which we view as a degenerate 4-gon having inputs θ_1, θ_2, θ_3, and output a degenerate vertex on one of the sides of the triangle, corresponding to 1.



Figure 1.6.2. The Lagrangian $L : S^1 \to \mathbb{CP}^1 \setminus \boldsymbol{D}$ can also be drawn as a 'trefoil', shown here in red on the pair of pants (upper left). It lies over the central point (shown in red) in the tropical pair of pants (bottom). The mirror is the union of coordinate axes in $\mathbb{C}^3$ (upper right), and the object corresponding to L is the structure sheaf of the origin (shown in red).

We next compute that the endomorphism algebra of $CF^*(L,L)$ in the first-order relative Fukaya category $\mathcal{F}(\mathbb{CP}^{n-2}, \boldsymbol{D})/\mathfrak{m}^2)$ is determined by structure maps of the form

$$\mu^1(\theta_j) = r_j \cdot 1,$$

corresponding to first-order deformation classes

$$r_j u_j \in \mathbb{C}[[u_1, \ldots, u_n]][\theta_1, \ldots, \theta_n] \otimes \mathfrak{m}/\mathfrak{m}^2.$$

When $n = 3$, we can see the corresponding holomorphic disks in Figure 3(b). The shaded 'teardrop' shape has one input θ_1, and a degenerate output vertex corresponding to 1. It intersects divisor D_1 exactly once, and does not intersect the other divisors, hence contributes with a coefficient r_1. Thus it gives rise to the term $\mu^1(\theta_1) = r_1 \cdot 1$.


D_1
$\theta_2, \theta_3 \wedge \theta_1$
$\theta_3, \theta_1 \wedge \theta_2$
D_3
$\theta_1, \theta_2 \wedge \theta_3$
D_2
D_1
$\theta_2, \theta_3 \wedge \theta_1$
$\theta_3, \theta_1 \wedge \theta_2$
D_3
$\theta_1, \theta_2 \wedge \theta_3$
D_2

(a) Disks contributing to $CF^*(L,L)$ in the affine Fukaya category, $\mathcal{F}(\mathbb{CP}^1 \setminus \boldsymbol{D})$.

(b) Disks contributing to the first-order deformation class of $CF^*(L,L)$ in the relative Fukaya category, $\mathcal{F}(\mathbb{CP}^1, \boldsymbol{D})$.

FIGURE 1.6.3. Holomorphic disks contributing to $CF^*(L,L)$.

It follows from the result described in Section 1.5 that the first-order deformation classes of the A_∞ algebra

$$\mathcal{A} := CF^*_{\mathcal{F}(\mathbb{CP}^{n-2}, \boldsymbol{D}, (n,\ldots,n))}(L,L)$$

in the orbifold Fukaya category, are $r_j u_j^n$. Thus, the full deformation class of $\mathcal{A}$ is

$$u_1 \ldots u_n + \sum_{j=1}^{n} r_j u_j^n + \mathcal{O}(r^2),$$

which we observe coincides with the defining polynomial w of N^n, to first order.

We prove a classification theorem (Theorem 3) which shows that this is enough information to determine the full $\boldsymbol{G}$-graded deformation, up to A_∞ quasi-isomorphism and formal change of variables.

We show that the A_∞ algebra

$$\mathcal{B} := \mathrm{Hom}_{MF^{\boldsymbol{G}}(S,w)}(\mathcal{O}_0, \mathcal{O}_0)$$

also has the same underlying algebra, $\boldsymbol{G} \cong \boldsymbol{G}(\mathbb{CP}^{n-2}, \boldsymbol{D})$-grading, and deformation classes (recall the deformation classes were given exactly by w itself). Therefore, by the above-mentioned classification theorem, we have a formal change of variables ψ, and an A_∞ quasi-isomorphism

$$\mathcal{B} \cong \psi \cdot \mathcal{A}.$$

Now the algebraic procedures of passing from $\mathcal{F}(\mathbb{CP}^{n-2}, \boldsymbol{D}, (n,\ldots,n))$ to the branched cover $\mathcal{F}(M^n, \boldsymbol{D})$, and of passing from $MF^{\boldsymbol{G}}(S,w)$ to $\boldsymbol{p}_1^* MF^{\boldsymbol{G}}(S,w)$, are equivalent: we have

$$\widetilde{\mathcal{B}} \cong \boldsymbol{p}_1^* \mathcal{B} \cong \psi \cdot \boldsymbol{p}_1^* \mathcal{A} \cong \psi \cdot \widetilde{\mathcal{A}}.$$

This completes the proof of Theorem 2. See Figure 1.6.4 for a picture in the case $n = 3$.



Figure 1.6.4. A fundamental domain of the elliptic curve M^3, with the divisor $\boldsymbol{D}$ consisting of nine points, indicated by black dots. The 9-fold cover $(M^3, \boldsymbol{D}) \to (\mathbb{CP}^1, \boldsymbol{D})$ has ramification of order 3 about each divisor. The black lines are the pullback of $\mathbb{RP}^1$. The lifts of L are the curves shown in red, with their intersection points in blue. We have shaded two of the holomorphic disks contributing to the A_∞ products between lifts of L. Note that the triangle of Figure 3(a), which contributes to the affine Fukaya category, lifts directly to M^3, whereas the 'teardrop' illustrated in Figure 3(b), which contributes the term $r_1 u_1$ to the deformation class of the relative Fukaya category, does not lift, but rather gives rise to the 'clover-leaf' shape, which contributes the deformation class $r_3 u_3^3$.

2. Graded and equivariant categories

The main purpose of this section is to introduce the relevant notions of graded and equivariant algebraic objects, and modify the results in [2, Section 3] to classify such objects.

2.1. Grading data.

For the purposes of this section, we fix an integer $n \geq 3$.

Definition 2.1.1. An **unsigned grading datum** $\boldsymbol{G}$ is an abelian group Y together with a morphism $f : \mathbb{Z} \to Y$. We will use the shorthand $\boldsymbol{G} = \{\mathbb{Z} \xrightarrow{f} Y\}$. We will often write $\boldsymbol{G}$ as

$$\mathbb{Z} \xrightarrow{f} Y \xrightarrow{g} X \longrightarrow 0,$$

where X is the cokernel of f. We say that $\boldsymbol{G}$ is **exact** if the map $\mathbb{Z} \to Y$ is injective.

Definition 2.1.2. A **morphism** of unsigned grading data, $\boldsymbol{p} : \boldsymbol{G}_1 \to \boldsymbol{G}_2$, is a morphism $p : Y_1 \to Y_2$ that makes the following diagram commute:

$$\begin{tikzcd}
\mathbb{Z} \arrow[r, "f_1"] \arrow[d, equal] & Y_1 \arrow[d, "p"'] \\
\mathbb{Z} \arrow[r, "f_2"] & Y_2,
\end{tikzcd}$$

Composition of morphisms is defined in the obvious way, and this defines a category of unsigned grading data. We say that a morphism of unsigned grading data is **injective** (respectively **surjective**) if the map p is injective (respectively surjective). We will sometimes write $\boldsymbol{p}$ as

$$\begin{tikzcd}
\mathbb{Z} \arrow[r, "f_1"] \arrow[d, equal] & Y_1 \arrow[r, "g_1"] \arrow[d, "p"'] & X_1 \arrow[r] \arrow[d, "p_X"'] & 0 \\
\mathbb{Z} \arrow[r, "f_2"] & Y_2 \arrow[r, "g_2"] & X_2 \arrow[r] & 0,
\end{tikzcd}$$

where p_X is the map induced by p.

Definition 2.1.3. We define the **sign grading datum**, $\boldsymbol{G}_\sigma := \{\mathbb{Z} \to \mathbb{Z}_2\}$.

Definition 2.1.4. We define a **grading datum** $(\boldsymbol{G}, \boldsymbol{\sigma})$ to be an unsigned grading datum $\boldsymbol{G}$, together with a **sign morphism**, which is a morphism of unsigned grading data,

$$\boldsymbol{\sigma} : \boldsymbol{G} \to \boldsymbol{G}_\sigma.$$

We define a morphism of grading data to be a morphism of unsigned grading data that is compatible with sign morphisms. Henceforth, we will often omit the sign morphism $\boldsymbol{\sigma}$ from the notation to avoid clutter.

The sign morphism is important because it allows us to define certain signs in our algebraic objects, which allows us to work over fields of characteristic not equal to 2.

Example 2.1.5. We define the grading datum $\boldsymbol{G}_{\mathbb{Z}} := \{\mathbb{Z} \overset{\mathrm{id}}{\to} \mathbb{Z}\}$, with the obvious morphism $\boldsymbol{\sigma}$. It is an initial object in the category of grading data.

In practice, it is often simpler to work with objects called pseudo-grading data.

Definition 2.1.6. A **pseudo-grading datum** $\boldsymbol{H}$ is a morphism of abelian groups $f : Z \to Y$, together with an element $c \in \mathrm{Hom}(Z, \mathbb{Z})$, whose image lies inside $2\mathbb{Z} \subset \mathbb{Z}$.

Definition 2.1.7. A **morphism** of pseudo-grading data,

$$\boldsymbol{p} : \boldsymbol{H}_1 \to \boldsymbol{H}_2,$$

consists of maps p_Z and p_Y that make the following diagram commute:

$$\begin{tikzcd}
Z_1 \arrow[r, "f_1"] \arrow[d, "p_Z"'] & Y_1 \arrow[d, "p_Y"'] \\
Z_2 \arrow[r, "f_2"] & Y_2,
\end{tikzcd}$$

together with an element $d \in \mathrm{Hom}(Y_1, \mathbb{Z})$, whose image lies inside $2\mathbb{Z} \subset \mathbb{Z}$, such that

$$c_1 = p_Z^*(c_2) + f_1^*(d).$$

Definition 2.1.8. Given a pseudo-grading datum $\boldsymbol{H}$:

$$Z \xrightarrow{f} Y,$$

together with c, we define a grading datum $\boldsymbol{G}(\boldsymbol{H})$ by

$$\mathbb{Z} \longrightarrow (\mathbb{Z} \oplus Y)/Z,$$

where we define the map $Z \to \mathbb{Z} \oplus Y$ by

$$z \mapsto c(-z) \oplus f(z)$$

and the other maps in the obvious way. We define the sign morphism $\boldsymbol{\sigma} : \boldsymbol{G}(\boldsymbol{H}) \to \boldsymbol{G}_\sigma$ by

$$\begin{aligned} \sigma : (\mathbb{Z} \oplus Y)/Z &\to \mathbb{Z}_2, \\ \sigma(j \oplus y) &:= j. \end{aligned}$$

Observe that the condition that the image of c lies in $2\mathbb{Z}$ ensures that σ is well-defined.

Definition 2.1.9. Given a morphism of pseudo-grading data $\boldsymbol{p} : \boldsymbol{H}_1 \to \boldsymbol{H}_2$ as in Definition 2.1.7, we define a corresponding morphism of grading data

$$\boldsymbol{G}(\boldsymbol{p}) : \boldsymbol{G}(\boldsymbol{H}_1) \to \boldsymbol{G}(\boldsymbol{H}_2),$$

where

$$\boldsymbol{G}(\boldsymbol{p})(j \oplus y_1) := (j + d(y_1)) \oplus p(y_1).$$

It is not hard to check that this defines a functor from the category of pseudo-grading data to the category of grading data.

Example 2.1.10. If we denote by $\boldsymbol{0} := \{0 \to 0\}$ the zero morphism, then $\boldsymbol{G}(\boldsymbol{0}) = \boldsymbol{G}_{\mathbb{Z}}$.

Example 2.1.11. Given $n \geq 0$, we define the pseudo-grading datum $\boldsymbol{H}_{MF(n)} := \{\mathbb{Z} \overset{\times n}{\to} \mathbb{Z}\}$, with $c = 2$. We denote the corresponding grading datum by $\boldsymbol{G}_{MF(n)}$. It is exact, and has corresponding short exact sequence

$$\begin{tikzcd} 0 \arrow[r] & \mathbb{Z} \arrow[r, "f"] & (\mathbb{Z} \oplus \mathbb{Z})/(2, -n) \arrow[r, "g"] & \mathbb{Z}_n \arrow[r] & 0, \end{tikzcd}$$

This grading datum is important because it controls Orlov's 'category of graded matrix factorizations' of a superpotential of degree n (hence our terminology).

Now we will introduce another important grading datum, although first we introduce a bit of convenient notation.

Definition 2.1.12. We denote $[k] := \{1, \ldots, k\}$ for any positive integer k and, for any $K \subset [k]$,

$$y_K := \sum_{j \in K} y_j \in \mathbb{Z}\langle y_1, \ldots, y_k \rangle \cong \mathbb{Z}^k.$$

Example 2.1.13. Given $n \geq 1$, we denote by $\boldsymbol{H}^n_a$ the pseudo-grading datum

$$\begin{tikzcd} \mathbb{Z} \arrow[r, "y_{[n]}"] & \mathbb{Z}\langle y_1, \ldots, y_n \rangle, \end{tikzcd}$$

together with $c = 2(n-a)$. We denote by $\boldsymbol{G}^n_a$ the corresponding grading datum. This grading datum is important because it controls both the Fukaya category and the category of equivariant matrix factorizations that we will consider.

Now we prove a Lemma relating some of the grading data that we have introduced. We will use it to relate the category of matrix factorizations to the category of coherent sheaves.

Lemma 2.1.14. *For any $n \geq 1$, there is a commutative square of grading data:*

$$\begin{tikzcd} \boldsymbol{G}^n_n \arrow[r, two heads, "\boldsymbol{q}_1"] \arrow[d, hook, "\boldsymbol{p}_1"'] & \boldsymbol{G}_{\mathbb{Z}} \arrow[d, hook, "\boldsymbol{p}_2"'] \\ \boldsymbol{G}^n_1 \arrow[r, two heads, "\boldsymbol{q}_2"] & \boldsymbol{G}_{MF(n)}, \end{tikzcd}$$

such that $\boldsymbol{p}_1$ and $\boldsymbol{p}_2$ are injective.

Proof. The morphism $\boldsymbol{p}_2$ is uniquely defined because $\boldsymbol{G}_{\mathbb{Z}}$ is an initial object. It is clearly injective. The morphism $\boldsymbol{q}_1$ comes from the zero morphism of pseudo-grading data, with $d = 0$ (note this is a morphism of pseudo-grading data because $c = 0$ in $\boldsymbol{H}^n_n$).

The morphism $\boldsymbol{p}_1$ comes from the morphism of pseudo-grading data,

$$\begin{tikzcd}
\mathbb{Z} \arrow[r, "y_{[n]}"] \arrow[d, "\times n"'] & \mathbb{Z}^n \arrow[d, "\times n"'] \\
\mathbb{Z} \arrow[r, "y_{[n]}"] & \mathbb{Z}^n,
\end{tikzcd}$$

with $d = 2(1-n)y_{[n]}$ (where here we denote by $y_{[n]}$ the element of the dual space $(\mathbb{Z}^n)^\vee \cong \mathbb{Z}^n$). It is clearly injective.

The morphism $\boldsymbol{q}_2$ comes from the morphism of pseudo-grading data,

$$\begin{tikzcd}
\mathbb{Z} \arrow[r, "y_{[n]}"] \arrow[d, "\times(-1)"'] & \mathbb{Z}^n \arrow[d, "-y_{[n]}"'] \\
\mathbb{Z} \arrow[r, "\times n"] & \mathbb{Z},
\end{tikzcd}$$

with $d = 2y_{[n]}$.

It is a simple exercise, applying Definition 2.1.9, to check that the diagram commutes. □

2.2. **Graded vector spaces.** We recall that, if Y is an abelian group, then a **Y-graded vector space** is a vector space V, together with a collection of vector spaces V_y, indexed by $y \in Y$, and an isomorphism

$$V \cong \bigoplus_{y \in Y} V_y.$$

Definition 2.2.1. Let $p : Y_1 \to Y_2$ be morphism of abelian groups. Given a Y_1-graded vector space V, we define the Y_2-graded vector space $p_* V$, where

$$(p_* V)_b := \bigoplus_{p(a)=b} V_a.$$

In particular, the underlying vector space V does not change.

Definition 2.2.2. Let $p : Y_1 \to Y_2$ be morphism of abelian groups. Given a Y_2-graded vector space V, we define the Y_1-graded vector space $p^* V$, where

$$(p^* V)_a := V_{p(a)}.$$

Remark 2.2.3. If $p : Y_1 \to Y_2$ is injective, then $p^* V$ is just the part of V whose Y_2-degree lies in $\operatorname{im}(p)$.

Definition 2.2.4. Let $\boldsymbol{G} = \{\mathbb{Z} \to Y\}$ be a grading datum. A $\boldsymbol{G}$-graded vector space V is the same thing as a Y-graded vector space V.

In fact, for the purposes of this section, '$\boldsymbol{G}$-graded' is virtually identical to 'Y-graded'. Things will get more complicated in the subsequent sections.

Definition 2.2.5. Given an element $y \in Y$, and a $\boldsymbol{G}$-graded vector space V, we define $V[y]$ to be V with grading shifted by y.

Definition 2.2.6. Given a morphism $\boldsymbol{p}$ of grading data, we define operations $\boldsymbol{p}_*$ and $\boldsymbol{p}^*$ on $\boldsymbol{G}$-graded vector spaces to be identical to p_* and p^*.

Definition 2.2.7. If V is a $\boldsymbol{G}$-graded vector space, then it automatically becomes a $\mathbb{Z}_2$-graded vector space:

$$V \cong \boldsymbol{\sigma}_* V,$$

where $\boldsymbol{\sigma}$ is the sign morphism of $\boldsymbol{G}$. Given $v \in V$ of pure degree with respect to this $\mathbb{Z}_2$-grading, we denote its $\mathbb{Z}_2$-degree by $\sigma(v)$.

Definition 2.2.8. A $\boldsymbol{G}$-graded algebra is a Y-graded algebra, i.e., one whose multiplication respects the Y-grading.

Example 2.2.9. Recall the grading datum $\boldsymbol{G}_1^n$ of Example 2.1.13. We define the n-dimensional $\boldsymbol{G}_1^n$-graded vector space

$$U_n := \mathbb{C}\langle u_1, \ldots, u_n \rangle,$$

where we equip u_j with degree $(-1, y_j) \in (\mathbb{Z} \oplus Y_n)/(2(1-n) \oplus y_{[n]})$.

Example 2.2.10. Let a be an integer. We define the $\boldsymbol{G}_1^n$-graded vector space

$$V_a^n := \mathbb{C}\langle r_1, \ldots, r_n \rangle,$$

where we equip r_j with degree $(2-2a, a y_j) \in (\mathbb{Z} \oplus Y_n)/(2(1-n) \oplus y_{[n]})$.

Remark 2.2.11. We observe that, if $\boldsymbol{p}_1$ and $\boldsymbol{q}_1$ are the morphisms of grading data defined in Lemma 2.1.14, then

$$\boldsymbol{q}_{1*}\boldsymbol{p}_1^* V_n^n \cong V_n^n$$

is a $\mathbb{Z}$-graded vector space concentrated in degree 0.

We remark that, if V is a $\boldsymbol{G}$-graded vector space, then the exterior algebra $\Lambda(V)$ and symmetric algebra $\mathrm{Sym}(V)$ have natural $\boldsymbol{G}$-graded algebra structures.

Definition 2.2.12. We define the $\boldsymbol{G}_1^n$-graded exterior algebra

$$A_n := \Lambda(U_n) \cong \mathbb{C}[\theta_1, \ldots, \theta_n],$$

where the variable θ_j anti-commute.

Definition 2.2.13. We define the $\boldsymbol{G}_1^n$-graded power series ring

$$R_a^n := \mathbb{C}[[V_a^\vee]] \cong \mathbb{C}[[r_1, \ldots, r_n]].$$

The next Lemma looks abstruse, but will be used in Section 7.5 to relate equivariant matrix factorizations to equivariant coherent sheaves:

Lemma 2.2.14. *Suppose we are given a commutative diagram of exact morphisms of grading data:*

$$\begin{tikzcd}
\boldsymbol{G}_1 \arrow[r, "\boldsymbol{q}_1"] \arrow[d, hook, "\boldsymbol{p}_1"'] & \boldsymbol{G}_{\mathbb{Z}} \arrow[d, hook, "\boldsymbol{p}_2"'] \\
\boldsymbol{G} \arrow[r, "\boldsymbol{q}_2"] & \boldsymbol{G}_2,
\end{tikzcd}$$

where $\boldsymbol{p}_1$ and $\boldsymbol{p}_2$ are injective, and a $\boldsymbol{G}$-graded vector space V. Taking the X-part of the grading data and morphisms, we find there are morphisms

$$X_1 \xrightarrow{p_{1,X}} X \xrightarrow{q_{2,X}} X_2$$

whose composition is 0 *(by commutativity of the diagram). We define the group* Γ *to be the homology of this sequence:*

$$\Gamma := \ker(q_{2,X})/\mathrm{im}(p_{1,X}).$$

Then $\boldsymbol{p}_2^*\boldsymbol{q}_{2*}V$ admits a Γ-grading, and hence an action of the character group Γ^*, and there is an isomorphism

$$(\boldsymbol{p}_2^*\boldsymbol{q}_{2*}V)^{\Gamma^*} \cong \boldsymbol{q}_{1*}\boldsymbol{p}_1^*V$$

as $\mathbb{Z}$-graded vector spaces, where the superscript Γ^* denotes the Γ^*-invariant part (or equivalently the part of degree $0 \in \Gamma$).

Proof. We have, by definition, the degree-j parts

$$(\boldsymbol{p}_2^*\boldsymbol{q}_{2*}V)^j := \bigoplus_{q_2(y)=p_2(j)} A_y,$$

and

$$(\boldsymbol{q}_1^*\boldsymbol{p}_{1*}V)^j := \bigoplus_{q_1(y)=j} A_{p_1(y)}.$$

Note that the first of these is equal to the part of A whose Y-grading lies in $q_2^{-1}(\mathrm{im}(p_2))$, while the second is equal to the part of A whose Y-grading lies in $\mathrm{im}(p_1)$ (using the exactness of $\boldsymbol{p}_1$ and $\boldsymbol{p}_2$). By commutativity of the diagram,

$$\begin{aligned} y &= p_1(y_1) \\ \Rightarrow q_2(y) &= p_2(q_1(y_1)) \\ \Rightarrow y &\in q_2^{-1}(\mathrm{im}(q_1)), \end{aligned}$$

so $\mathrm{im}(p_1) \subset q_2^{-1}(\mathrm{im}(p_2))$. Therefore, we have

$$\boldsymbol{q}_1^*\boldsymbol{p}_{1*}V \subset \boldsymbol{p}_2^*\boldsymbol{q}_{2*}V.$$

Furthermore, the left-hand side is exactly equal to the part of the right-hand side whose Y-grading lies in

$$\mathrm{im}(p_1) \subset q_2^{-1}(\mathrm{im}(p_2)),$$

so we can equip the right-hand side with a grading in

$$q_2^{-1}(\mathrm{im}(p_2))/\mathrm{im}(p_1) \cong \Gamma,$$

and the left-hand side is equal to the part of degree $0 \in \Gamma$.

The fact that the $\mathbb{Z}$-gradings match up also follows from commutativity of the diagram:

$$q_1(y) = j \Rightarrow p_2(j) = q_2(p_1(y)).$$

This completes the proof. □

2.3. **$\boldsymbol{G}$-graded A_∞ algebras and Hochschild cohomology.** We now define appropriate notions of $\boldsymbol{G}$-graded A_∞ algebras and Hochschild cohomology. For the purposes of this section, let $\boldsymbol{G}$ be an exact grading datum.

Definition 2.3.1. Let R be a $\boldsymbol{G}$-graded algebra, and let A, B be $\boldsymbol{G}$-graded R-bimodules. For each $s \ge 0$, we define a $Y \oplus \mathbb{Z}$-graded R-bimodule, whose degree-(y,s) part is

$$CC^y_c(A,B|R)^s := \mathrm{Hom}_{R-bimod}(A^{\otimes_R s}, B)_{y-f(s)},$$

called **compactly supported Hochschild cochains of length s and degree y**. Note that the Y-grading is not quite the obvious one: if ϕ^s changes Y-degree by y', then we define the Y-grading of ϕ^s to be $y := f(s) + y'$. We will omit the '$|R$' from the notation unless it is necessary to avoid confusion. There is a natural filtration of $CC^*_c(A,B)$, called the **length filtration**, given by

$$(CC^*_c(A,B))^s := \bigoplus_{s' \ge s} (CC^*_c(A,B)^s).$$

We also define a filtered $\boldsymbol{G}$-graded R-bimodule, whose degree-y part is

$$CC^y(A,B) := \prod_{s \ge 0} CC^y(A,B)^s,$$

which is the $\boldsymbol{G}$-graded completion of $CC^*_c(A,B)$ with respect to the length filtration. Note that we call it a '$\boldsymbol{G}$-graded' completion because it is the completion in the category of $\boldsymbol{G}$-graded R-bimodules – this is different from taking the completion of $CC^*_c(A,B)$ in the category of R-bimodules, which would no longer be Y-graded (elements could be sums of non-zero elements of infinitely many different Y-degrees). If $B = A$, we denote

$$CC^*(A) := CC^*(A,A).$$

Given $\phi \in CC^*(A,B)$, we write ϕ^s for the length-s component of ϕ.

Definition 2.3.2. We consider the morphism $f : \mathbb{Z} \to Y$ coming from the grading datum $\boldsymbol{G}$, and define

$$\begin{array}{rcl} p : \mathbb{Z} \oplus \mathbb{Z} & \to & Y \oplus \mathbb{Z} \\ (j,k) & \mapsto & (f(j), j-k). \end{array}$$

Then we define the $\mathbb{Z} \oplus \mathbb{Z}$-graded R-bimodule

$$CC^*_{c,\boldsymbol{G}}(A,B) := p^* CC^*_c(A,B).$$

We denote the degree-$(s+t,t)$ part of $CC^*_{\boldsymbol{G}}(A,B)$ by

$$CC^{s+t}_{c,\boldsymbol{G}}(A,B)^t := \text{ the degree-}f(t)\text{ part of } CC^s(A,B).$$

We also define the $\mathbb{Z}$-graded R-bimodule

$$CC^*_{\boldsymbol{G}}(A,B) := f^* CC^*(A,B),$$

which is the completion of $CC^*_{c,\boldsymbol{G}}(A,B)$ with respect to the length filtration.

Remark 2.3.3. Note that $CC^*_{\boldsymbol{G}}(A,B)$ is the part of $CC^*(A,B)$ whose Y-grading lies in the image of $\mathbb{Z} \to Y$. In our applications, $CC^*_{\boldsymbol{G}}(A,B)$ will be easier to compute (being smaller), and most of our deformation theory problems take place inside it. It is this observation that makes many of our deformation theory problems tractable.

Definition 2.3.4. We also define a 'truncated' version of $CC^*_{\boldsymbol{G}}(A,B)$, by restricting to elements of non-positive degree t. Namely,

$$TCC^*_{\boldsymbol{G}}(A,B) := \prod_{s\geq 0}\bigoplus_{t\leq 0} CC^{s+t}_{\boldsymbol{G}}(A,B)^t.$$

Definition 2.3.5. Suppose that R is a $\boldsymbol{G}$-graded algebra, and A and B are $\boldsymbol{G}$-graded R-bimodules. The **Gerstenhaber product** is a map of degree $(f(-1),-1) \in Y \oplus \mathbb{Z}$,

$$\begin{array}{rcl} CC^*_c(A,B)\otimes_R CC^*_c(A) & \to & CC^*_c(A,B), \text{ which we denote by} \\ \phi\otimes\psi & \mapsto & \phi\circ\psi, \text{ and which is defined by} \\ \phi\circ\psi(a_n,\ldots,a_1) & := & \\ \displaystyle\sum_{i+j+k=n} & (-1)^{\dagger}\phi^{i+k+1}(a_{i+j+k},\ldots,a_{i+j+1},\psi^j(a_{i+j},\ldots,a_{i+1}),a_i,\ldots,a_1), & \end{array}$$

where

$$\dagger = (\sigma(\psi)+1)(\sigma(a_1)+\ldots+\sigma(a_i)-i)$$

(recalling Definition 2.2.7). If the left and right actions of R on the R-bimodules A and B coincide, then the Gerstenhaber product is R-bilinear; otherwise it is only R-linear in ϕ. Because the Gerstenhaber product respects the length filtration, it defines a product

$$CC^*(A,B)\otimes_R CC^*(A) \to CC^*(A,B)$$

of degree $f(-1)$, also called the Gerstenhaber product.

Definition 2.3.6. If R is a $\boldsymbol{G}$-graded algebra and A is a $\boldsymbol{G}$-graded R-bimodule, then we define the **Gerstenhaber bracket**, which is a Lie bracket of degree $f(-1)$ on $CC^*(A)$, by

$$[\phi,\psi] := \phi\circ\psi - (-1)^{(\sigma(\phi)+1)(\sigma(\psi)+1)}\psi\circ\phi.$$

Definition 2.3.7. If R is a $\boldsymbol{G}$-graded algebra, then a **$\boldsymbol{G}$-graded associative R-algebra** is a $\boldsymbol{G}$-graded R-bimodule A, together with an element

$$\mu^2 \in CC^2_{c,\boldsymbol{G}}(A)^0,$$

satisfying the associativity relation,

$$\mu^2\circ\mu^2 = 0.$$

Remark 2.3.8. If A is a $\boldsymbol{G}$-graded associative R-algebra, then the product

$$a_2\cdot a_1 := (-1)^{\sigma(a_1)}\mu^2(a_1,a_2)$$

is associative and respects the $\boldsymbol{G}$-grading, and makes A into a $\boldsymbol{G}$-graded associative R-algebra in the usual sense.

Remark 2.3.9. If R is semisimple:

$$R := \bigoplus_{i\in I}\mathbb{C}u_i, \;\; u_iu_j = \delta_{ij}u_i$$

(where I is finite) and A is unital, then this is equivalent to a category with objects indexed by I.

Definition 2.3.10. If (A,μ^2) is a $\boldsymbol{G}$-graded associative R-algebra, then we define the **Hochschild differential**

$$\begin{array}{rcl} \delta : CC^*(A) & \to & CC^*(A) \\ \delta(\tau) & := & \left[\mu^2,\tau\right]. \end{array}$$

It has degree $f(1) \in Y$ and increases length by 1 (i.e., it is induced by a similar differential of degree $(f(1), 1)$ on $CC^*_c(A)$). It follows from the fact that $\mu^2 \circ \mu^2 = 0$ that δ is a differential, i.e., $\delta^2 = 0$. We define the **Hochschild cohomology** of A to be its cohomology,

$$HH^*(A) := H^*(CC^*(A), \delta),$$

which is a $\boldsymbol{G}$-graded R-bimodule (as δ has pure degree in Y), and similarly

$$HH^*_{\boldsymbol{G}}(A) := H^*(CC^*_{\boldsymbol{G}}(A), \delta),$$

which is $\mathbb{Z}$-graded. Furthermore, because δ is pure of degree $(1, f(1))$ on $CC^*_c(A)$, we can also define compactly-supported versions, $HH^*_c(A)$ which is $Y \oplus \mathbb{Z}$-graded and $HH^*_{c,\boldsymbol{G}}(A)$ which is $\mathbb{Z} \oplus \mathbb{Z}$-graded. As before, we denote the degree-(j,k) part by $HH^j_{c,\boldsymbol{G}}(A)^k$.

Definition 2.3.11. If R is a $\boldsymbol{G}$-graded algebra, then a **$\boldsymbol{G}$-graded A_∞ algebra** over R is a $\boldsymbol{G}$-graded R-bimodule A, together with an element

$$\mu \in CC^2_{\boldsymbol{G}}(A),$$

satisfying $\mu^0 = 0$, and such that the A_∞ **associativity relation**

$$\mu \circ \mu = 0$$

is satisfied. We denote A_∞ algebras by $\mathcal{A} := (A, \mu^*)$. If $\mu^1 = 0$ (or equivalently, if μ sits inside $TCC^2_{\boldsymbol{G}}(A)$), we say that $\mathcal{A}$ is **minimal**. If we have μ^* such that $\mu^0 \neq 0$, but still $\mu \circ \mu = 0$, then (A, μ^*) is called a **curved** A_∞ algebra.

Remark 2.3.12. In other words, A is a Y-graded R-bimodule, equipped with R-multilinear maps

$$\mu^s : A^{\otimes_R s} \to A$$

of degree $f(2-s) \in Y$, for all $s \geq 1$, satisfying the A_∞ associativity relations.

Definition 2.3.13. If A and B are $\boldsymbol{G}$-graded R-bimodules, we define a new 'product'

$$\begin{aligned} CC^*(B) \otimes_R CC^*(A,B) &\to CC^*(A,B), \text{ denoted by} \\ \phi \otimes \psi &\mapsto \phi \diamond \psi, \text{ and which we define by} \\ (\phi \diamond \psi)^n(a_n, \ldots, a_1) &:= \end{aligned}$$
$$\sum_{i_1+\ldots+i_j=n} \phi^j(\psi^{i_1}(a_{i_1+\ldots+i_j}, \ldots, a_{1+i_2+\ldots+i_j}), \psi^{i_2}(a_{i_2+\ldots+i_j}, \ldots), \ldots, \psi^{i_j}(a_{i_j}, \ldots, a_1)).$$

It is R-linear only in the first variable ϕ. Note that $\diamond$ is clearly associative: $(F \diamond G) \diamond H = F \diamond (G \diamond H)$.

Remark 2.3.14. We remark that $\diamond$ descends to a product

$$CC^*_{\boldsymbol{G}}(B) \otimes CC^*_{\boldsymbol{G}}(A,B) \to CC^*_{\boldsymbol{G}}(A,B),$$

because the Y-degree of a summand of $\phi \diamond \psi$ clearly lies in the image of $f : \mathbb{Z} \to Y$. Although $\diamond$ does not have a pure grading, it is an easy exercise to check that it maps

$$CC^s_{\boldsymbol{G}}(B) \otimes CC^1_{\boldsymbol{G}}(A,B) \to CC^s_{\boldsymbol{G}}(A,B).$$

Definition 2.3.15. If $\mathcal{A} = (A, \mu)$ and $\mathcal{B} = (B, \eta)$ are $\boldsymbol{G}$-graded A_∞ algebras over R, then an A_∞ morphism from $\mathcal{A}$ to $\mathcal{B}$ is an element $F \in TCC^1_{\boldsymbol{G}}(A,B)$ such that

$$F \circ \mu - \eta \diamond F = 0 \in CC^2_{\boldsymbol{G}}(A,B)$$

(note that the class lives in degree 2 by Remark 2.3.14). Composition of two A_∞ morphisms is defined using the product $\diamond$. F is called **strict** if $F^j = 0$ for all $j \geq 2$.

If an A_∞ morphism induces an isomorphism on the level of cohomology, then it is said to be a **quasi-isomorphism**. In fact, when our A_∞ algebras are minimal, there is an easier notion, that of **formal diffeomorphism** (see [3, Section 1c]).

Definition 2.3.16. If R is a $\boldsymbol{G}$-graded $\mathbb{C}$-algebra, and A and B are $\boldsymbol{G}$-graded R-modules, then a **$\boldsymbol{G}$-graded formal diffeomorphism** from A to B is an element

$$F \in TCC^1_{\boldsymbol{G}}(A, B)$$

such that

$$F^1 : A \to B$$

is an isomorphism of R-modules.

Lemma 2.3.17. *If (A, μ) is a minimal $\boldsymbol{G}$-graded A_∞ algebra over R, and $F \in TCC^1_{\boldsymbol{G}}(A, B)$ is a formal diffeomorphism from A to B, then there exists a unique $\boldsymbol{G}$-graded minimal A_∞ structure $F_*\mu$ on B, such that F defines an A_∞ morphism from (A, μ) to $(B, F_*\mu)$.*

Proof. Briefly, $F_*\mu$ is determined inductively in the length filtration: if $F_*\mu$ is determined to length $\le s-1$, then at order s we have (schematically):

$$(F_*\mu)^s(F^1(a_s), F^1(a_{s-1}), \dots, F^1(a_1)) = \sum_{s' \le s-1} (F_*\mu)^{s'}(F(\dots), F(\dots), \dots, F(\dots)) + \sum_j \mu^j(\dots, F(\dots), \dots),$$

and since F^1 is an isomorphism, this determines F^s uniquely. One can quickly check that $F_*\mu$ satisfies the A_∞ equations. $\square$

Lemma 2.3.18. *Formal diffeomorphisms can be composed using $\diamond$: If F is a $\boldsymbol{G}$-graded formal diffeomorphism from A to B and G is a $\boldsymbol{G}$-graded formal diffeomorphism from B to C, then $G \diamond F$ is a $\boldsymbol{G}$-graded formal diffeomorphism from A to C. Furthermore, if μ is a $\boldsymbol{G}$-graded A_∞ structure on A, then*

$$(G \diamond F)_*\mu = G_*(F_*\mu).$$

Furthermore, $\diamond$ is associative.

Lemma 2.3.19. *$\boldsymbol{G}$-graded formal diffeomorphisms can be strictly inverted: if F is a $\boldsymbol{G}$-graded formal diffeomorphism from A to B, then there exists a unique $\boldsymbol{G}$-graded formal diffeomorphism G from B to A such that $G \diamond F = \mathrm{id}$ (where id denotes the formal diffeomorphism whose length-1 component is the identity, and all other components vanish).*

Proof. The construction of G is by induction on length, as in the proof of Lemma 2.3.17. $\square$

Definition 2.3.20. If A is a $\boldsymbol{G}$-graded vector space over $\mathbb{C}$, then there is a group

$$\mathfrak{G}(A) := \{F \in TCC^1_{\boldsymbol{G}}(A) : F^1 = \mathrm{id}\},$$

called $\boldsymbol{G}$-graded formal diffeomorphisms from A to itself. It follows from Lemmata 2.3.17, 2.3.18 and 2.3.19 that $\mathfrak{G}(A)$ is a group, and that it acts on the space of minimal $\boldsymbol{G}$-graded A_∞ structures on A. Note that this action preserves the underlying algebra (A, μ^2), and that F defines an A_∞ quasi-isomorphism from μ to $F_*\mu$.

Definition 2.3.21. If we are given a $\boldsymbol{G}$-graded associative algebra A over $\mathbb{C}$, we define $\mathfrak{A}(A)$, the set of $\boldsymbol{G}$-graded minimal A_∞ algebras $\mathcal{A} = (A, \mu)$ over $\mathbb{C}$, with μ^2 coinciding with the product on A.

Definition 2.3.22. Suppose that $\mathcal{A} = (A, \mu) \in \mathfrak{A}(A)$, and $\mu^s = 0$ for $2 < s < d$. Then the A_∞ associativity relations $\mu \circ \mu = 0$ imply that μ^d is a Hochschild cocycle for A. The class

$$\left[\mu^d\right] \in HH^2_{c,\boldsymbol{G}}(A)^{2-d}$$

is called the **order-d deforming class** of $\mathcal{A}$.

Remark 2.3.23. In [2] and [16], the class $[\mu^d]$ is called an order-d **deformation class**. However a large part of this paper is devoted to studying different objects (see Definition 2.4.8), also elements of Hochschild cohomology, and also called deformation classes in [2]. That is why, to avoid confusing the reader, and only for the purposes of the current paper, we use the terminology 'deforming class' to distinguish this object.

We recall a versality result from [3], appropriately modified to take into account $\boldsymbol{G}$-grading:

Proposition 2.3.24. *Suppose that A is a $\boldsymbol{G}$-graded associative algebra over $\mathbb{C}$, and there exists $d > 2$ such that*

$$HH^2_{c,\boldsymbol{G}}(A)^{2-s} \cong \begin{cases} \mathbb{C} & \text{for } s = d \\ 0 & \text{for all } s > 0,\ s \neq d. \end{cases}$$

Suppose that $\mathcal{A}_1 = (A, \mu_1)$ and $\mathcal{A}_2 = (A, \mu_2)$ both lie in $\mathfrak{A}(A)$, satisfy $\mu_1^s = \mu_2^s = 0$ for all $2 < s < d$, and have non-trivial order-d deforming class in $HH^2_{c,\boldsymbol{G}}(A)^{2-d}$. Then $\mathcal{A}_1$ and $\mathcal{A}_2$ are quasi-isomorphic.

Proof. The proof is by a straightforward order-by-order construction of a formal diffeomorphism F such that $F_* \mu_1 = \mu_2$, showing that all obstructions to the existence of F vanish (see also [2, Lemma 3.2]). □

Now we define Hochschild cohomology of an A_∞ algebra:

Definition 2.3.25. Suppose that $\mathcal{A} = (A, \mu)$ is a $\boldsymbol{G}$-graded A_∞ algebra. We define the Hochschild differential

$$\begin{aligned} \delta : CC^*(A) &\rightarrow CC^*(A), \\ \delta(\tau) &:= [\mu, \tau]. \end{aligned}$$

It follows from the fact that $\mu \circ \mu = 0$ that δ is a differential, i.e., $\delta^2 = 0$. We define the Hochschild cohomology of $\mathcal{A}$ to be the cohomology,

$$HH^*(\mathcal{A}) := H^*(CC^*(A), \delta).$$

The Hochschild differential has pure degree $f(1)$, so $HH^*(\mathcal{A})$ is $\boldsymbol{G}$-graded. However, note that the Hochschild differential is no longer pure with respect to the length, so we can not define the $Y \oplus \mathbb{Z}$-graded compactly-supported version, as we could for an associative algebra. However, δ does always increase (or preserve) the length; therefore, we do still have the length filtration on the Hochschild cochain complex.

Definition 2.3.26. Similarly, we can define the $\boldsymbol{G}$-graded Hochschild cohomology,

$$HH^*_{\boldsymbol{G}}(\mathcal{A}) := H^*(CC^*_{\boldsymbol{G}}(A), \delta).$$

It is $\mathbb{Z}$-graded and admits a length filtration.

Definition 2.3.27. If $\mathcal{A} = (A, \mu)$ is a $\boldsymbol{G}$-graded **minimal** A_∞ algebra (i.e., $\mu \in TCC^2_{\boldsymbol{G}}(A)$), then the Hochschild differential preserves the truncated Hochschild cochains. Thus it makes sense to define the **truncated Hochschild complex** $(TCC^*_{\boldsymbol{G}}(\mathcal{A}), \delta)$, and call its cohomology the **truncated Hochschild cohomology** $THH^*_{\boldsymbol{G}}(\mathcal{A})$.

Remark 2.3.28. Suppose that A is a $\boldsymbol{G}$-graded associative algebra and $\mathcal{A} \in \mathfrak{A}(A)$. The length filtration on the Hochschild cochain complex $CC^*(\mathcal{A})$ yields a '$\boldsymbol{G}$-graded spectral sequence' $(E_d^{*,*}, \delta_d^{*,*})$. The E_d page is $Y \oplus \mathbb{Z}$ graded, and δ_d has degree $(f(1), d)$. We have

$$E_2^* = HH_c^*(A).$$

When we need to prove that it converges to $HH^*(\mathcal{A})$, we will apply the 'complete convergence theorem' [27, Theorem 5.5.10]. The length filtration is clearly bounded above, because all Hochschild cochains have length $s \geq 0$, and hence it is also exhaustive. It is also complete in the category of $\boldsymbol{G}$-graded R-bimodules (because the Hochschild cochain complex is defined to be a direct product). Therefore, to prove that the spectral sequence converges to $HH^*(\mathcal{A})$, we must show that it is **regular**: for each $(y, s) \in Y \oplus \mathbb{Z}$, the differentials

$$\delta_d : E_d^{(y,s)} \to E_d^{(y+f(1),s+d)}$$

vanish for sufficiently large d. When we need to prove that the spectral sequence converges, we will in fact show that the differentials δ_d vanish whenever d is sufficiently large (independent of y, s).

Similarly, the length filtration on $CC^*_{\boldsymbol{G}}(\mathcal{A})$ induces a cohomological spectral sequence $(E_d^{*,*}, \delta_d^{*,*})$ with the standard grading convention, and

$$E_2^{s,t} = HH^{s+t}_{c,\boldsymbol{G}}(A)^t.$$

Remark 2.3.29. Suppose that A is a $\boldsymbol{G}$-graded associative algebra over $\mathbb{C}$, and $\mathcal{A} \in \mathfrak{A}(A)$ has $\mu^s = 0$ for $2 < s < d$, and order-d deforming class $[\mu^d]$. Then the first non-zero differential in the spectral sequence is δ_{d-1}, and it is given by

$$\begin{aligned} \delta_{d-1} : HH_c^{(y,s)}(A) &\to HH_c^{(y+f(1),s+d-1)}(A), \\ \delta_{d-1}(\phi) &= \left[\left[\mu^d\right], \phi\right], \end{aligned}$$

where we observe that the Gerstenhaber bracket $[-,-]$ descends to the cohomology.

Now we will define the appropriate notions of $\boldsymbol{G}$-graded A_∞ categories and their Hochschild cohomology.

Definition 2.3.30. Let $\boldsymbol{G} = \{\mathbb{Z} \to Y\}$ be a grading datum, and R a $\boldsymbol{G}$-graded algebra. A **$\boldsymbol{G}$-graded pre-category** $\mathcal{C}$ over R is a set of objects $Ob(\mathcal{C})$, together with morphism spaces

$$\mathrm{Hom}_{\mathcal{C}}(K, L)$$

which are $\boldsymbol{G}$-graded R-bimodules, and an action of Y on $Ob(\mathcal{C})$ by 'shifts' $[y]$, together with (compatible) isomorphisms

$$\mathrm{Hom}_{\mathcal{C}}(K[y_1], L[y_2]) \cong \mathrm{Hom}_{\mathcal{C}}(K, L)[y_2 - y_1].$$

We can think of this as equipping $\mathcal{C}$ with an $R[Y]$-bimodule structure. We define the group $CC^*(\mathcal{C})$, and the $\boldsymbol{G}$-graded version $CC^*_{\boldsymbol{G}}(\mathcal{C})$, by analogy with Definitions 2.3.1 and 2.3.2, restricting to the parts that respect the above isomorphisms. We can think of this strict equivariance requirement as taking $CC^*(\mathcal{C}|R[Y])$, where $\mathcal{C}$ acquires an $R[Y]$-linear structure from the Y-action. Explicitly, it means that for $\phi \in CC^s(\mathcal{C})$, and any $y_0, \ldots, y_s \in Y$, the following diagram commutes:

$$\begin{tikzcd}
\mathrm{Hom}(L_{s-1}, L_s) \otimes \ldots \otimes \mathrm{Hom}(L_0, L_1) \arrow[r, "\phi^s"] \arrow[d] & \mathrm{Hom}(L_0, L_s) \arrow[d] \\
\mathrm{Hom}(L_{s-1}[y_{s-1}], L_s[y_s]) \otimes \ldots \otimes \mathrm{Hom}(L_0[y_0], L_1[y_1])[y_s - y_0] \arrow[r, "\phi^s"] & \mathrm{Hom}(L_0[y_0], L_s[y_s])[y_s - y_0].
\end{tikzcd}$$

Definition 2.3.31. We define a ***G*-graded** A_∞ **category** $\mathcal{C}$ over R to be a $\boldsymbol{G}$-graded pre-category together with

$$\mu \in CC^2_{\boldsymbol{G}}(\mathcal{C})$$

satisfying $\mu^0 = 0$ and $\mu \circ \mu = 0$. An A_∞ category is said to be **cohomologically unital** if its cohomological category is unital. We define the Hochschild cohomology $HH^*(\mathcal{C})$ and $HH^*_{\boldsymbol{G}}(\mathcal{C})$ by analogy with Definition 2.3.25. We define $\boldsymbol{G}$-graded A_∞ functors by analogy with $\boldsymbol{G}$-graded A_∞ morphisms. We also consider the case where $\mu^0 \neq 0$; in this case we say that $\mathcal{C}$ is a **curved** A_∞ **category**.

Definition 2.3.32. A $\boldsymbol{G}$-graded A_∞ category is said to be **minimal** if μ lies in $TCC^2_{\boldsymbol{G}}(\mathcal{C})$.

Remark 2.3.33. The notions of unitality and equivalence for minimal A_∞ categories are simpler than for non-minimal categories. Because there is no differential μ^1 on the morphisms spaces, μ^2 is strictly associative and therefore defines a category. Thus, if $(\mathcal{C}, \mu)$ is minimal and cohomologically unital, then $(\mathcal{C}, \mu^2)$ is a category, in particular is unital. We say that two objects of $\mathcal{C}$ are **quasi-isomorphic** if they are quasi-isomorphic as objects of $(\mathcal{C}, \mu^2)$. We say that an A_∞ functor $\mathcal{F} : (\mathcal{C}, \mu) \to (\mathcal{D}, \eta)$ between minimal A_∞ categories is a **quasi-equivalence** if the functor $\mathcal{F}^1 : (\mathcal{C}, \mu^2) \to (\mathcal{D}, \eta^2)$ is a quasi-equivalence.

Lemma 2.3.34. *If $\mathcal{F} : (\mathcal{C}, \mu) \to (\mathcal{D}, \eta)$ is an A_∞ quasi-equivalence between minimal A_∞ categories, then there exists an A_∞ functor $\mathcal{G} : (\mathcal{D}, \eta) \to (\mathcal{C}, \mu)$ such that $\mathcal{F}^1$ and $\mathcal{G}^1$ are mutually inverse quasi-equivalences.*

Proof. Follows from Lemma 2.3.19. □

Definition 2.3.35. If $\mathcal{A}$ is a $\boldsymbol{G}$-graded A_∞ algebra, then we denote by $\underline{\mathcal{A}}$ the smallest $\boldsymbol{G}$-graded A_∞ category with an object whose endomorphism algebra is $\mathcal{A}$. Namely, $\underline{\mathcal{A}}$ has objects $K[y]$ indexed by Y, and morphism spaces

$$\mathrm{Hom}(K[y_1], K[y_2]) := \mathcal{A}[y_2 - y_1].$$

By definition, there are isomorphisms

$$CC^*(\mathcal{A}) \cong CC^*(\underline{\mathcal{A}})$$

and

$$CC^*_{\boldsymbol{G}}(\mathcal{A}) \cong CC^*_{\boldsymbol{G}}(\underline{\mathcal{A}}).$$

We define the A_∞ structure maps μ^* on $\underline{\mathcal{A}}$ to be the image of those on $\mathcal{A}$ under the latter isomorphism.

Now we explain how $\boldsymbol{G}$-graded A_∞ categories can be 'pulled back' along injective morphisms of grading data, and how this operation affects the Hochschild cohomology.

Definition 2.3.36. Let $\boldsymbol{p} : \boldsymbol{G}_1 \hookrightarrow \boldsymbol{G}_2$ be an injective morphism of grading data, R a $\boldsymbol{G}_1$-graded algebra, and $\mathcal{C}$ a $\boldsymbol{G}_2$-graded pre-category over $\boldsymbol{p}_* R$. We define $\boldsymbol{p}^*\mathcal{C}$, a $\boldsymbol{G}_1$-graded pre-category over R, to have the same objects as $\mathcal{C}$, but $\boldsymbol{G}_1$-graded morphism spaces

$$\mathrm{Hom}_{\boldsymbol{p}^*\mathcal{C}}(K,L) := \boldsymbol{p}^*\mathrm{Hom}_{\mathcal{C}}(K,L)$$

(thus it is a faithful but not full sub-pre-category). We note that $\boldsymbol{p}^*\mathcal{C}$ still has an action of Y_2 by shifts, so the subgroup $Y_1 \subset Y_2$ acts, equipping $\boldsymbol{p}^*\mathcal{C}$ with the structure of a $\boldsymbol{G}_1$-graded pre-category over R.

Because Y_2 acts on $\boldsymbol{p}^*\mathcal{C}$ by shifts (shifting all objects simultaneously by the same $y \in Y_2$), it acts on $CC^*(\boldsymbol{p}^*\mathcal{C})$. However the action of the subgroup Y_1 is trivial by definition, since we restrict to Hochschild cochains that respect the shifts by Y_1. So there is an action of the group $\Gamma := Y_2/Y_1$ on $CC^*(\boldsymbol{p}^*\mathcal{C})$. It is not hard to see that the Γ-fixed part is isomorphic to $\boldsymbol{p}^*CC^*(\mathcal{C})$. Thus, we have

$$\boldsymbol{p}^*CC^*(\mathcal{C}) \cong CC^*(\boldsymbol{p}^*\mathcal{C})^\Gamma,$$

and it follows that

$$CC^*_{\boldsymbol{G}_2}(\mathcal{C}) \cong CC^*_{\boldsymbol{G}_1}(\boldsymbol{p}^*\mathcal{C})^\Gamma \subset CC^*_{\boldsymbol{G}_1}(\boldsymbol{p}^*\mathcal{C}).$$

Thus we can make the following:

Definition 2.3.37. If $\mathcal{C}$ is a $\boldsymbol{G}_2$-graded A_∞ category over R, with structure maps μ^*, then we define $\boldsymbol{p}^*\mathcal{C}$, a $\boldsymbol{G}_1$-graded A_∞ category over R, whose A_∞ structure maps are given by the image of μ^* under the inclusion

$$CC^2_{\boldsymbol{G}_2}(\mathcal{C}) \cong CC^2_{\boldsymbol{G}_1}(\boldsymbol{p}^*\mathcal{C})^\Gamma \subset CC^2_{\boldsymbol{G}_1}(\boldsymbol{p}^*\mathcal{C}).$$

Remark 2.3.38. It follows that there are isomorphisms

$$\boldsymbol{p}^*HH^*(\mathcal{C}) \cong HH^*(\boldsymbol{p}^*\mathcal{C})^\Gamma$$

and

$$HH^*_{\boldsymbol{G}_2}(\mathcal{C}) \cong HH^*_{\boldsymbol{G}_1}(\boldsymbol{p}^*\mathcal{C})^\Gamma \subset HH^*_{\boldsymbol{G}_1}(\boldsymbol{p}^*\mathcal{C}).$$

Now we explain how a $\boldsymbol{G}$-graded A_∞ category can be 'pushed forward' along a surjective morphism of grading data.

Definition 2.3.39. Let $\boldsymbol{p} : \boldsymbol{G}_1 \to \boldsymbol{G}_2$ be a surjective morphism of grading data, and $\mathcal{C}$ a $\boldsymbol{G}_1$-graded A_∞ category over a $\boldsymbol{G}_1$-graded algebra R. We now define a $\boldsymbol{G}_2$-graded pre-category $\boldsymbol{p}_*\mathcal{C}$ over $\boldsymbol{p}_*R$, as follows: First, observe that there are canonical isomorphisms of $\boldsymbol{G}_2$-graded vector spaces

$$\boldsymbol{p}_*\mathrm{Hom}_{\mathcal{C}}(K[y],L) \cong \boldsymbol{p}_*\mathrm{Hom}_{\mathcal{C}}(K,L)$$

for any $y \in \Gamma$. We define the set of objects of $\boldsymbol{p}_*\mathcal{C}$ to be the quotient of the set of objects of $\mathcal{C}$ by the action of $\Gamma \subset Y_1$. We define the $\boldsymbol{G}_2$-graded morphism spaces to be

$$\mathrm{Hom}_{\boldsymbol{p}_*\mathcal{C}}(K,L) := \boldsymbol{p}_*\mathrm{Hom}_{\mathcal{C}}(K,L).$$

This is well-defined by our previous remarks. Furthermore, because $\boldsymbol{p}$ is surjective, there is an obvious action of Y_2 on $\boldsymbol{p}_*\mathcal{C}$ by shifts, so $\boldsymbol{p}_*\mathcal{C}$ is a $\boldsymbol{G}_2$-graded pre-category. In some sense we have

$$\boldsymbol{p}_*\mathcal{C} = \mathcal{C} \otimes_{R[Y_1]} R[Y_2].$$

Now it is not hard to see that $CC^y(\boldsymbol{p}_*\mathcal{C})$ is just the completion of $(\boldsymbol{p}_*CC^*(\mathcal{C}))^y$ with respect to the length filtration, for all $y \in Y_2$ (observe that the completion is only needed when $\boldsymbol{p}$ has an infinite kernel). It follows that there is an inclusion

$$CC^*_{\boldsymbol{G}_1}(\mathcal{C}) \hookrightarrow CC^*_{\boldsymbol{G}_2}(\boldsymbol{p}_*\mathcal{C}).$$

Definition 2.3.40. If $\mathcal{C}$ is a $\boldsymbol{G}_1$-graded A_∞ category over R, with structure maps μ^*, and $\boldsymbol{p} : \boldsymbol{G}_1 \to \boldsymbol{G}_2$ a surjective morphism of grading data, then we define $\boldsymbol{p}_*\mathcal{C}$, a $\boldsymbol{G}_2$-graded A_∞ category over $\boldsymbol{p}_*R$, whose A_∞ structure maps are given by the image of μ^* under the inclusion

$$CC^2_{\boldsymbol{G}_1}(\mathcal{C}) \hookrightarrow CC^2_{\boldsymbol{G}_2}(\boldsymbol{p}_*\mathcal{C}).$$

Remark 2.3.41. It follows that $HH^y(\boldsymbol{p}_*\mathcal{C})$ is the completion of $(\boldsymbol{p}_*HH^*(\mathcal{C}))^y$ with respect to the length filtration, for all $y \in Y_2$. Observe that the completion is only needed when $\boldsymbol{p}$ has infinite kernel and $HH^*(\mathcal{C})$ has infinite rank.

2.4. **Deformations of A_∞ algebras.**

For the purposes of this section, let us fix a grading datum $\boldsymbol{G}$, and a power series ring

$$R \cong \mathbb{C}[[r_1, \ldots, r_k]]$$

with a $\boldsymbol{G}$-grading.

We also equip R with its natural filtration by order. We denote the order-j part of R by R^j. There is a natural projection $R \to R^0 \cong \mathbb{C}$, given by setting all $r_j = 0$, and we denote the kernel of this projection by $\mathfrak{m} \subset R$.

We also denote by R_0 the part of R of degree $0 \in Y$.

Definition 2.4.1. Given $\psi \in R_0$, there is a $\boldsymbol{G}$-graded algebra homomorphism

$$\begin{aligned} f_\psi : R &\to R, \\ f_\psi(r^j) &:= \psi^j \cdot r^j, \text{ where } r^j \in R^j \text{ has order } j. \end{aligned}$$

We now define a group, which by abuse of notation we call $\mathrm{Aut}(R)$, by setting

$$\begin{aligned} \mathrm{Aut}(R) &:= \{\psi \in R_0 : \psi(0) \neq 0\}, \\ \phi \cdot \psi &:= f_\phi(\psi)\phi, \end{aligned}$$

and an action of $\mathrm{Aut}(R)$ on R by $\boldsymbol{G}$-graded algebra isomorphisms. Note that the condition $\psi(0) \neq 0$ ensures that ψ has a unique inverse in $\mathrm{Aut}(R)$, which can be constructed order-by-order.

Definition 2.4.2. If A is a $\boldsymbol{G}$-graded vector space, then $A \otimes R$ is a $\boldsymbol{G}$-graded R-bimodule, and we have an isomorphism

$$CC^*(A \otimes R, A \otimes R | R) \cong CC^*(A, A \otimes R | \mathbb{C}).$$

If we have

$$\begin{aligned} \phi &\in CC^*(A, A \otimes R), \\ \phi &= \sum_{j \geq 0} \phi_j, \text{ where} \\ \phi_j &\in CC^*(A, A \otimes R^j), \end{aligned}$$

then we call ϕ_j the **order-j component** of ϕ.

Definition 2.4.3. Let $\mathcal{A} = (A, \mu_0)$ be a $\boldsymbol{G}$-graded minimal A_∞ algebra over $\mathbb{C}$. A **$\boldsymbol{G}$-graded deformation of $\mathcal{A}$ over R** is an element

$$\mu \in CC^2_{\boldsymbol{G}}(A \otimes R | R) \cong CC^2_{\boldsymbol{G}}(A, A \otimes R | \mathbb{C})$$

that makes $A \otimes R$ into an A_∞ algebra over R (i.e., $\mu \circ \mu = 0$), and whose order-0 component is μ_0. If $\mu \in TCC^2_{\boldsymbol{G}}(A, A \otimes R)$, then the deformation is said to be **minimal**.

Remark 2.4.4. Observe that $CC^2_{\boldsymbol{G}}(A, A \otimes R)$ and $TCC^2_{\boldsymbol{G}}(A, A \otimes R)$ are R_0-modules in the obvious way.

Definition 2.4.5. The group $\mathrm{Aut}(R)$ acts on the set of $\boldsymbol{G}$-graded deformations of $\mathcal{A}$ over R, via its action on R. We write $\psi \cdot \mu$ for the action of $\psi \in \mathrm{Aut}(R)$ on a deformation μ.

Now suppose that (A, μ_0) and (B, η_0) are $\boldsymbol{G}$-graded A_∞ algebras over $\mathbb{C}$, and (A, μ) and (B, η) are $\boldsymbol{G}$-graded A_∞ deformations of these over R. We recall (from Definition 2.3.15) that a $\boldsymbol{G}$-graded A_∞ morphism from (A, μ) to (B, η) over R is an element

$$F \in TCC^1_{\boldsymbol{G}}(A \otimes R, B \otimes R | R) \cong TCC^1_{\boldsymbol{G}}(A, B \otimes R | \mathbb{C})$$

such that

$$F \circ \mu - \eta \diamond F = 0.$$

Once again, we write

$$F = \sum_{j \geq 0} F_j,$$

where F_j is the order-j component of F. Observe that F_0 is a $\boldsymbol{G}$-graded A_∞ morphism from (A, μ_0) to (B, η_0).

The notion of A_∞ morphisms over R is not very well-behaved (for example, it is not clear that A_∞ quasi-isomorphisms can be inverted over R). It turns out that it will be sufficient for us to work with **minimal** A_∞ algebras, and formal diffeomorphisms. We recall the notion of $\boldsymbol{G}$-graded formal diffeomorphisms from Definition 2.3.16, and make appropriate modifications for the case of A_∞ algebras defined over R:

Definition 2.4.6. If A and B are $\boldsymbol{G}$-graded vector spaces over $\mathbb{C}$, and R a $\boldsymbol{G}$-graded power series ring, then a **$\boldsymbol{G}$-graded formal diffeomorphism** from $A \otimes R$ to $B \otimes R$ is an element

$$F \in TCC^1_{\boldsymbol{G}}(A, B \otimes R)$$

such that

$$F^1_0 : A \to B$$

is an isomorphism of vector spaces (to clarify: this is the order-0 component of the length-1 part of F).

As before (see Lemmata 2.3.17, 2.3.18, 2.3.19), formal diffeomorphisms can be composed, used to push forward minimal $\boldsymbol{G}$-graded A_∞ structures over R, and they can be strictly inverted. This last point is particularly important, because (as we stated above), there is no reason for an arbitrary A_∞ quasi-isomorphism over R to be invertible.

Now we introduce an analogue of Definition 2.3.20 for minimal deformations of A_∞ algebras.

Definition 2.4.7. If $\mathcal{A} = (A, \mu_0)$ is a $\boldsymbol{G}$-graded minimal A_∞ algebra over $\mathbb{C}$, we consider the group of $\boldsymbol{G}$-graded formal diffeomorphisms from A to itself, whose leading-order term is the identity:

$$\mathfrak{G}_R(A) := \{F \in TCC^1_{\boldsymbol{G}}(A, A \otimes R) : F^1 = \mathrm{id} + \mathfrak{m}\}.$$

By the analogues of Lemmata 2.3.17, 2.3.18 and 2.3.19, $\mathfrak{G}_R(A)$ forms a group, and this group acts on the set of $\boldsymbol{G}$-graded minimal A_∞ structures on $A \otimes R$. The action of F on $\mu \in TCC^2_{\boldsymbol{G}}(A, A \otimes R)$ is denoted by $F_*\mu \in TCC^2_{\boldsymbol{G}}(A, A \otimes R)$, and it is the unique element such that F defines an A_∞ morphism from μ to $F_*\mu$.

Definition 2.4.8. Suppose that μ is a $\boldsymbol{G}$-graded deformation of the A_∞ algebra $\mathcal{A} = (A, \mu_0)$ over R. The first-order component of the A_∞ relation $\mu \circ \mu = 0$ tells us that

$$\mu_1 \in CC^2_{\boldsymbol{G}}\left(A, A \otimes R^1\right)$$

is a Hochschild cochain. Thus, we obtain an element

$$[\mu_1] \in HH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^1),$$

which we call the **first-order deformation class** of μ.

Definition 2.4.9. If μ is a $\boldsymbol{G}$-graded **minimal** deformation of the **minimal** A_∞ algebra $\mathcal{A}$ over R, then the first-order component of μ defines an element in the **truncated** Hochschild cohomology,

$$[\mu_1] \in THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^1),$$

which we also call the **first-order deformation class** of the minimal deformation μ.

We are now almost ready to prove our main classification result for deformations of A_∞ algebras. It turns out that in our particular situation, we need to incorporate a finite group action into the picture, so we now briefly explain how to do that.

Definition 2.4.10. Let H be a finite group. An **action** of H on a grading datum $\boldsymbol{G}$ is an action α of H on Y by group homomorphisms,

$$\alpha : H \to \mathrm{Hom}_{Ab}(Y, Y),$$

which preserves $\mathbb{Z}$. We will denote $\alpha(h)$ by α_h.

Example 2.4.11. In the case of Example 2.1.13, there is an action of the symmetric group $H = S_n$ on $\boldsymbol{G}_n$, by permuting the generators of $\mathbb{Z}^n$.

Definition 2.4.12. Suppose that H acts on the grading datum $\boldsymbol{G}$, and V is a $\boldsymbol{G}$-graded vector space, and we have an action

$$\rho : H \to \mathrm{Hom}_{Vect}(V, V).$$

We will denote $\rho(h)$ by ρ_h. We say that the action ρ is **$\boldsymbol{G}$-graded** if $\rho(h)$ maps V_y to $V_{\alpha_h(y)}$.

Definition 2.4.13. Suppose that we have compatible $\boldsymbol{G}$-graded actions of H on the $\boldsymbol{G}$-graded algebra R, and on the $\boldsymbol{G}$-graded R-bimodules A and B. Then H acts on the $\mathbb{Z}$-graded vector space $CC^*_{\boldsymbol{G}}(A, B)$ (preserving the grading), via

$$(h \cdot \phi)^s(a_s, \ldots, a_1) := h^{-1} \cdot \phi^s(h \cdot a_s, \ldots, h \cdot a_1).$$

We denote the H-invariant part of $CC^*_{\boldsymbol{G}}(A, B)$ by

$$CC^*_{\boldsymbol{G}}(A, B)^H := \{\phi \in CC^*_{\boldsymbol{G}}(A, B) : h \cdot \phi = \phi \text{ for all } h \in H\}.$$

Definition 2.4.14. We say that a $\boldsymbol{G}$-graded A_∞ algebra $\mathcal{A} = (A, \mu)$ over R is **strictly H-equivariant** if μ lies in $CC^2_{\boldsymbol{G}}(A)^H \subset CC^2_{\boldsymbol{G}}(A)$. Equivalently, we have

$$\mu^k(h \cdot a_k, \ldots, h \cdot a_1) = h \cdot \mu^k(a_k, \ldots, a_1)$$

for all k and all $h \in H$.

Remark 2.4.15. We remark that $TCC^2_{\boldsymbol{G}}(A, A \otimes R)^H$ is naturally a R_0^H-module (compare Remark 2.4.4).

Now we prove our main classification result for $\boldsymbol{G}$-graded deformations of A_∞ algebras over R:

Proposition 2.4.16. *Suppose that $\mathcal{A} = (A, \mu_0)$ is a $\boldsymbol{G}$-graded minimal A_∞ algebra over $\mathbb{C}$, and furthermore is strictly H-equivariant with respect to the action of some finite group H on A. Suppose that*

$$THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R)^H$$

is generated, as an R_0^H-module, by its first-order part

$$THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^1)^H,$$

and this first-order part is one-dimensional as a $\mathbb{C}$-vector space. Then any two strictly H-equivariant $\boldsymbol{G}$-graded minimal deformations of $\mathcal{A}$, whose first-order deformation classes are non-zero in

$$THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^1)^H,$$

are related by an element of $\mathrm{Aut}(R)^H$ *composed with a $\boldsymbol{G}$-graded formal diffeomorphism.*

Proof. Suppose that (A, μ) and (A, η) are two such deformations. We will construct, order-by-order, elements $\psi \in \mathrm{Aut}(R)^H$ and $F \in TCC^1_{\boldsymbol{G}}(A, A \otimes R)^H$ so that $\psi \cdot \mu = F_* \eta$. The equation that ψ and F must satisfy is

$$\psi \cdot \mu = F_* \eta.$$

We call this the A_∞ relation for the purposes of this proof.

We denote

$$\begin{aligned} \psi &= \sum_{j \geq 0} \psi_j, \text{ where } \psi_j \in (R^j)_0^H, \text{ and} \\ F &= \sum_{j \geq 0} F_j, \text{ where } F_j \in TCC^1_{\boldsymbol{G}}(A, A \otimes R^j)^H. \end{aligned}$$

We start with $F_0 = \mathrm{id}$. The order-zero component of the A_∞ equation says that $\mu_0 = \eta_0$, which is true by assumption.

Now suppose, inductively, that we have determined F_j and ψ_{j-1} for all $j \leq k-1$, H-invariant and $\boldsymbol{G}$-graded, and that

$$(\psi \cdot \mu - F_* \eta)_j = 0 \text{ for all } j \leq k-1.$$

We show that it is possible to choose F_k and ψ_{k-1} so that

$$(\psi \cdot \mu - F_* \eta)_k = 0.$$

The left hand side lies in $TCC^2_{\boldsymbol{G}}(A, A \otimes R^k)^H$.

First, we observe that

$$[\psi \cdot \mu - F_* \eta, \psi \cdot \mu + F_* \eta] = 0,$$

by expanding out the brackets: the cross-terms vanish by symmetry ($\psi \cdot \mu$ and $F_*\eta$ both have degree 2), and the other terms vanish because $\psi \cdot \mu$ and $F_*\eta$ are A_∞ structures. Now note that

$$(\psi \cdot \mu + F_* \eta)_0 = \mu_0 + \eta_0 = 2\mu_0,$$

so taking the order-k component of the previous equation gives us

$$\begin{aligned} [\psi \cdot \mu - F_* \eta, \mu_0]_k &= 0 \\ \Rightarrow \delta\left((\psi \cdot \mu - F_* \eta)_k\right) = 0, \end{aligned}$$

regardless of what F_k and ψ_{k-1} are.

Now we pick out the terms in $(\psi \cdot \mu - F_* \eta)_k$ that involve F_k and ψ_{k-1}. After a little calculation along the lines of the proof of Lemma 2.3.17, we see that it has the form

$$(\psi \cdot \mu - F_* \eta)_k = \psi_{k-1} \cdot \mu_1 + F_k \circ \mu_0 - \eta_0 \circ F_k + D_k,$$

where D_k contains all the terms that do not involve ψ_{k-1} or F_k. We recall that $\eta_0 = \mu_0$, and identify the Hochschild differential:

$$(\psi \cdot \mu - F_* \eta)_k = \psi_{k-1} \cdot \mu_1 + \delta(F_k) + D_k.$$

Note that if we set $F_k = 0$ and $\psi_{k-1} = 0$, our previous argument shows that $\delta(D_k) = 0$, so D_k defines a class

$$[D_k] \in THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^k)^H.$$

Thus, we need to choose ψ_{k-1} so that $\psi_{k-1} \cdot [\mu_1] = -[D_k]$ in the truncated Hochschild cohomology $THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^k)^H$. We can do this by our assumption that $[\mu_1]$ is non-zero, hence generates the one-dimensional first-order component $THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^1)^H$, which generates $THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R)^H$ as a R^H_0-module. We then choose F_k to effect the Hochschild coboundary between $\psi_{k-1} \cdot \mu_1$ and $-D_k$. We can make F_k H-invariant by averaging over H.

Finally, note that at first order, we have

$$\psi_0 \cdot [\mu_1] = [\eta_1],$$

from which it follows that $\psi_0 \neq 0$, because $[\mu_1]$ and $[\eta_1]$ are both non-zero, so indeed $\psi \in \mathrm{Aut}(R)^H$. □

2.5. **Computations.** In this section, we introduce specific $\boldsymbol{G}$-graded A_∞ algebras and deformations, and use the results of the previous sections to prove classification theorems for them.

Let us introduce some notation. We fix an integer $n \geq 4$ (we will be considering hypersurfaces in $\mathbb{CP}^{n-1}$) and $a \geq 1$ (this will be the degree of the hypersurface in $\mathbb{CP}^{n-1}$ that we will consider). In our intended applications in the current paper, a will be either 1 or n.

Throughout this section, we will be using the grading datum $\boldsymbol{G} := \boldsymbol{G}^n_1$ from Example 2.1.13. For an element $y \in Y$, we will denote

$$|y| := y_{[n]} \cdot y.$$

We will denote by H the symmetric group on n elements, and recall that it acts on $\boldsymbol{G}$ (see Example 2.4.11).

We recall the $\boldsymbol{G}$-graded exterior algebra

$$A := A_n := \Lambda(U_n)$$

of Definition 2.2.12. For each subset $K \subset [n]$, we denote the corresponding element of A by

$$\theta^K := \bigwedge_{j \in K} \theta^j.$$

We equip the vector space U_n with an H-action, which up to sign is the obvious action by permuting basis elements. In other words,

$$h(u_j) = \pm u_{h(j)}.$$

We will not need to specify the actual signs. There is an induced action of H on A.

We recall the $\boldsymbol{G}$-graded ring

$$R_a := R_a^n := \mathbb{C}[[r_1, \ldots, r_n]]$$

from Definition 2.2.13. We give a name to one important element of R_a: we denote

$$T := r^{y_{[n]}} = r_1 \ldots r_n.$$

We equip R_a with an H action, which up to sign is the obvious action by permuting basis elements. We furthermore denote

$$R := R_n$$

because $a = n$ is the most important case we will consider. Finally, we will denote by R_a^j the order-j part of R_a. Note the change of notation from Definition 2.2.13, where the superscript denoted the number of generators. We hope this does not cause confusion.

Definition 2.5.1. Suppose that $A \cong \Lambda(U)$ is an exterior algebra over $\mathbb{C}$, and R is an arbitrary commutative ring. We define the R-algebra $A \otimes R$, and the **Hochschild-Kostant-Rosenberg map**

$$\begin{aligned} \Phi : CC^*(A \otimes R|R) &\to R[[U]] \otimes A, \\ \phi &\mapsto \sum_{s=0}^{\infty} \phi^s(\boldsymbol{u}, \ldots, \boldsymbol{u}), \end{aligned}$$

where

$$\boldsymbol{u} := \sum_{j=1}^{n} u_j \theta^j.$$

It has the following properties:

- If R is a field, then Φ induces an isomorphism on the level of cohomology (the Hochschild-Kostant-Rosenberg isomorphism [28]);
- If S is an R-algebra, then there is a commutative diagram

$$\begin{tikzcd} CC^*(A \otimes R|R) \otimes_R S \arrow[r, "\Phi_R \otimes \mathbf{1}"] \arrow[d] & (R[[U]] \otimes A) \otimes_R S \arrow[d] \\ CC^*(A \otimes S|S) \arrow[r, "\Phi_S"] & S[[U]] \otimes A. \end{tikzcd}$$

We call the latter property 'naturality' of the HKR map.

Definition 2.5.2. Let us fix $a = n$. We say that a $\boldsymbol{G}$-graded A_∞ algebra $\mathcal{A}$ over R has **type A** if it satisfies the following properties:

- Its underlying R-module and order-0 cohomology algebra is
$$(\mathcal{A}, \mu_0^2) \cong A \otimes R;$$
- It is strictly H-equivariant;
- It satisfies
$$\Phi(\mu^{\geq 3}) = u_1 \ldots u_n + \sum_{j=1}^n r_j u_j^n + \mathcal{O}(r^2).$$

Now we state the main result we will prove in this section:

Theorem 3. *Suppose that* $\mathcal{A}_1 = (A \otimes R, \mu)$ *and* $\mathcal{A}_2 = (A \otimes R, \eta)$ *are two* $\boldsymbol{G}$*-graded* A_∞ *algebras over* R *of type A. Then there exists* $\psi \in R_0^H$*, of the form*

$$\psi(r_1, \ldots, r_n) = 1 + \sum_{j=1}^\infty c_j T^j,$$

and a $\boldsymbol{G}$*-graded formal diffeomorphism* $F \in \mathfrak{G}_R(A)$*, such that*

$$\psi \cdot \mathcal{A}_1 = F_* \mathcal{A}_2.$$

We will now give a brief outline of the proof. The first step is to prove (in Corollary 2.5.7, using Proposition 2.3.24) that the order-0 parts (A, μ_0) and (A, η_0) are quasi-isomorphic. The classification of the order-0 part is governed by the Hochschild cohomology $HH^*_{\boldsymbol{G}}(A)$, which is determined via the Hochschild-Kostant-Rosenberg (HKR) isomorphism. We thereafter denote by $\boldsymbol{\mathcal{A}} \cong (A, \mu_0)$ this (unique up to formal diffeomorphism) order-0 part.

We then study deformations of $\boldsymbol{\mathcal{A}}$ over R. The classification of such deformations is governed by the Hochschild cohomology $HH^*_{\boldsymbol{G}}(\boldsymbol{\mathcal{A}}, \boldsymbol{\mathcal{A}} \otimes R)$, which is also determined from the HKR isomorphism, via a spectral sequence. We apply Proposition 2.4.16 to show that such deformations are unique up to A_∞ quasi-isomorphism and the action of $\mathrm{Aut}(R)^H$.

Now let us begin. We first explain how to use the HKR isomorphism [28] to calculate $HH^*_{\boldsymbol{G}}(A)$, and more generally $HH^*_{\boldsymbol{G}}(A, A \otimes R_a)$. The HKR isomorphism show that, if $A = \Lambda(U)$ is a $\boldsymbol{G}$-graded exterior algebra over $\mathbb{C}$, then the map

$$\Phi : (CC^*_c(A), \delta) \to (\mathbb{C}[U] \otimes \Lambda(U), 0),$$

is a quasi-isomorphism of $Y \oplus \mathbb{Z}$-graded chain complexes. This allows us to determine

$$HH^*(A) \cong \mathbb{C}[[U]] \otimes \Lambda(U)$$

by taking the completion with respect to the length filtration, and also $HH^*_{\boldsymbol{G}}(A)$, where now we equip A with its $\boldsymbol{G}$-grading. We will make the $\boldsymbol{G}$-grading of the right-hand side explicit in Lemma 2.5.4.

More generally, it follows that there is an isomorphism of $Y \oplus \mathbb{Z}$-graded vector spaces,

$$\Phi : HH^*_c(A, A \otimes R_a) \to \mathbb{C}[U] \otimes \Lambda(U) \otimes R_a.$$

$\mathbb{C}[U] \otimes \Lambda(U) \otimes R_a$ is generated by terms $u^{\boldsymbol{b}} \theta^K r^{\boldsymbol{c}}$, where $\boldsymbol{b}, \boldsymbol{c} \in Y_{\geq 0}$ (recall $Y \cong \mathbb{Z}^n$, so we define $Y_{\geq 0} := \mathbb{Z}^n_{\geq 0}$) and $K \subset [n]$. If $\boldsymbol{b} = \sum_j b_j e_j$, then this is the image under Φ of a Hochschild cochain which sends

$$\bigotimes_j (\theta^j)^{\otimes b_j} \mapsto r^{\boldsymbol{c}} \theta^K.$$

We start by examining what the various gradings on a Hochschild cochain tell us.

Lemma 2.5.3. *If a generator* $\tau \in CC^{s+t}_{c,\boldsymbol{G}} \left(A, A \otimes R^j_a \right)^t$ *sends*

$$\bigotimes_{i=1}^s \theta^{K_i} \mapsto r^{\boldsymbol{c}} \theta^{K_0},$$

then we have

$$j = |\boldsymbol{c}| \tag{2.5.1}$$

$$a\boldsymbol{c} + y_{K_0} - \sum_{i=1}^s y_{K_i} = q y_{[n]} \text{ in } Y, \text{ for some } q \tag{2.5.2}$$

$$t = (n-2)q + (2-a)j. \tag{2.5.3}$$

Proof. Equation (2.5.1) follows by definition. To prove Equations (2.5.2) and (2.5.3), we recall that the grading of θ_j is $(-1, y_j)$ (Example 2.2.9) and the grading of r_j is $(2-2a, a y_j)$ (Example 2.2.10). We alter the grading datum $\boldsymbol{G}$ by an automorphism sending

$$(j, y) \mapsto (j + y_{[n]} \cdot y, y).$$

This is equivalent to considering the pseudo-grading datum with the same exact sequence as $\boldsymbol{H}_n$, but

$$c = 2(n-1) - y_{[n]} \cdot y_{[n]} = n - 2.$$

Then θ_j has grading $(0, y_j)$ and r_j has grading $(2-a, a y_j)$.

If the grading of τ is $f(t) = (t, 0) \in (\mathbb{Z} \oplus Y)/Z$, then

$$(0, y_{K_0}) + ((2-a)|\boldsymbol{c}|, a\boldsymbol{c}) - \sum_{j=1}^s (0, y_{K_j}) \cong (t, 0) \text{ modulo } Z.$$

Recalling that the image of Z in $\mathbb{Z} \oplus Y$ is given by

$$((2-n)q, q y_{[n]})$$

(in the altered grading datum), we have

$$(0, y_{K_0}) + ((2-a)|\boldsymbol{c}|, a\boldsymbol{c}) - \sum_{j=1}^s (0, y_{K_j}) = (t, 0) + q(2-n, y_{[n]}),$$

from which the result follows. □

Now recalling the Hochschild-Kostant-Rosenberg isomorphism, a generator of $HH^{s+t}_{c,\boldsymbol{G}}(A, A \otimes R_a)^t$ has the form $u^{\boldsymbol{b}} \theta^K r^{\boldsymbol{c}}$, where $\boldsymbol{b}$ and $\boldsymbol{c}$ are elements of $Y_{\geq 0}$, and $K \subset [n]$. We examine what the gradings tell us about such a generator of the Hochschild cohomology.

Lemma 2.5.4. *If $u^{\boldsymbol{b}}\theta^K r^{\boldsymbol{c}}$ is a generator of $HH^{s+t}_{c,\boldsymbol{G}}\left(A, A \otimes R^j_a\right)^t$, then the following equations hold:*

$$\begin{aligned} j &= |\boldsymbol{c}| && (2.5.4)\\ s &= |\boldsymbol{b}| && (2.5.5)\\ y_K + a\boldsymbol{c} - \boldsymbol{b} &= q y_{[n]} \text{ in } Y, \text{ for some } q && (2.5.6)\\ t &= (n-2)q + (2-a)j && (2.5.7)\\ |K| &= s + t + 2(q-j). && (2.5.8)\\ (n-2)|K| &= (n-2)s + nt + 2(a-n)j && (2.5.9)\end{aligned}$$

Proof. Equations (2.5.4) and (2.5.5) hold by definition. Equation (2.5.6) follows from Equation (2.5.2). Equation (2.5.7) follows from Equation (2.5.3). The first step in proving Equations (2.5.8) and (2.5.9) is to take the dot product of Equation (2.5.6) with $y_{[n]}$:

$$\begin{aligned} y_{[n]} \cdot (y_K + a\boldsymbol{c} - \boldsymbol{b}) &= q y_{[n]} \cdot y_{[n]} \\ \Rightarrow |K| &= |\boldsymbol{b}| - a|\boldsymbol{c}| + nq \\ &= s - aj + nq. \end{aligned}$$

To prove equation (2.5.8), we use equation (2.5.7) to substitute for nq:

$$\begin{aligned} |K| &= s - aj + t + (a-2)j + 2q \\ &= s + t + 2(q-j). \end{aligned}$$

We use the same equation to prove equation (2.5.9), but this time we first multiply by n then substitute in equation (2.5.7):

$$\begin{aligned} (n-2)|K| &= (n-2)(s - aj + nq) \\ &= (n-2)s - (n-2)aj + n(t + (a-2)j) \\ &= (n-2)s + nt + 2(a-n)j. \end{aligned}$$

□

We now set about determining the order-0 part of an A_∞ algebra of type A. We identify the possible $\boldsymbol{G}$-graded A_∞ algebras $\mathcal{A} = (A, \mu_0)$ over $\mathbb{C}$ with underlying vector space and μ_0^2 given by A.

Lemma 2.5.5. *We have*

$$CC^*_{c,\boldsymbol{G}}(A)^t \cong 0$$

unless t is divisible by $(n-2)$.

Proof. Follows from Equation (2.5.7) with $j = 0$. □

Lemma 2.5.6. *The Hochschild cohomology of A satisfies, for $d > 0$,*

$$HH^2_{c,\boldsymbol{G}}(A)^{2-d} \cong \begin{cases} \mathbb{C} \cdot u^{y_{[n]}} & \text{if } d = n, \\ 0 & \text{otherwise,} \end{cases}$$

where $u^{y_{[n]}} = u_1 \dots u_n$.

Proof. Let $u^{\boldsymbol{b}}\theta^K$ be a generator of $HH^2_{c,\boldsymbol{G}}(A)^{2-d}$. Equation (2.5.7), with $j = 0$, yields

$$d = 2 - (n-2)q.$$

We want $d > 0$, so we have $q < 0$. Now equation (2.5.8), with $j = 0$, yields

$$2 + 2q = |K| \geq 0.$$

Thus, $-1 \leq q < 0$, so $q = -1$ and $|K| = 0$, so $K = \phi$.

Equation (2.5.6) now yields $\boldsymbol{b} = y_{[n]}$. Therefore $u^{\boldsymbol{b}}\theta^K = u^{y_{[n]}}$, as required. □

Corollary 2.5.7. *There is an A_∞ quasi-isomorphism between the 0th-order parts of $\mathcal{A}_1$ and $\mathcal{A}_2$:*

$$(A, \mu_0) \cong (A, \eta_0).$$

Proof. First, we observe that $CC^2_{c,\boldsymbol{G}}(A)^1 \cong 0$ by Equation (2.5.3) (with $j = 0$, and assuming $n \geq 4$), so the A_∞ algebras are minimal. Furthermore, $\mu_0^d = 0$ for $2 < d < n$ because $CC^2_{c,\boldsymbol{G}}(A)^{2-d} \cong 0$ by Lemma 2.5.5. The result now follows from Proposition 2.3.24, because the deforming class of an algebra of type A is

$$\Phi_0(\mu_0) = u_1 \dots u_n = u^{y_{[n]}}$$

by definition. □

Henceforth, we will denote by $\mathcal{A}$ any $\boldsymbol{G}$-graded minimal A_∞ algebra over $\mathbb{C}$ with cohomology algebra given by A, and whose order-n deforming class is a non-zero multiple of $u^{y_{[n]}}$. The previous lemma says that $\mathcal{A}$ is well-defined up to quasi-isomorphism.

We will consider deformations of $\mathcal{A}$ over R_a, which are controlled by the Hochschild cohomology with coefficients in R_a,

$$HH^*_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R_a).$$

Recall from Remark 2.3.28 that the filtration by length on the Hochschild complex $CC(A)$ yields a spectral sequence for the Hochschild cohomology, with

$$E_2^{*,*} = HH^*_c(A, A).$$

Lemma 2.5.8. *The spectral sequence induced by the length filtration on the Hochschild cochain complex $CC^*(\mathcal{A})$ converges to the Hochschild cohomology $HH^*(\mathcal{A})$.*

Proof. By Remark 2.3.28, it suffices to prove that the spectral sequence is regular.

A generator $u^{\boldsymbol{b}}\theta^K$ of $HH^*_c(A)$ changes the Y-degree by

$$(-|K| + |\boldsymbol{b}|, -\boldsymbol{b} + y_K) \in \mathbb{Z} \oplus Y/(2, y_{[n]}),$$

and its length is $s = |\boldsymbol{b}|$. Therefore, its grading is

$$(-|K| + 2|\boldsymbol{b}|, -\boldsymbol{b} + y_K)$$

(recalling the conventions for grading Hochschild cochains). Using a suitably altered pseudo-grading datum as in the proof of Lemma 2.5.3, this is equivalent to a grading

$$(|K|, -\boldsymbol{b} + y_K)$$

(note: $c = 2$ in this altered pseudo-grading datum).

Now recall that the differential on page d of the spectral sequence maps

$$\delta_d : E_d^{(y,s)} \to E_d^{(y+f(1),s+d)}.$$

If both domain and codomain of the differential are to be non-zero, then we must have generators $u^{\boldsymbol{b}_1}\theta^{K_1}$ and $u^{\boldsymbol{b}_2}\theta^{K_2}$ such that

$$(|K_1| + 1, -\boldsymbol{b}_1 + y_{K_1}) = (|K_2|, -\boldsymbol{b}_2 + y_{K_2}) + q(2, y_{[n]})$$

and

$$|\boldsymbol{b}_2| = |\boldsymbol{b}_1| + d.$$

Because $0 \le |K_1|, |K_2| \le n$, and $|K_1| + 1 = |K_2| + 2q$, we must have $q \le (n+1)/2$. We also have

$$\boldsymbol{b}_2 = \boldsymbol{b}_1 + y_{K_2} - y_{K_1} + qy_{[n]},$$

and hence that

$$|\boldsymbol{b}_2| = |\boldsymbol{b}_1| + |K_2| - |K_1| + nq = |\boldsymbol{b}_1| + 1 + (n-2)q.$$

Therefore, for the differential δ_d to be non-zero, we must have

$$d = |\boldsymbol{b}_2| - |\boldsymbol{b}_1| = (n-2)q + 1 \le \frac{(n+1)(n-2)}{2} + 1.$$

Hence, for d sufficiently large, the differential vanishes, so the spectral sequence is regular. □

Now, if we are to apply the deformation theory of Section 2.4, we need to know that our deformations are minimal.

Lemma 2.5.9. *If $a = n$, and $n \ge 5$, then any $\boldsymbol{G}$-graded deformation of $\mathcal{A}$ over R is minimal.*

Proof. Equation (2.5.3) with $a = n$ shows that

$$CC^2_{c,\boldsymbol{G}}(A, A \otimes R)^t \cong 0$$

unless t is divisible by $(n-2)$. The μ^0 and μ^1 terms live in the spaces with $t = 2$ and $t = 1$, respectively. So when $n \ge 5$, any deformation of $\mathcal{A}$ over R satisfies $\mu^0 = 0$ and $\mu^1 = 0$, hence is minimal. □

Remark 2.5.10. Lemma 2.5.9 is the only place where we need the assumption $n \ge 5$, as opposed to $n \ge 4$. If $n = 4$, then $CC^2_{c,\boldsymbol{G}}(A, A \otimes R)^0$ is one-dimensional, generated by the element $T\theta^{[4]}$. These deformations can be ruled out in our situation with some additional input (namely, by building a strictly unital model for our category), but we prefer to avoid this additional complication. See [2].

The next step is to determine the first-order deformation space.

Lemma 2.5.11. *If $n \ge 4$, then the vector space*

$$HH^2_{\boldsymbol{G}}(A, A \otimes R^1_a)$$

is generated by the elements

$$r_j u_j^a,$$

for $j = 1, \ldots, n$.

Proof. We apply Lemma 2.5.4 with $s+t=2$ and $j=1$. Equation (2.5.8) yields

$$|K| = 2q \geq 0.$$

Furthermore, Equation (2.5.7) gives

$$\begin{aligned} 2-s &= (n-2)q+2-a \\ \Rightarrow a-s &= (n-2)q. \end{aligned}$$

Furthermore, we have $\boldsymbol{c} = y_k$ for some k. Taking the dot product of y_k with Equation (2.5.6) gives us

$$\begin{aligned} y_k \cdot (y_K + a\boldsymbol{c} - \boldsymbol{b}) &= q y_k \cdot y_{[n]} \\ \Rightarrow (0 \text{ or } 1) &= -a + y_k \cdot \boldsymbol{b} + q \\ &\leq -a+s+q \\ &= (3-n)q. \end{aligned}$$

If $n \geq 4$, then $3-n<0$ and $q \geq 0$, so we must have $q=0$. Thus, we have $|K| = 2q = 0$, so $K = \phi$. From Equation (2.5.6), we obtain $\boldsymbol{b} = a y_j$, so the generator has the form

$$r_j u_j^a.$$

□

Having determined the first-order deformation space, we now seek to understand the higher-order parts of the deformation. The following lemma will be useful:

Lemma 2.5.12. *If $r^{\boldsymbol{c}} u^{\boldsymbol{b}} \theta^K$ is a generator of $THH^2_{\boldsymbol{G}}(A, A \otimes R_a)$, and $j = |\boldsymbol{c}| \geq 2$, then $\boldsymbol{c} \geq y_{[n]}$.*

Proof. We apply Lemma 2.5.4, with $s+t=2$. Equation (2.5.8) yields

$$0 \leq |K| = 2(1+q-j).$$

It follows that $q \geq j-1$. Now we split into two cases, depending on whether $q = j-1$ or $q > j-1$.

Case 1: $q = j-1$. In this case, we have $|K| = 0$ so $K = \phi$. Thus Equation (2.5.6) gives

$$a\boldsymbol{c} = q y_{[n]} + \boldsymbol{b}.$$

Taking the dot product with y_k yields

$$a c_k = q + b_k \geq q = j-1 > 0,$$

since $j \geq 2$. Hence $c_k \geq 1$ for all k, and the result is proved.

Case 2: $q > j-1$. In this case, we have $q > j-1 \geq 1$, so $q \geq 2$. Thus, taking the dot product of Equation (2.5.6) with y_k gives

$$a c_k = q + b_k - y_k \cdot y_K \geq q-1 > 0.$$

Hence $c_k \geq 1$ for all k, and the result is proved. □

Now if $a = n$, then the grading of $T = r^{y_{[n]}} \in R_0$ is 0 in $\mathbb{Z}$, and furthermore T is clearly H-invariant. Thus $\mathbb{C}[[T]] \subset R_0^H$ (in fact one can show they are equal). It follows that $THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R)^H$ is a $\mathbb{C}[[T]]$-module.

Lemma 2.5.13. *Suppose that $a = n$, and $n \geq 4$. Then the H-invariant part of the truncated Hochschild cohomology group*

$$THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R)^H$$

is generated, as a $\mathbb{C}[[T]]$-module, by its first-order part

$$THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^1)^H.$$

Proof. We prove the result by induction on the order j, and using the spectral sequence induced by the length filtration.

When $j = 0$, we must show that $THH^2_{\boldsymbol{G}}(\mathcal{A}) \cong 0$. Let $u^{\boldsymbol{b}}\theta^K$ be a generator of $THH^2_{\boldsymbol{G}}(A)$. By equation (2.5.7) with $j = 0$, we have

$$t = (n-2)q \leq 0$$

(since we are considering the truncated Hochschild cohomology). By equation (2.5.8) with $j = 0$, we have

$$|K| = 2(1+q) \geq 0.$$

Thus we have $-1 \leq q \leq 0$. We have two cases:

Case 1: $q = 0$. In this case, $|K| = 2$, $t = 0$, so $s = 2$. Equation (2.5.6) yields

$$y_K = \boldsymbol{a},$$

so our generator has the form $u_i u_k \theta^i \wedge \theta^k$. But now, if $\sigma_{ik} \in H$ denotes the transposition of elements i and k, we have

$$\sigma_{ik}(u_i u_k \theta^i \wedge \theta^k) = u_k u_i \theta^k \wedge \theta^i = -u_i u_k \theta^i \wedge \theta^k.$$

Hence, the coefficient of this term in any H-invariant element of the truncated Hochschild cohomology must be 0.

Remark 2.5.14. Recall that H may differ from the obvious action by permutation of basis elements, by some signs. However, each sign appears twice (once for u_i, once for θ^i), so they do not affect this computation.

Case 2: $q = -1$. In this case, $|K| = 0$, so $K = \phi$. Equation (2.5.6) yields

$$y_{[n]} = \boldsymbol{b}.$$

Thus, the generator is $u^{y_{[n]}}$, the deforming class identified in Lemma 2.5.6. However, this generator gets killed by the first non-trivial differential of the spectral sequence. To see this, observe that, because $\mu^d = 0$ for $2 < d < n$ by Lemma 2.5.5, the first non-trivial differential is δ_{n-1}, and is given by

$$\delta_{n-1}(\phi) = [[\mu^n], \phi] = [u^{y_{[n]}}, \phi]$$

(see Remark 2.3.29). The Gerstenhaber bracket on $CC^*(A)$ gets carried to the Schouten bracket on polyvector fields $\mathbb{C}[[U]] \otimes \Lambda(U)$. It follows quickly from the explicit form of the Schouten bracket (see [4, Equation (3.7)]) that for any $W \in \mathbb{C}[[U]]$,

$$[W, \eta] = \iota_{dW}(\eta),$$

and therefore that the cohomology of $[W, -]$ is the cohomology of the Koszul complex associated to dW. In particular, the class W itself gets killed by taking the cohomology of this differential. In our case, this means that $u^{y_{[n]}}$ is killed by δ_{n-1}.

This completes the proof that $THH^2_{\boldsymbol{G}}(\mathcal{A}) \cong 0$.

When $j = 1$, the statement is simply that the first-order part is generated by the first-order part. So it remains to prove that, for $j \geq 2$, the order-j part is generated by the first-order part.

Suppose, inductively, that

$$THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^k) \subset \mathbb{C}[[T]] \cdot THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^1)$$

for all $k \leq j-1$, for some $j \geq 2$. We prove that the same holds for $k = j$. Let

$$r^{\boldsymbol{c}} u^{\boldsymbol{b}} \theta^K \in THH^2_{\boldsymbol{G}}(A, A \otimes R^j)$$

be a generator. Because $j \geq 2$, it follows from Lemma 2.5.12 that $r^{\boldsymbol{c}} = T \cdot r^{\boldsymbol{c}'}$, where

$$\boldsymbol{c}' = \boldsymbol{c} - y_{[n]} \geq 0.$$

Because $T \in R_0$, we have

$$r^{\boldsymbol{c}'} u^{\boldsymbol{b}} \theta^K \in THH^2_{\boldsymbol{G}}(A, A \otimes R^{j-n}).$$

It follows that

$$THH^2_{\boldsymbol{G}}(A, A \otimes R^j) = T \cdot THH^2_{\boldsymbol{G}}(A, A \otimes R^{j-n}),$$

and hence, from the spectral sequence induced by the length filtration, that

$$THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^j) = T \cdot THH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes R^{j-n}).$$

The result now follows by taking H-invariant parts of this equation.

This completes the inductive step, and hence the proof. $\square$

Corollary 2.5.15. *There exists a $\psi \in \mathbb{C}[[T]]$, with $\psi(0) = 1$, and an $\boldsymbol{G}$-graded formal diffeomorphism F, such that*

$$\psi \cdot \mathcal{A}_1 = F_* \mathcal{A}_2.$$

Proof. By Corollary 2.5.7, there is a formal diffeomorphism F_0 from (A, η_0) to (A, μ_0). Therefore we can push forward η by F_0, and reduce to the case where $\mu_0 = \eta_0$. The result then follows from Proposition 2.4.16 and Lemma 2.5.13. $\square$

This completes the proof of Theorem 3.

We now make one final computation of Hochschild cohomology. For this computation, we will be interested in an extra structure on Hochschild cohomology: the **Yoneda product**, which makes it into an associative algebra. In fact, together with the Gerstenhaber bracket, this makes the Hochschild cohomology into a **Gerstenhaber algebra** (see [29]).

Definition 2.5.16. The Hochschild cohomology $HH^*(\mathcal{A})$ of a $\boldsymbol{G}$-graded A_∞ algebra $\mathcal{A}$ carries the $\boldsymbol{G}$-graded, associative **Yoneda product**, which has the form

$$(\phi \bullet \psi)^n(x_n, \ldots, x_1) :=$$

$$\sum_{i+j+k+l+m=n} (-1)^{\dagger} \mu^{m+k+i+2}(x_{i+j+k+l+m}, \ldots, \phi^l(x_{i+j+k+l}, \ldots), x_{i+j+k}, \ldots, \psi^j(x_{i+j}, \ldots), x_i, \ldots),$$

where $\dagger$ is some sign.

Remark 2.5.17. We record the following useful information about the Yoneda product:

- If $\boldsymbol{k}$ is a field, and U a $\boldsymbol{k}$-vector space, then the HKR isomorphism
$$\Phi : HH^*(\Lambda(U)) \to \boldsymbol{k}[[U]] \otimes \Lambda(U),$$
is an isomorphism of $\boldsymbol{k}$-algebras, where the product on $HH^*(\Lambda(U))$ is the Yoneda product, and the product on polyvector fields is the wedge product (see [30, Section 8]);
- The spectral sequence $E_d^{s,t} \Rightarrow HH^*(\mathcal{A})$ induced by the length filtration (Remark 2.3.28) respects the multiplicative structure of the Yoneda product.

We now carry out our final Hochschild cohomology computation. Suppose that $\mathcal{A}$ is an A_∞ algebra of type A, and $\underline{\mathcal{A}}$ its extension to a $\boldsymbol{G}_1^n$-graded category. We denote
$$\widetilde{\mathcal{A}} := \boldsymbol{p}_1^* \underline{\mathcal{A}},$$
where $\boldsymbol{q}_1$ and $\boldsymbol{p}_1$ are the morphisms of grading data defined in Lemma 2.1.14. Note that $\boldsymbol{q}_{1*}\widetilde{\mathcal{A}}$ is a $\boldsymbol{G}_{\mathbb{Z}}$-graded (or equivalently, $\mathbb{Z}$-graded) A_∞ category, over the $\mathbb{Z}$-graded coefficient ring $\boldsymbol{q}_{1*}\boldsymbol{p}_1^* R \cong R$, which has degree $0 \in \mathbb{Z}$, by Remark 2.2.11. Therefore, the homomorphism
$$\begin{array}{rcl} R & \to & \Lambda, \\ r_j & \mapsto & r \text{ for all } j \end{array}$$
respects the $\mathbb{Z}$-grading, and we can form the $\boldsymbol{G}_{\mathbb{Z}}$-graded Λ-linear A_∞ category
$$\widetilde{\mathcal{A}}_{nov} := \boldsymbol{q}_{1*}\widetilde{\mathcal{A}} \otimes_R \Lambda.$$
Our aim now is to compute a certain part of
$$HH^*\left(\widetilde{\mathcal{A}}_{nov}\middle| \Lambda\right).$$

We observe that Y_1^n acts on $\underline{\mathcal{A}}$ by shifts, hence also acts on $\widetilde{\mathcal{A}}_{nov}$, hence also acts on $CC^*(\widetilde{\mathcal{A}}_{nov})$. The action of $\ker(Y_n^n \to \mathbb{Z}) \subset Y_1^n$ is trivial, because shifts in this subgroup become trivial when we take $\boldsymbol{q}_{1*}$ of our category. The action of $\mathbb{Z}$ is also trivial, because we consider only $\mathbb{Z}$-equivariant cochains, by definition. It follows that the action of the image of Y_n^n in Y_1^n is trivial, so there is an action of
$$\tilde{\Gamma}_n^* \cong Y_1^n / Y_n^n$$
on $CC^*(\widetilde{\mathcal{A}}_{nov})$. Note that this is the same $\tilde{\Gamma}_n^*$ as in Definition 1.2.4.

Lemma 2.5.18. *There is an isomorphism of $\mathbb{Z}$-graded Λ-algebras,*
$$HH^*\left(\widetilde{\mathcal{A}}_{nov}\right)^{\tilde{\Gamma}_n^*} \cong \Lambda[\alpha]/\alpha^{n-1},$$
where α has degree 2.

Proof. We consider the spectral sequence induced by the length filtration on
$$CC^*\left(\widetilde{\mathcal{A}}_{nov}\right)^{\tilde{\Gamma}_n^*}.$$
As we saw in Remark 2.3.28, the filtration is bounded above, hence exhaustive, and is also complete. So if we can prove that it is regular, then it must converge to the Hochschild cohomology.

We observe that there is an obvious morphism of chain complexes,

$$CC^*\left(\boldsymbol{q}_{1*}\boldsymbol{p}_1^*\underline{\mathcal{A}}\right)\otimes_R\Lambda\to CC^*\left(\widetilde{\mathcal{A}}_{nov}\right),$$

and indeed the A_∞ structure maps on $\widetilde{\mathcal{A}}_{nov}$ are the image of the A_∞ structure maps of $\mathcal{A}$ under this morphism. However, this is **not** necessarily a quasi-isomorphism, because the Hochschild cochain complex is defined as a direct product over cochains of all lengths $s\geq 0$, and arbitrary direct products do not commute with $\otimes_R\Lambda$.

On the other hand, **finite** direct products **do** commute with $\otimes_R\Lambda$, so the above morphism of chain complexes does induce an isomorphism after we quotient by chains of length greater than some fixed s. It follows that there is an isomorphism between the spectral sequences induced by the length filtrations on

$$CC^*\left(\boldsymbol{q}_{1*}\boldsymbol{p}_1^*\underline{\mathcal{A}}\right)\otimes_R\Lambda$$

and on

$$CC^*\left(\widetilde{\mathcal{A}}_{nov}\right)$$

(it is just that the filtration on the former chain complex is not complete, so the associated spectral sequence may not converge to its cohomology).

Now, applying Remarks 2.3.41 and 2.3.38, we have isomorphisms

$$\begin{aligned}CC^*\left(\boldsymbol{q}_{1*}\boldsymbol{p}_1^*\underline{\mathcal{A}}\right)^{\tilde{\Gamma}_n^*} &\cong \boldsymbol{q}_{1*}CC^*\left(\boldsymbol{p}_1^*\underline{\mathcal{A}}\right)^{\tilde{\Gamma}_n^*}\text{ (up to completing with respect to the length filtration)}\\ &\cong \boldsymbol{q}_{1*}\boldsymbol{p}_1^*CC^*\left(\underline{\mathcal{A}}\right)\\ &\cong \boldsymbol{q}_{1*}\boldsymbol{p}_1^*CC^*\left(\mathcal{A}\right).\end{aligned}$$

It follows from the preceding discussion that, if it is regular, then the spectral sequence induced by the length filtration on

$$\boldsymbol{q}_{1*}\boldsymbol{p}_1^*CC^*\left(\mathcal{A}\right)\otimes_R\Lambda$$

converges to

$$HH^*\left(\widetilde{\mathcal{A}}_{nov}\right)^{\tilde{\Gamma}_n^*}.$$

Now, note that $\mu^s=0$ in $\mathcal{A}$ unless $s-2$ is divisible by $(n-2)$ (by Equation (2.5.3) with $a=n$ and $s+t=2$). In particular, $\mathcal{A}$ is minimal. Next, we show that there is an algebra isomorphism $(\mathcal{A},\mu_0^2)\cong(\mathcal{A},\mu^2)$. I.e., the higher-order terms in the product μ^2 can be absorbed into the order-0 product μ_0^2, which we know to be the exterior product. This follows immediately from the calculation

$$THH^2_{\boldsymbol{G}}(A,A\otimes R)^{0,H}\cong 0,$$

which was carried out in the proof of Lemma 2.5.13 (as 'Case 1'), together with a deformation theory argument. Thus, we can replace $\mathcal{A}$ by a quasi-isomorphic A_∞ algebra $\mathcal{A}'$, whose underlying algebra is the exterior algebra

$$\left(\mathcal{A}',\mu^2\right)\cong A\otimes R.$$

Furthermore, $\mathcal{A}'$ and $\mathcal{A}$ coincide to order n, because the higher-order terms in μ^2 were functions of $T=r_1\ldots r_n$. So $\mathcal{A}'$ is still of type A. Henceforth we'll simply write $\mathcal{A}$ for this replacement.

Now the E_2 page of the spectral sequence induced by the length filtration on

$$CC^*(\mathcal{A}) \otimes_R \Lambda$$

is the Hochschild cohomology of the associative algebra $(\mathcal{A} \otimes_R \Lambda, \mu^2 \otimes 1)$. By the preceding arguments, this is exactly $A \otimes \Lambda$, the exterior algebra over the field Λ. Its Hochschild cohomology is given by the HKR isomorphism (because Λ is a field), so we have

$$E_2 = \boldsymbol{q}_{1*}\boldsymbol{p}_1^*(\Lambda[[U]] \otimes A).$$

Because $\mu^s = 0$ for $2 < s < n$, the first non-trivial differential is δ_{n-1}, which is given by Gerstenhaber bracket with the order-n deformation class $[\mu^n]$. The Gerstenhaber bracket gets carried to the Schouten bracket under the HKR isomorphism, and μ^n gets carried to

$$w_{nov} := \Phi([\mu^n]) \in \Lambda[[U]] \otimes A,$$

where Φ is the HKR isomorphism. By the 'naturality' property of the HKR map (see Definition 2.5.1), w_{nov} is the image of the element

$$w \otimes 1 \in (R[[U]] \otimes A) \otimes_R \Lambda$$

under the natural map

$$(R[[U]] \otimes A) \otimes_R \Lambda \to \Lambda[[U]] \otimes A,$$

where w is the image of $[\mu^n]$ in $R[[U]] \otimes A$ under the HKR map.

Now, by the grading computations in the proof of Lemma 2.5.13, w has the form

$$w = f_1(T)u_1 \ldots u_n + f_2(T)\sum_{j=1}^{n} r_j u_j^n \in R[[u_1, \ldots, u_n]],$$

where $f_1, f_2 \in \mathbb{C}[[T]]$, and we recall that $T := r_1 \ldots r_n$. By the definition of 'type A', we have $f_1(0) = 1$ and $f_2(0) = 1$. It follows that

$$w_{nov} = f_1(r^n)u_1 \ldots u_n + f_2(r^n)\sum_{j=1}^{n} r u_j^n \in \Lambda[[u_1, \ldots, u_n]].$$

In particular, w_{nov} lies in $\Lambda[[U]]$, i.e., there are no non-trivial polyvector field terms appearing. Therefore, as we saw in the proof of Lemma 2.5.13,

$$\delta_{n-1}(-) = [\Phi(\mu^n), -] = \iota_{dw_{nov}}(-)$$

gives the Koszul complex for the sequence

$$\frac{\partial w_{nov}}{\partial u_1}, \ldots, \frac{\partial w_{nov}}{\partial u_n} \in \Lambda[[U]].$$

Now we show that this sequence is regular. This follows because w_{nov} has an isolated singularity at the origin. To see this, observe that we have relations

$$\begin{aligned} f_1(r^n)\frac{u_1 \ldots u_n}{u_j} &\equiv -nrf_2(r^n)u_j^{n-1} \\ \Rightarrow \frac{u_1 \ldots u_n}{u_j} &\equiv rf(r)u_j^{n-1}, \end{aligned}$$

in the ring

$$\Lambda[[u_1, \ldots, u_n]]/(\partial_1 w_{nov}, \ldots, \partial_n w_{nov}),$$

where

$$f(r) = -n\frac{f_2(r^n)}{f_1(r^n)} \in \mathbb{C}[[r^n]],\ f(0) \neq 0.$$

Taking the product of these relations gives

$$(u_1 \dots u_n)^{n-1} \equiv r^n f(r)^n (u_1 \dots u_n)^{n-1},$$

and hence that

$$(u_1 \dots u_n)^{n-1} \equiv 0,$$

because $1 + \mathcal{O}(r)$ is invertible in Λ. Returning to the original relation, we have

$$(rf(r)u_j^n)^{n-1} = (u_1 \dots u_n)^{n-1} = 0,$$

and hence sufficiently high powers of each generator u_j vanish (recalling $f \neq 0$, so $f \in \Lambda^*$ is invertible). Therefore w_{nov} has an isolated singularity at 0, so the sequence is regular, so the cohomology of the Koszul complex is the Jacobian ring.

Now we observe that

$$\boldsymbol{q}_{1*}\boldsymbol{p}_1^*\Lambda[[U]]$$

is generated, as a Λ-algebra, by the elements

$$u_1 \dots u_n, u_1^n, \dots, u_n^n,$$

and each of these has degree $2 \in \mathbb{Z}$. To see this, note that for $u^{\boldsymbol{b}}$ to be in $\boldsymbol{p}_1^*\Lambda[[U]]$, its degree must be in the image of Y_n^n in Y_1^n. From the definition, this means that

$$(|\boldsymbol{b}|, -\boldsymbol{b}) = (t + 2(n-1)|\boldsymbol{m}|, -n\boldsymbol{m}) - q(2(1-n), y_{[n]})$$

for some $t, q \in \mathbb{Z}$ and $\boldsymbol{m} \in \mathbb{Z}^n$, or in other words, setting $s = |\boldsymbol{b}|$ for the length,

$$\begin{aligned} s &= t + 2(n-1)|\boldsymbol{m}| + 2(n-1)q, \\ \boldsymbol{b} &= n\boldsymbol{m} + qy_{[n]}. \end{aligned}$$

It is not hard to show that, since $\boldsymbol{b} \geq \boldsymbol{0}$, we can arrange that $\boldsymbol{m} \geq \boldsymbol{0}$ and $q \geq 0$, so $u^{\boldsymbol{b}}$ is a product of the generators $u_1 \dots u_n, u_1^n, \dots, u_n^n$, as claimed. Furthermore, one can show that $s + t = 2q + 2|\boldsymbol{m}|$, where $s = |\boldsymbol{b}|$ is the length and t is the $\mathbb{Z}$-grading. It follows that the degree of each generator is 2, when regarded as an element of Hochschild cohomology.

It follows that the E_n page of our spectral sequence is given by

$$\Lambda[[u_1, \dots, u_n]]/(\partial_1 w_{nov}, \dots, \partial_n w_{nov}) \cap \Lambda[[u_1 \dots u_n, u_1^n, \dots, u_n^n]].$$

From the relations in the Jacobian ring, we have

$$u_1 \dots u_n = rf(r)u_j^n \text{ where } f(r) \neq 0,$$

so we only need a single generator $\alpha := u_1 \dots u_n$, and furthermore this generator satisfies

$$\alpha^{n-1} = 0,$$

as we showed above. It is easy to check (for example using Gröbner bases) that α^{n-2} does not lie in the ideal generated by the partial derivatives $\partial_j w_{nov}$. It follows that the E_n page of the spectral sequence is given by

$$\Lambda[\alpha]/\alpha^{n-1}.$$

Because α has degree $2 \in \mathbb{Z}$, E_n is graded in even degrees so all subsequent differentials in the spectral sequence vanish, so the spectral sequence is regular. Since the length filtration is bounded above and complete, it follows by [27, Theorem 5.5.10] that the spectral sequence converges to the Hochschild cohomology. Thus the E_n page is isomorphic to the associated graded algebra of the Hochschild cohomology; but since it is one-dimensional in each degree, it is in fact isomorphic to the Hochschild cohomology. This completes the proof. □

Now we observe that the element

$$r_j \frac{\partial \mu^*}{\partial r_j} \in CC^2_{\boldsymbol{G}^n_n}\left(\widetilde{\mathcal{A}}\right)$$

is a Hochschild cochain (this follows by applying $r_j \partial / \partial r_j$ to the A_∞ relation $\mu \circ \mu = 0$). Hence it defines an element in $HH^*(\widetilde{\mathcal{A}})$. We denote by β the element of $HH^*(\widetilde{\mathcal{A}}_{nov})$ that is the image of

$$\left(r_j \frac{\partial \mu^*}{\partial r_j}\right) \otimes 1 \in HH^*\left(\widetilde{\mathcal{A}}\right) \otimes_R \Lambda$$

under the obvious map.

Lemma 2.5.19. *The element β lies in the $\tilde{\Gamma}^*_n$-equivariant part of $HH^*(\widetilde{\mathcal{A}}_{nov})$, and corresponds to $g \cdot \alpha$ for some invertible $g \in \Lambda^*$ under the isomorphism of Lemma 2.5.18.*

Proof. The fact that β is $\tilde{\Gamma}^*_n$-equivariant follows immediately from the fact that μ^* is. We recall from the proof of Lemma 2.5.18 that, firstly, we arrange that μ^2 is independent of r_j and, secondly, the image of $\mu^{\ge 3}$ under the HKR map to $R[[U]] \otimes A$ has the form

$$w = f_1(T) u_1 \ldots u_n + f_2(T) \sum_{j=1}^n r_j u_j^n.$$

It follows that the image of $r_j \partial \mu^* / \partial r_j$ under the HKR map has the form

$$r_j \frac{\partial w}{\partial r_j} = T f_1'(T) u_1 \ldots u_n + f_2(T) r_j u_j^n + T f_2'(T) \sum_{j=1}^n r_j u_j^n.$$

It follows from naturality of the HKR map that the image of $\beta = (r_j \partial \mu^* / \partial r_j) \otimes 1$ on the E_2 page of the spectral sequence has the form

$$r^n f_1'(r^n) u_1 \ldots u_n + f_2(r^n) r u_j^n + r^n f_2'(r^n) \sum_{j=1}^n r u_j^n.$$

We now recall from the proof of Lemma 2.5.18 that the E_n page of the spectral sequence is the Jacobian ring $\Lambda[[u_1, \ldots, u_n]]/(\partial_1 w_{nov}, \ldots, \partial_n w_{nov})$. We recall (from the proof of Lemma 2.5.18) that, in the Jacobian ring, we have relations

$$f_1(r^n) u_1 \ldots u_n \equiv -n f_2(r^n) r u_j^n,$$

and we set $\alpha := u_1 \ldots u_n$, so the image of β on the E_n page of the spectral sequence is equivalent to

$$\left(r^n f_1'(r^n) - \frac{1}{n} f_1(r^n) - r^n \frac{f_2'(r^n) f_1(r^n)}{f_2(r^n)}\right) \alpha = g(r^n) \alpha,$$

where $g(r^n) \in \mathbb{C}[[r^n]]$ and $g(0) \neq 0$.

Recalling that the spectral sequence degenerates at the E_n page, this completes the proof. □

2.6. **First-order deformations.** In this section, we will consider a very specific situation. Let $\mathcal{A} = (A, \mu_0)$ be a $\boldsymbol{G}$-graded minimal A_∞ algebra over $\mathbb{C}$, and V a $\boldsymbol{G}$-graded vector space. We consider the $\boldsymbol{G}$-graded $\mathbb{C}$-algebra

$$R = \mathbb{C}[V]/\mathfrak{m}^2,$$

where $\mathfrak{m} \subset \mathbb{C}[V]$ is the maximal ideal corresponding to 0. A $\boldsymbol{G}$-graded **first-order deformation** of $\mathcal{A}$ over V is an element

$$\mu \in CC^2_{\boldsymbol{G}}(A \otimes R, A \otimes R),$$

whose order-0 component agrees with μ_0, and such that $\mu \circ \mu = 0$. Equivalently, we have

$$\mu = \mu_0 + \mu_1,$$

where

$$\mu_1 \in CC^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes V^\vee)$$

is a Hochschild cocycle. Its class in $HH^2_{\boldsymbol{G}}(\mathcal{A}, \mathcal{A} \otimes V^\vee)$ is called the **deformation class** of the deformation. Two first-order deformations are A_∞ quasi-isomorphic if and only if their deformation classes coincide.

Let $\mathcal{C}$ be an A_∞ category over $\mathbb{C}$ with two quasi-isomorphic objects, which we call L_0 and L_1. We denote the A_∞ endomorphism algebras of the two objects by $\mathcal{A}_j := \mathrm{Hom}(L_j, L_j)$ for $j = 0, 1$. It is standard that $\mathcal{A}_0$ and $\mathcal{A}_1$ are A_∞ quasi-isomorphic.

Let $\mathcal{C}_j$ be the full subcategory with a single object L_j, for $j = 0, 1$. Then the inclusions

$$\mathcal{C}_0 \hookrightarrow \mathcal{C} \hookleftarrow \mathcal{C}_1$$

are quasi-equivalences. It follows by Morita invariance (see, for example, [3, Lemma 2.6]) that the restriction maps

$$HH^*_{\boldsymbol{G}}(\mathcal{C}_0, \mathcal{C}_0) \leftarrow HH^*_{\boldsymbol{G}}(\mathcal{C}, \mathcal{C}) \rightarrow HH^*_{\boldsymbol{G}}(\mathcal{C}_1, \mathcal{C}_1)$$

are isomorphisms.

Now suppose that we have a $\boldsymbol{G}$-graded first-order deformation of $\mathcal{C}$ over V. As we observed above, this is equivalent (up to quasi-isomorphism) to a choice of deformation class

$$[\mu_1] \in HH^2_{\boldsymbol{G}}(\mathcal{C}, \mathcal{C} \otimes V^\vee).$$

The deformation classes of the induced first-order deformations of the endomorphism algebras $\mathcal{A}_j$ are given by the images of $[\mu_1]$ under the restriction maps

$$HH^2_{\boldsymbol{G}}(\mathcal{C}, \mathcal{C} \otimes V^\vee) \rightarrow HH^2_{\boldsymbol{G}}(\mathcal{A}_j, \mathcal{A}_j \otimes V^\vee),$$

and hence they correspond under the isomorphism

$$HH^2_{\boldsymbol{G}}(\mathcal{A}_0, \mathcal{A}_0 \otimes V^\vee) \cong HH^2_{\boldsymbol{G}}(\mathcal{A}_1, \mathcal{A}_1 \otimes V^\vee)$$

induced by the quasi-isomorphism of $\mathcal{A}_0$ with $\mathcal{A}_1$.

3. The affine Fukaya category

In this section, we introduce the affine Fukaya category $\mathcal{F}(M)$ of an exact symplectic manifold, and explain its relationship with the exact Fukaya category as defined in [3], which we will denote by $\mathcal{F}'(M)$. The difference is essentially that the affine Fukaya category has less objects, but a richer grading structure.

3.1. **Grading data from the Lagrangian Grassmannian.** We recall some notions from [3, Chapters 11, 12]. Let M be a symplectic manifold. We denote by $\mathcal{G}M$ the bundle of Lagrangian subspaces of TM. Observe that, because we have a fibration

$$\begin{tikzcd}
\mathcal{G}_p M \arrow[r, hook] & \mathcal{G}M \arrow[d] \\
 & M,
\end{tikzcd}$$

there is an exact sequence

$$\ldots \to \pi_1(\mathcal{G}_p M) \to \pi_1(\mathcal{G}M) \to \pi_1(M) \to *.$$

We recall that

$$\pi_1(\mathcal{G}_p M) \cong \mathbb{Z}.$$

Assumption 3.1.1. For the purposes of this paper, we will only consider manifolds M such that $\pi_1(M)$ is abelian. The reason for this assumption is that we will use $\pi_1(M)$ to define a grading datum $\boldsymbol{G}(M)$, and the affine Fukaya category will be $\boldsymbol{G}(M)$-graded. However we have only set up the theory of grading data (Section 2) for abelian groups. One should be able to define a sensible theory of non-abelian grading data, and apply it to study Fukaya categories of exact symplectic manifolds M with non-abelian fundamental group. On the other hand, when one studies the the relative Fukaya category (Section 5), it becomes absolutely necessary to consider abelian grading data. In this case, one should consider the universal **abelian** cover of $\mathcal{G}M$, rather than the universal cover. Since these issues do not concern us in this paper, we ignore them by making this assumption.

In particular, Assumption 3.1.1 implies that that $\pi_1(M) \cong H_1(M)$ and $\pi_1(\mathcal{G}M) \cong H_1(\mathcal{G}M)$.

Definition 3.1.2. We define a grading datum $\boldsymbol{G}(M)$:

$$\begin{tikzcd}
\mathbb{Z} \arrow[r, "f"] \arrow[d, equal] & Y(M) \arrow[r, "g"] \arrow[d, equal] & X(M) \arrow[r] \arrow[d, equal] & 0 \\
H_1(\mathcal{G}_p M) \arrow[r] & H_1(\mathcal{G}M) \arrow[r] & H_1(M) \arrow[r] & 0.
\end{tikzcd}$$

To define the sign morphism $\boldsymbol{\sigma}$, we must specify a map

$$\sigma : H_1(\mathcal{G}M) \to \mathbb{Z}_2.$$

To do this, we consider the real vector bundle $\mathcal{L} \to \mathcal{G}M$, whose fibre over a point is identified with the Lagrangian subspace at that point. The first Stiefel-Whitney class defines an element

$$w_1(\mathcal{L}) \in H^1(\mathcal{G}M; \mathbb{Z}_2),$$

and σ is defined by pairing with this element.

We will see that it is natural to define the Fukaya category as a $\boldsymbol{G}(M)$-graded category. That is because of the relationship between $\boldsymbol{G}(M)$ and index theory of Cauchy-Riemann operators in M, which we now explain.

We recall (from [31, Appendix C.3]) the definition of a **bundle pair** (Σ, E, F) over a Riemann surface with boundary Σ. It is a complex vector bundle $E \to \Sigma$ together with a totally real subbundle $F \subset E|_{\partial\Sigma}$. We recall that a bundle pair defines a Cauchy-Riemann operator

$$D : \Omega^0_F(\Sigma, E) \to \Omega^{0,1}_F(\Sigma, E),$$

whose zeroes are holomorphic sections of E whose boundary values lie in F. We recall also the **boundary Maslov index** of a bundle pair, which is an integer $\mu(\Sigma, E, F)$ such that the index of (an appropriate Sobolev-space version of) the Cauchy-Riemann operator D associated to the bundle pair (Σ, E, F) is

$$\mathrm{ind}(D) = n\chi(\Sigma) + \mu(\Sigma, E, F),$$

where n is the complex dimension of a fibre of E.

Now suppose we are given a map $u : \Sigma \to M$, together with a Lagrangian subbundle $F \subset u^*TM|_{\partial\Sigma}$. F defines a lift

$$\begin{tikzcd} & \mathcal{G}M \arrow[d] \\ \partial\Sigma \arrow[ur, "\rho"] \arrow[r, "u|_{\partial\Sigma}"'] & M, \end{tikzcd}$$

which defines a class $[\rho] \in H_1(\mathcal{G}M) = Y(M)$, which is a lift of

$$u|_{\partial\Sigma} = \partial u = 0 \in H_1(M) = X(M).$$

So by exactness of the sequence $\boldsymbol{G}(M)$, $[\rho]$ lies in the image of $\mathbb{Z}$.

Lemma 3.1.3. *We have*

$$[\rho] = f(\mu(\Sigma, u^*TM, F)),$$

in $Y(M)$, where $f : \mathbb{Z} \to Y(M)$ comes from the grading datum.

Proof. Define a decomposition of bundle pairs,

$$(\Sigma, u^*TM, F) = (\Sigma_1, u^*TM, F) \cup (\Sigma_2, u^*TM, F),$$

where Σ_1 is a small ball in the interior of Σ, Σ_2 is its complement, and the totally real subbundle of $u^*TM|_{\partial\Sigma_1}$ is defined by a lift

$$\rho' : \partial\Sigma_1 \to \mathcal{G}M,$$

chosen in such a way that there is a trivialization

$$(\Sigma_2, u^*TM, F) \cong (\Sigma_2, \Sigma_2 \times \mathbb{C}^n, \Sigma_2 \times \mathbb{R}^n).$$

It follows quickly from the properties of the boundary Maslov index (see [31, Theorem C.3.5]) that

$$\mu(\Sigma_2, u^*TM, F) = 0.$$

Then, by the composition property of the boundary Maslov index,

$$\mu(\Sigma, u^*TM, F) = \mu(\Sigma_1, u^*TM, F) + \mu(\Sigma_2, u^*TM, F) = \mu(\Sigma_1, u^*TM, F).$$

By our definition of (Σ_2, u^*TM, F), there is a lift

$$\begin{tikzcd} & \mathcal{G}M \arrow[d] \\ \Sigma_2 \arrow[ur, "P"] \arrow[r, "u|_{\Sigma_2}"'] & M, \end{tikzcd}$$

and P defines a homology between the class $[\rho]$ and the class $[\rho']$. If the ball Σ_1 is centred on a point p, then there is an obvious isomorphism

$$u^*TM|_{\Sigma_1} \cong \Sigma_1 \times T_pM,$$

so ρ' defines a class in $H_1(\mathcal{G}_pM) \cong \mathbb{Z}$, which is exactly equal to $\mu(\Sigma_1, u^*TM, F)$ (essentially from the definition of the Maslov class). The result follows. $\square$

3.2. **Anchored branes.** In this section, we will define the notion of an **anchored Lagrangian brane** in M. These will be the objects of the affine Fukaya category $\mathcal{F}(M)$ (when M is exact). First we recall the notion of a (non-anchored) graded Lagrangian brane from [3, Chapter 11], which is an object of $\mathcal{F}'(M)$.

If $2c_1(M) = 0$, and M is equipped with a quadratic complex volume form η, then we can construct a **phase map**

$$\alpha_M : \mathcal{G}M \to S^1.$$

Now if $i : L \to M$ is a Lagrangian immersion, there is a canonical lift of

$$\begin{tikzcd} & \mathcal{G}M \arrow[d] \\ L \arrow[ur, "i_*"] \arrow[r, "i"'] & M. \end{tikzcd}$$

Hence, given η, there is a map

$$\begin{aligned} \alpha_L : L &\to S^1, \\ \alpha_L &:= \alpha_M \circ i_*. \end{aligned}$$

A (non-anchored) **graded Lagrangian brane** in M is a compact embedded Lagrangian $L \subset M$, together with a lift $\alpha_L^{\#}$ of α_L to $\mathbb{R}$, and a Pin structure on L.

Now we introduce a new notion. Let

$$\pi : \widetilde{\mathcal{G}M} \to \mathcal{G}M$$

denote the universal cover of the manifold $\mathcal{G}M$.

Definition 3.2.1. An **anchored Lagrangian brane** $L^{\#}$ in M is a Lagrangian immersion $i : L \to M$ of a compact manifold L into M, together with a lift $i^{\#}$ as follows:

$$\begin{tikzcd} & \widetilde{\mathcal{G}M} \arrow[d, "\pi"] \\ L \arrow[ur, "i^{\#}"] \arrow[r, "i_*"'] & \mathcal{G}M, \end{tikzcd}$$

and a Pin structure. Note that we do **not** need a quadratic complex volume form to define the notion of an anchored Lagrangian brane.

We observe that there is a natural action of the covering group $\pi_1(\mathcal{G}M) \cong Y(M)$ on $\widetilde{\mathcal{G}M}$, and hence on anchored Lagrangian branes. We denote the action of $y \in Y(M)$ on $L^{\#}$ by $y \cdot L^{\#}$.

Now we explain the relationship between anchored Lagrangian branes and (non-anchored) graded Lagrangian branes. We observe that, given a quadratic complex volume form η_M, there exists a lift of the squared phase map

$$\begin{tikzcd}
\widetilde{\mathcal{G}}M \arrow[r, "\tilde{\alpha}_M"] \arrow[d] & \mathbb{R} \arrow[d] \\
\mathcal{G}M \arrow[r, "\alpha_M"] & S^1.
\end{tikzcd}$$

Definition 3.2.2. Given a quadratic volume form η_M, a lift $\tilde{\alpha}_M$ as above, and an embedded anchored Lagrangian brane $L^\# = (L, i^\#)$, we define a Lagrangian brane $\mathfrak{f}(L^\#)$ in M ($\mathfrak{f}$ for 'forgetting' the anchored structure), with the same underlying Lagrangian L and Pin structure, and

$$\alpha_L^\# = \tilde{\alpha}_M \circ i^\#.$$

3.3. **The affine Fukaya category.** Let M be an exact symplectic manifold with convex boundary, with $\pi_1(M)$ abelian (see Assumption 3.1.1). In this section, we define the **affine Fukaya category**, $\mathcal{F}(M)$. It will be a $\boldsymbol{G}(M)$-graded A_∞ category. The definition is very closely related to the definition of the affine Fukaya category $\mathcal{F}'(M)$, given in [3, Section 12], to which the reader is referred for all technical details. We will see that $\mathcal{F}(M)$ is essentially a full subcategory of $\mathcal{F}'(M)$ with a richer grading structure – in particular, the analytic details of defining moduli spaces of pseudoholomorphic disks to define the A_∞ structure maps are completely analogous.

Objects of $\mathcal{F}(M)$ are embedded anchored Lagrangian branes. For each pair of objects, we choose a Floer datum on M, and for all moduli spaces of boundary-punctured holomorphic disks with boundary components labelled by anchored Lagrangian branes, we make a consistent universal choice of perturbation data on M. We assume that the action of $Y(M)$ on anchored Lagrangian branes lifts to an action on Floer and perturbation data (i.e., if we change some of the boundary labels of a boundary-punctured holomorphic disk by the action of $Y(M)$, then the perturbation datum does not change).

We now define the $\boldsymbol{G}(M)$-graded morphism spaces in $\mathcal{F}(M)$. Now, given anchored Lagrangian branes $L_0^\#, L_1^\#$, we define the morphism space $CF^*(L_0^\#, L_1^\#)$ to be generated by paths $p : [0,1] \to M$ satisfying $p(0) \in L_0$, $p(1) \in L(1)$, which are flowlines of the Hamiltonian vector field associated with the corresponding Floer datum.

Given such a p, we define its grading $y \in Y(M)$ to be the unique element such that p lifts to a path from $L_0^\#$ to $y \cdot L_1^\#$ in $\widetilde{\mathcal{G}}M$, which has Maslov index $0 \in \mathbb{Z}$. To explain what this means, we observe that there is a commutative diagram

$$\begin{tikzcd}
\widetilde{\mathcal{G}}M \arrow[r] \arrow[d] & \widetilde{M} \arrow[d] \\
\mathcal{G}M \arrow[r] & M,
\end{tikzcd}$$

where $\widetilde{M} \to M$ is the universal cover of M. Thus, associated with any anchored Lagrangian brane $L^\#$ is a lift, $\widetilde{L}$, of L to $\widetilde{M}$. The fact that p must lift to $\widetilde{\mathcal{G}}M$ implies that it must lift to

$\widetilde{M}$; this already defines $y \in Y(M)$ up to addition of an element in the image of $\mathbb{Z} \to Y(M)$. Now we observe that the fibres of the bundle

$$\widetilde{\mathcal{G}}M \to \widetilde{M}$$

are the universal covers of the fibres of the Lagrangian Grassmannian $\mathcal{G}\widetilde{M}$. Thus, the anchored brane structures $L_0^{\#}, L_1^{\#}$ equip $\widetilde{L}_0, \widetilde{L}_1$ with the structure of 'abstract Lagrangian branes' (see [3, Section 12a]). Therefore, if the path p lifts to a path $\tilde{p}$ from $\widetilde{L}_0$ to $y \cdot \widetilde{L}_1$ in $\widetilde{M}$, then we can define the Maslov index i of any lift of $\tilde{p}$ to $\widetilde{\mathcal{G}}M$, and it is equal to the relative Maslov index of the abstract linear Lagrangian branes at either end of $\tilde{p}$. It is this index that we require to be 0. Given p, it is clear that we can find $y' \in Y(M)$ such that the path p lifts to a path $\tilde{p}$ from $\widetilde{L}_0$ to $y' \cdot \widetilde{L}_1$, but the Maslov index i may not be zero. However, we then necessarily have

$$y = y' - f(i),$$

so the $Y(M)$-grading of p is well-defined.

We define the A_∞ structure maps in $\mathcal{F}(M)$ by counting rigid pseudo-holomorphic disks in M. That is, given objects

$$L_0^{\#}, \ldots, L_s^{\#}$$

and morphisms

$$p_j \in CF^*(L_{j-1}^{\#}, L_j^{\#}) \text{ for } j = 1, \ldots, s$$

and

$$p_0 \in CF^*(L_0^{\#}, L_s^{\#}),$$

we define the coefficient of p_0 in $\mu^s(p_s, \ldots, p_1)$ to be the count of rigid pseudolomorphic disks in M with boundary conditions on L_j, asymptotic to the generators p_j.

We now explain why these structure maps are $\boldsymbol{G}(M)$-graded. Firstly, observe that the structure maps respect the action of $Y(M)$ on objects, because we chose the perturbation data to do so. From [3, Section 11l], we recall the definition of an orientation operator D_p corresponding to a generator p of $CF^*(L_0^{\#}, L_1^{\#})$. We lift p to a path

$$\rho : [0,1] \to \widetilde{\mathcal{G}}M$$

connecting $L_0^{\#}(p(0))$ to $y \cdot L_1^{\#}(p(1))$, where $y \in Y(M)$ is the degree of p. Now define a smooth, non-decreasing function $\psi : \mathbb{R} \to [0,1]$ such that $\psi(s) = 0$ for $s \ll 0$ and $\psi(s) = 1$ for $s \gg 0$. We consider the Hermitian vector bundle over the upper half plane $\mathbb{R} \times \mathbb{R}_{\geq 0}$, with fibre over (s,t) given by $T_{p(\psi(s))}M$. We introduce Lagrangian boundary conditions along the real axis, given by $\rho(\psi(s))$. These Lagrangian boundary conditions define a Cauchy-Riemann operator, which we denote by D_p. D_p is Fredholm, its index is equal to the relative Maslov index of the abstract Lagrangian branes at either end of p, which is 0 by the definition of y, and its determinant line is canonically isomorphic to the orientation line o_p of p.

Given a holomorphic disk u contributing to an A_∞ product μ^s, we denote the linearized operator of the perturbed holomorphic curve equation at u (with fixed domain S) by $D_{S,u}$. It is a Cauchy-Riemann operator on the trivial Hermitian vector bundle u^*TM over S. We can glue the orientation operators $D_{p_1}, \ldots, D_{p_s}$ and $D_{p_0}^{\vee}$ to $D_{S,u}$ along the strip-like ends to obtain a new Cauchy-Riemann operator over the closed disk. We denote this operator by $\overline{D}$. The gluing formula then implies that

$$\begin{aligned} i(\overline{D}) &= i(D_{S,u}) + i(p_1) + \ldots + i(p_s) + (n - i(p_0)) \\ &= i(D_{S,u}) + n, \end{aligned}$$

and there is a canonical isomorphism

$$\det(\overline{D}) \cong \det(D_u) \otimes o_{p_1} \otimes \ldots \otimes o_{p_s} \otimes o_{p_0}^{\vee}.$$

Now the Cauchy-Riemann operator $\overline{D}$ is given by a bundle pair (D^2, E, F), which is equivalent to a bundle pair (D^2, u^*TM, F), where $u : D^2 \to M$ is obtained from the original disk u (which had strip-like ends converging to the generators p_j) by gluing the orientation operators onto the ends. The boundary conditions for $\overline{D}$ define a map $\rho : \partial D^2 \to \mathcal{G}M$ which lifts the boundary map ∂u.

If we think of ρ as a map

$$\rho : [0,1] \to \mathcal{G}M$$

such that $\rho(0) = \rho(1)$ lies on L_0, then we have a lift

$$\begin{tikzcd} & \widetilde{\mathcal{G}}M \arrow[d] \\ {[0,1]} \arrow[ur, "\tilde{\rho}"] \arrow[r, "\rho"] & \mathcal{G}M. \end{tikzcd}$$

The boundary conditions L_j lift to $(y_1 + \ldots + y_j) \cdot L_j^{\#}$, and finally $\rho(1)$ lands on $(y_1 + \ldots + y_s - y_0) \cdot L_0^{\#}$. It follows that

$$[\rho] = -y_0 + \sum_{j=1}^{s} y_j.$$

Lemma 3.1.3 now implies that

$$\begin{aligned} f\left(i(\overline{D}) - n\chi\left(D^2\right)\right) &= -y_0 + \sum_{j=1}^{s} y_j \\ \Rightarrow f(i(D_{S,u})) &= -y_0 + \sum_{j=1}^{s} y_j \text{ in } Y(M). \end{aligned}$$

We now recall that, for the disk to be rigid, the **extended** linearized operator D_u (in which the modulus of the domain is allowed to vary, as well as the map) should have index zero. The dimension of the moduli space of disks with $s+1$ marked boundary points is $s-2$, so this means that

$$y_0 = f(2-s) + \sum_{j=1}^{s} y_j$$

in $Y(M)$. It follows that the affine Fukaya category is a $\boldsymbol{G}(M)$-graded A_∞ category (see Remark 2.3.12). We observe that the A_∞ associativity equations are satisfied, by the same argument as for $\mathcal{F}'(M)$ ([3, Proposition 12.3]).

We will now explain how $\mathcal{F}(M)$ is related to the $\mathbb{Z}$-graded exact Fukaya category $\mathcal{F}'(M)$, as defined in [3]. We recall that, to define $\mathbb{Z}$-gradings on $\mathcal{F}'(M)$, we require that $2c_1(M) = 0$ and equip η with a quadratic volume form η_M.

Recall from Definition 3.2.2 that, if we equip M with a quadratic volume form η_M, then we obtain a squared phase map

$$\alpha_M : \mathcal{G}M \to S^1,$$

and if we define a lift

$$\tilde{\alpha}_M : \widetilde{\mathcal{G}}M \to \mathbb{R},$$

then we obtain a forgetful map $\mathfrak{f}$ from anchored Lagrangian branes to (non-anchored) Lagrangian branes.

Now on the level of H_1, α_M induces a map

$$Y(M) \to \mathbb{Z}.$$

This defines a morphism of grading data, $\boldsymbol{p}^\eta : \boldsymbol{G}(M) \to \boldsymbol{G}_{\mathbb{Z}}$. It follows that $\boldsymbol{p}^\eta_* \mathcal{F}(M)$ is a $\mathbb{Z}$-graded A_∞ category. We have:

Lemma 3.3.1. *The forgetful map $\mathfrak{f}$ on objects extends to a fully faithful embedding of $\mathbb{Z}$-graded A_∞ categories,*

$$\mathfrak{f} : \boldsymbol{p}^\eta_* \mathcal{F}(M) \to \mathcal{F}'(M).$$

Remark 3.3.2. The image of this embedding consists of all (non-anchored) Lagrangian branes L such that the image of $H_1(L)$ in $H_1(M)$ is zero.

3.4. **Covers.** We explain how the affine Fukaya category behaves with respect to finite covers (essentially following [2, Section 8b]). Suppose that M, N are exact symplectic manifolds with convex boundary, with assumptions as in Section 3.3, and $\phi : N \to M$ is an exact symplectic covering (i.e., a covering such that the Liouville one-form on N is pulled back from that on M via ϕ). Then we have an induced covering

$$\begin{tikzcd}
\mathcal{G}N \arrow[r] \arrow[d, "\phi_*"'] & N \arrow[d, "\phi"] \\
\mathcal{G}M \arrow[r] & M,
\end{tikzcd}$$

and hence an injective morphism of grading data, $\boldsymbol{p} : \boldsymbol{G}(N) \to \boldsymbol{G}(M)$, given by

$$\begin{tikzcd}
H_1(\mathcal{G}_p N) \arrow[r] \arrow[d, equal, "\phi_*"'] & H_1(\mathcal{G}N) \arrow[r] \arrow[d, "\phi_*"'] & H_1(N) \arrow[r] \arrow[d, "\phi_*"'] & 0 \\
H_1(\mathcal{G}_{\phi(p)} M) \arrow[r] & H_1(\mathcal{G}M) \arrow[r] & H_1(M) \arrow[r] & 0.
\end{tikzcd}$$

Proposition 3.4.1. *There is a fully faithful embedding of $\boldsymbol{G}(N)$-graded categories,*

$$\boldsymbol{p}^*(\mathcal{F}(M)) \hookrightarrow \mathcal{F}(N).$$

Proof. We remark that the universal covers of $\mathcal{G}N$ and $\mathcal{G}M$ are isomorphic, so there is an obvious correspondence between anchored Lagrangian branes on N and on M. One can similarly set up a correspondence between morphism spaces and moduli spaces of pseudoholomorphic disks defining the A_∞ structure maps, and show that the gradings correspond, so the categories are strictly equivalent. $\square$

3.5. **The relative case.** Now we specialize to a particular type of exact symplectic manifold with corners.

Definition 3.5.1. A **Kähler pair** $(M, \boldsymbol{D})$ consists of:

- A smooth complex projective variety M, equipped with a positive holomorphic line bundle $\mathcal{L}$ with a Hermitian metric, so that the curvature of $\mathcal{L}$ defines a Kähler form ω, with $[\omega] = c_1(\mathcal{L})$;
- A tuple $\boldsymbol{D} = (D_1, D_2, \ldots, D_k)$ of smooth irreducible divisors $D_j \subset M$ with normal crossings, each corresponding to some positive power d_j of the bundle $\mathcal{L}$.

That is, for each $j = 1, 2, \ldots, k$, we have a section

$$h_j \in \Gamma(M, \mathcal{L}^{\otimes d_j})$$

such that

$$D_j = \{h_j = 0\}.$$

We furthermore assume that

- $\pi_1(M) = 0$;
- $\pi_1(M \setminus \boldsymbol{D})$ is abelian;
- $k \geq n + 1$, where k is the number of divisors and n is the complex dimension of M.

We define the **affine part**,

$$M \setminus \boldsymbol{D} := M \setminus \bigcup_j D_j.$$

We equip $M \setminus \boldsymbol{D}$ with a Kähler potential

$$h := \frac{\sum_{j=1}^k \log\left(\|h_j\|^2\right)}{\sum_{j=1}^k d_j},$$

so that $\alpha = -dh \circ J_0$ is a Liouville one-form (i.e., $\omega = d\alpha$), and α is convex at infinity (h is exhausting and bounded below on $M \setminus \boldsymbol{D}$).

Example 3.5.2. We consider the Fermat hypersurfaces,

$$M_a^n := \left\{ \sum_{j=1}^n z_j^a = 0 \right\} \subset \mathbb{CP}^{n-1},$$

with the ample divisors

$$D_j := \{z_j = 0\}$$

for $j = 1, \ldots, n$.

Now recall that there is a grading datum $\boldsymbol{G}(M \setminus \boldsymbol{D})$ associated to $M \setminus \boldsymbol{D}$, with exact sequence

$$\mathbb{Z} \to H_1(\mathcal{G}(M \setminus \boldsymbol{D})) \to H_1(M \setminus \boldsymbol{D}) \to 0.$$

We will denote

$$\boldsymbol{G}(M, \boldsymbol{D}) := \boldsymbol{G}(M \setminus \boldsymbol{D})$$

in the relative case, and write the exact sequence as

$$\mathbb{Z} \to \widetilde{Y}(M, \boldsymbol{D}) \to \widetilde{X}(M, \boldsymbol{D}) \to 0.$$

We now introduce a pseudo-grading datum $\boldsymbol{H}(M, \boldsymbol{D})$, whose associated grading datum is $\boldsymbol{G}(M, \boldsymbol{D})$. Compare [32, Section 5].

If we set U to be a neighbourhood of the union of the divisors $\boldsymbol{D}$ in M, and V to be $M \setminus \boldsymbol{D}$, then part of the Mayer-Vietoris long exact sequence for reduced homology gives

$$H_2(M) \to H_1(U \cap V) \to H_1(U) \oplus H_1(V) \to H_1(M) \to 0,$$

which becomes

$$H_2(M) \to \mathbb{Z}\langle y_1, \ldots, y_k \rangle \to H_1(M \setminus \boldsymbol{D}) \to 0,$$

since we are assuming $H_1(M) = 0$. Here $y_j \in H_1(U \cap V)$ is the class of a meridian loop around divisor D_j.

Definition 3.5.3. We define a pseudo-grading datum $\boldsymbol{H}(M, \boldsymbol{D})$, as follows: the exact sequence is

$$\begin{tikzcd}
Z(M, \boldsymbol{D}) \arrow[r, "f"] \arrow[d, equal] & Y(M, \boldsymbol{D}) \arrow[r, "g"] \arrow[d, equal] & X(M, \boldsymbol{D}) \arrow[r] \arrow[d, equal] & 0 \\
H_2(M; \mathbb{Z}) \arrow[r, "f"] & \mathbb{Z}\langle y_1, \ldots, y_k \rangle \arrow[r, "g"] & H_1(M \setminus \boldsymbol{D}; \mathbb{Z}) \arrow[r] & 0,
\end{tikzcd}$$

where we define

$$\begin{aligned} f(u) &= \sum_{i=1}^{k} (u \cdot D_i) y_i, \text{ and} \\ g(y_i) &= \text{the class of a loop around } D_i. \end{aligned}$$

We define the element $c \in \mathrm{Hom}(Z, \mathbb{Z})$ to be given by $2c_1(TM) \in H^2(M)$.

We now define a morphism of pseudo-grading data, $\boldsymbol{p} : \boldsymbol{H}(M, \boldsymbol{D}) \to \boldsymbol{G}(M, \boldsymbol{D})$. We define $p_Z : H_2(M) \to \mathbb{Z}$ to be given by $2c_1(TM)$, and p_X to be the identity. We define the map d to be 0. To define the map

$$p_Y : \mathbb{Z}\langle y_1, \ldots, y_k \rangle \to H_1(\mathcal{G}(M \setminus \boldsymbol{D})),$$

it is sufficient to define the action on the generators y_i. We denote the image of y_i by $\bar{y}_i$. To define $\bar{y}_i$, we consider a disk

$$u_i : (D^2, \partial D^2) \to (M, M \setminus \boldsymbol{D})$$

such that $u_i \cdot D_j = \delta_{ij}$. We trivialize the symplectic vector bundle $u_i^* TM$, and choose a lift of ∂u_i:

$$\begin{tikzcd}
 & \mathcal{G}M \arrow[d] \\
S^1 \arrow[ur, "\bar{y}_i"] \arrow[r, "\partial u_i"'] & M,
\end{tikzcd}$$

so that the boundary Maslov index

$$\mu(D^2, u_i^* TM, \bar{y}_i) = 0.$$

This defines the element $p_Y(y_i) := \bar{y}_i \in H_1(\mathcal{G}(M \setminus \boldsymbol{D}))$.

Lemma 3.5.4. *The diagram*

$$\begin{tikzcd}
H_2(M) \arrow[r] \arrow[d, "2c_1"'] & \mathbb{Z}\langle y_1, \dots, y_k\rangle \arrow[r] \arrow[d, "p_Y"'] & H_1(M \setminus \boldsymbol{D}) \arrow[r] \arrow[d, equal] & 0 \\
\mathbb{Z} \arrow[r] & H_1(\mathcal{G}(M \setminus \boldsymbol{D})) \arrow[r] & H_1(M \setminus \boldsymbol{D}) \arrow[r] & 0
\end{tikzcd}$$

commutes.

Proof. The only non-trivial thing to check is the commutativity of the left square. Suppose that $u : \Sigma \to M$ is a surface representing a homology class in $H_2(M)$, and intersecting the divisors D_j transversely. One side of the square maps

$$u \mapsto \sum_{i=1}^{k} (u \cdot D_i)\bar{y}_i,$$

while the other maps

$$u \mapsto f(2c_1(u)).$$

We consider the bundle pair (Σ, E, ϕ) with empty boundary, simply given by the complex vector bundle $E := u^*(TM)$. Its Maslov index is $\mu(\Sigma, E, \phi) = 2c_1(u)$. We now define a decomposition of this bundle pair: $\Sigma = \Sigma_1 \cup \Sigma_2$, where Σ_1 is a union of small balls around each of the intersection points of u with divisors D_i, and Σ_2 is the rest of Σ. We define the Lagrangian boundary conditions along $\partial\Sigma_1$ by requiring that the bundle pair over each ball around a single intersection point has boundary Maslov index zero. Then the composition property for bundle pairs (see [31, Appendix C.3]) says that

$$2c_1(u) = \mu(\Sigma, E, \phi) = \mu(\Sigma_1, E, F) + \mu(\Sigma_2, E, F) = \mu(\Sigma_2, E, F).$$

We note that the boundary conditions we have associated to a small ball around an intersection point of u with the divisor D_i define a map

$$S^1 \to \mathcal{G}(M \setminus \boldsymbol{D})$$

representing the class $\bar{y}_i$, by definition. Because u maps Σ_2 into $M \setminus \boldsymbol{D}$, it follows from Lemma 3.1.3 and the previous argument that

$$f(2c_1(u)) = f(\mu(\Sigma_2, E, F)) = \sum_{i=1}^{k} (u \cdot D_i)\bar{y}_i,$$

which proves that the left square commutes. □

Corollary 3.5.5. *The grading datum corresponding to the pseudo-grading datum $\boldsymbol{H}(M, \boldsymbol{D})$ is*

$$\boldsymbol{G}(\boldsymbol{H}(M, \boldsymbol{D})) \cong \boldsymbol{G}(M, \boldsymbol{D}).$$

Now suppose that we equip M with a meromorphic n-form η (i.e., an $(n, 0)$-form), whose zeroes and poles lie along the divisors D_j. Then we obtain a quadratic complex volume form η^2 on $M \setminus \boldsymbol{D}$, and recall that this defines a morphism $\boldsymbol{p}^\eta : \boldsymbol{G}(M, \boldsymbol{D}) \to \boldsymbol{G}_{\mathbb{Z}}$, allowing us to equip our category with a $\mathbb{Z}$-grading.

Lemma 3.5.6. *The morphism of grading data $\boldsymbol{p}^\eta$ is induced by the morphism of pseudo-grading data defined by*

$$\begin{aligned} p_Y : \mathbb{Z}\langle y_1, \dots, y_k\rangle &\to \mathbb{Z}, \\ p_Y(y_j) &= 2p_j, \end{aligned}$$

where p_j is the order of the pole of η along divisor D_j.

Proof. Follows essentially from the definition of the boundary Maslov index, see [31, Theorem C.3.5, 'Normalization' property]. □

Remark 3.5.7. If M is Calabi-Yau, then it admits a nowhere-vanishing holomorphic volume form η, so there is a canonical morphism of grading data

$$\boldsymbol{G}(M, \boldsymbol{D}) \to \boldsymbol{G}_{\mathbb{Z}},$$

which is induced by the zero morphism of pseudo-grading data, in accordance with Lemma 3.5.6.

Lemma 3.5.8. *The pseudo-grading datum associated to the Fermat hypersurfaces with coordinate divisors, $(M^n_a, \boldsymbol{D})$ (see Example 3.5.2) is*

$$\boldsymbol{H}(M^n_a, \boldsymbol{D}) \cong \boldsymbol{H}^n_a,$$

where $\boldsymbol{H}^n_a$ is the pseudo-grading datum of Example 2.1.13.

Proof. Follows from the fact that $H_2(M^n_a) \cong \mathbb{Z}$, generated by the class of a line $[P]$, that

$$[P] \cdot D_j = 1$$

for all j, and that

$$c_1([P]) = n - a.$$

□

Definition 3.5.9. Suppose that $(N, \boldsymbol{E})$ and $(M, \boldsymbol{D})$ are Kähler pairs (each with k divisors), and $\boldsymbol{a} = (a_1, \ldots, a_k)$ is a tuple of positive integers. An ***a*-branched cover** of Kähler pairs,

$$\phi : (N, \boldsymbol{E}) \to (M, \boldsymbol{D}),$$

is a branched cover $\phi : N \to M$ which maps divisor E_j to D_j, and has branching of order a_j along divisor E_j (and no branching anywhere else).

Example 3.5.10. There is an $(a, \ldots, a)$-branched cover of Fermat hypersurfaces

$$\begin{aligned} \phi_a : (M^n_a, \boldsymbol{D}) &\to (M^n_1, \boldsymbol{D}), \\ \phi_a([z_1 : \ldots : z_n]) &= [z_1^a : \ldots : z_n^a]. \end{aligned}$$

Lemma 3.5.11. *Let $\phi : (N, \boldsymbol{E}) \to (M, \boldsymbol{D})$ be an $\boldsymbol{a}$-branched cover. The unbranched cover*

$$\phi : N \setminus \boldsymbol{E} \to M \setminus \boldsymbol{D}$$

induces an injective morphism of grading data

$$\boldsymbol{p} : \boldsymbol{G}(N, \boldsymbol{E}) \to \boldsymbol{G}(M, \boldsymbol{D}),$$

as in Section 3.4. This morphism of grading data is induced by a morphism of pseudo-grading data

$$\tilde{\boldsymbol{p}} : \boldsymbol{H}(N, \boldsymbol{E}) \to \boldsymbol{H}(M, \boldsymbol{D}),$$

where

$$\begin{aligned} \tilde{p}_Y(y_j) &= a_j y_j, \text{ and} \\ d(y_j) &= 2(1 - a_j) \end{aligned}$$

Proof. It suffices to prove that

$$p_Y(0 \oplus y_j) = (2(1-a_j) \oplus a_j y_j).$$

This follows easily from the definition, and the local form

$$(z_1, z_2, \ldots, z_n) \mapsto (z_1^{a_j}, z_2, \ldots, z_n)$$

of ϕ near divisor E_j. □

Corollary 3.5.12. *If* $\phi_n : (M_n^n, \boldsymbol{D}) \to (M_1^n, \boldsymbol{D})$ *is the* $(n, \ldots, n)$*-branched cover of Fermat hypersurfaces introduced in Example 3.5.10, then the induced morphism of grading data*

$$\boldsymbol{p} : \boldsymbol{G}(M_n^n, \boldsymbol{D}) \to \boldsymbol{G}(M_1^n, \boldsymbol{D})$$

Coincides with the morphism

$$\boldsymbol{p}_1 : \boldsymbol{G}_n^n \to \boldsymbol{G}_1^n$$

of Lemma 2.1.14 (recall also Lemma 3.5.8).

4. Moduli Spaces of Disks

In this section we introduce the various moduli spaces of pseudo-holomorphic disks that we will need to define the versions of the Fukaya category that we will consider.

4.1. Moduli spaces of holomorphic spheres and disks.

Definition 4.1.1. Given an unordered set $\boldsymbol{E}$, with $|\boldsymbol{E}| \geq 3$, we define $\mathcal{R}_0(\boldsymbol{E})$, the moduli space of holomorphic spheres with distinct marked points q_e indexed by $e \in \boldsymbol{E}$, up to biholomorphism preserving marked points.

Definition 4.1.2. Given an ordered tuple $\boldsymbol{L} = (L_0, \ldots, L_d)$, a **disk with boundary labels** $\boldsymbol{L}$ is a disk with $d+1$ distinct boundary marked points, $\zeta_0, \zeta_1, \ldots, \zeta_d$, with the boundary component between ζ_i and ζ_{i+1} labelled L_i (understood modulo $d+1$).

We define three types of moduli spaces of disks:

Definition 4.1.3. Given a tuple $\boldsymbol{L}$, and a set $\boldsymbol{E}$, with $|\boldsymbol{L}| + 2|\boldsymbol{E}| \geq 3$, we define $\mathcal{R}(\boldsymbol{L}, \boldsymbol{E})$ to be the moduli space of holomorphic disks S with boundary labels $\boldsymbol{L}$, together with distinct internal marked points q_e indexed by $e \in \boldsymbol{E}$. We consider these objects up to biholomorphism preserving all marked points.

Definition 4.1.4. Given a tuple $\boldsymbol{L}$, we define $\mathcal{R}_1(\boldsymbol{L}) := \mathcal{R}(\boldsymbol{L}, \{1\})$, the moduli space of holomorphic disks S with boundary labels $\boldsymbol{L}$ and a single interior marked point q.

Definition 4.1.5. Given a tuple $\boldsymbol{L}$, we define $\mathcal{R}_2(\boldsymbol{L}) \subset \mathcal{R}(\boldsymbol{L}, \{1,2\})$ to be the moduli space of holomorphic disks S with Lagrangian labels $\boldsymbol{L}$, together with interior marked points q_1, q_2, such that there is a biholomorphism of S with the unit disk $\{|w| \leq 1\} \subset \mathbb{C}$ sending

$$\begin{aligned} \zeta_0 &\mapsto -i \\ q_1 &\mapsto -t \\ q_2 &\mapsto t \end{aligned}$$

for some $t \in (0,1) \subset \mathbb{R}$ (see Figure 4.1.1).



FIGURE 4.1.1. The moduli space $\mathcal{R}_2(\boldsymbol{L})$, where $|\boldsymbol{L}| = 6$.

Given a point r in one of these moduli spaces, we denote by S_r the corresponding (marked) disk, with all boundary marked points removed.

4.2. **Deligne-Mumford compactifications.** We make a universal choice of strip-like and cylindrical ends for each of these moduli spaces. We denote by $\overline{\mathcal{R}}_0(\boldsymbol{E})$, $\overline{\mathcal{R}}(\boldsymbol{L}, \boldsymbol{E})$, $\overline{\mathcal{R}}_1(\boldsymbol{L})$, $\overline{\mathcal{R}}_2(\boldsymbol{L})$ the Deligne-Mumford compactifications of these moduli spaces by stable spheres and disks. We now describe these compactifications.

The Deligne-Mumford compactification of $\mathcal{R}_0(\boldsymbol{E})$ consists of stable trees of spheres. Boundary strata are indexed by stable trees T, with semi-infinite edges indexed by $\boldsymbol{E}$. We denote by $V(T)$ the set of vertices of T, and $E(T)$ the set of edges of T. A tree is called **stable** if each vertex has valence ≥ 3. For each vertex v of T, we denote by $\boldsymbol{E}_v$ the set of edges of T incident to v. The boundary stratum indexed by T is

$$\mathcal{R}_0^T(\boldsymbol{E}) := \prod_v \mathcal{R}_0(\boldsymbol{E}_v).$$

Points in this stratum correspond to trees of sphere bubbles, with semi-infinite edges corresponding to marked points, and finite edges corresponding to nodes (see [31, Section D.3, Figure 4]). The Deligne-Mumford (or Grothendieck-Knudsen) compactification, as a set, is the



Figure 4.2.1. If $\boldsymbol{L}$ is some tuple, then a k-leafed stable tree T is said to have labels $\boldsymbol{L}$ if the connected components of $\mathbb{R}^2 \setminus T$ are labeled by the elements of $\boldsymbol{L}$, in order. In this figure, $\boldsymbol{L} = (L_0, L_0, L_0, L_1, L_2, L_2, L_1, L_0, L_3)$. A labeling $\boldsymbol{L}$ of T induces a labeling $\boldsymbol{L}_v$ of the regions surrounding each vertex v. In this figure, the induced labeling of the regions surrounding the uppermost vertex v is $\boldsymbol{L}_v = (L_0, L_1, L_2, L_2)$.

union of all such strata. It is a smooth manifold (see [31, Section D.5]). The codimension of the stratum indexed by T is $2(|V(T)| - 1)$.

Definition 4.2.1. A **directed d-leafed planar tree** is a directed d-leafed tree T embedded in $\mathbb{R}^2$. It consists of the following data:

- a finite set of vertices $V(T)$;
- a set of d semi-infinite **outgoing edges**;
- a single semi-infinite **incoming edge**, connected to a vertex $v \in V(T)$ called the **root** of T;
- a set $E(T)$ of internal edges.

A vertex is allowed to have zero outgoing edges, but must always have exactly one incoming edge. We say that a vertex $v \in V(T)$ is **stable** if it has ≥ 2 outgoing edges, and **semi-stable** if it has ≥ 1 outgoing edges. Given a tuple $\boldsymbol{L}$, we say that T **has labels $\boldsymbol{L}$** if the connected components of $\mathbb{R}^2 \setminus T$ are labeled by the elements of $\boldsymbol{L}$, in order. A labeling of T induces a labeling $\boldsymbol{L}_v$ of the regions surrounding each vertex $v \in V(T)$ (see Figure 4.2.1).

The Deligne-Mumford compactification of $\overline{\mathcal{R}}(\boldsymbol{L}, \boldsymbol{E})$ consists of stable trees of disk and sphere bubbles with appropriate markings (see [33, Section 2.3]). It is a smooth manifold with corners. Each boundary stratum is indexed by a directed tree T, together with a directed planar subtree T_L with labels $\boldsymbol{L}$. We denote $T_E := T \setminus T_L$. We require that the semi-infinite edges of T_E are indexed by $\boldsymbol{E}$. For each vertex $v \in T_L$, we have a labeling $\boldsymbol{L}_v$ as above, and denote by $\boldsymbol{E}_v$ the set of edges incident to v in T_E. For each vertex $v \in T_E$, we denote by $\boldsymbol{E}_v$ the set of edges incident to v in T_E. We require that the tree is stable, in the sense that for each vertex $v \in T_L$,

$$|\boldsymbol{L}_v| + 2|\boldsymbol{E}_v| \geq 3,$$

while for each vertex $v \in T_E$,

$$|\boldsymbol{E}_v| \geq 3.$$

The tree T corresponds to the stratum

$$\mathcal{R}^T(\boldsymbol{L}, \boldsymbol{E}) \cong \prod_{v \in V(T_L)} \mathcal{R}(\boldsymbol{L}_v, \boldsymbol{E}_v) \times \prod_{v \in V(T_E)} \mathcal{R}_0(\boldsymbol{E}_v).$$

Points in this stratum correspond to nodal disks, with semi-infinite edges of T_L corresponding to boundary marked points, finite edges of T_L corresponding to boundary nodes, semi-infinite edges of T_E corresponding to internal marked points, and finite edges of T_E corresponding to internal nodes. The codimension of this stratum is

$$|V(T_L)| + 2|V(T_E)| - 1.$$

The boundary strata of $\overline{\mathcal{R}}_2(\boldsymbol{L})$ fall into three types (we have illustrated the codimension-1 part of each stratum in Figure 4.2.2):

- strata indexed by directed planar trees T with boundary labels $\boldsymbol{L}$, together with a distinguished vertex v_1, so that all vertices other than possibly v_1 have valence ≥ 3; these correspond to codimension-$(|V(T)| - 1)$ strata
$$\mathcal{R}_2^{1,T}(\boldsymbol{L}) \cong \mathcal{R}_2(\boldsymbol{L}_{v_1}) \times \prod_{v \in V(T) \setminus \{v_1\}} \mathcal{R}(\boldsymbol{L}_v)$$
which consist of nodal disks glued together in the obvious way (see Figure 2(a));
- another set of strata indexed by directed planar trees T with boundary labels $\boldsymbol{L}$, together with a distinguished vertex v_1, so that all vertices other than possibly v_1 have valence ≥ 3; these correspond to codimension-$|V(T)|$ strata
$$\mathcal{R}_2^{2,T}(\boldsymbol{L}) \cong \mathcal{R}_1(\boldsymbol{L}_{v_1}) \times \prod_{v \in V(T) \setminus \{v_1\}} \mathcal{R}(\boldsymbol{L}_v)$$
which consist of nodal disks glued in the obvious way, together with a sphere with three marked points, two of which are the marked points q_1, q_2 and one of which is a node, identified to the internal marked point q in the disk coming from the factor $\mathcal{R}_1(\boldsymbol{L}_{v_1})$ (see Figure 2(b));
- strata indexed by directed planar trees T with boundary labels $\boldsymbol{L}$ and two (different) distinguished vertices v_1, v_2, so that all vertices other than possibly v_1 and v_2 have valence ≥ 3, and the branch of T containing v_1 lies strictly to the left of the branch



(a) The codimension-1 part of $\mathcal{R}_2^{1,T}(\boldsymbol{L})$.

(b) The codimension-1 part of $\mathcal{R}_2^{3,T}(\boldsymbol{L})$.



(c) The codimension-1 part of $\mathcal{R}_2^{2,T}(\boldsymbol{L})$.

FIGURE 4.2.2. The codimension-1 boundary components of $\mathcal{R}_2(\boldsymbol{L})$, where $|\boldsymbol{L}| = 6$.

containing v_2; these correspond to codimension-$(|V(T)| - 2)$ strata

$$\mathcal{R}_2^{3,T}(\boldsymbol{L}) \cong \mathcal{R}_1(\boldsymbol{L}_{v_1}) \times \mathcal{R}_1(\boldsymbol{L}_{v_2}) \times \prod_{v \in V(T) \setminus \{v_1, v_2\}} \mathcal{R}(\boldsymbol{L}_v)$$

consisting of nodal disks glued together in the obvious way, where the internal marked point in the disk coming from the factor $\mathcal{R}_1(\boldsymbol{L}_{v_j})$ corresponds to the marked point q_j, for $j = 1, 2$ (see Figure 2(c)).

4.3. **Moduli spaces of pseudoholomorphic disks.** Let $(M, \boldsymbol{D})$ be a Kähler pair (see Definition 3.5.1). Thus, M is a Kähler manifold, and $\boldsymbol{D} = D_1 \cup \ldots \cup D_k$ is a union of smooth ample divisors with normal crossings. In this section we will define moduli spaces of pseudoholomorphic disks mapping into M.

Definition 4.3.1. Let $\boldsymbol{F}$ be a finite set, and let

$$\ell : \boldsymbol{F} \to [k]$$

be a function from $\boldsymbol{F}$ to the set $[k] := \{1, \ldots, k\}$ indexing the divisors $D_1, \ldots, D_k$. We call such a function a **labelling** of $\boldsymbol{F}$. Recalling the definition of the pseudo-grading datum $\boldsymbol{H}(M, \boldsymbol{D})$ from Definition 3.5.3, we denote

$$\boldsymbol{d}(\ell) := \sum_{j=1}^{k} |\ell^{-1}(j)| y_j \in Y_{\geq 0}$$

(where $Y_{\geq 0} := \mathbb{Z}_{\geq 0}\langle y_1, \ldots, y_k \rangle$). If $\boldsymbol{a} = (a_1, \ldots, a_k)$ is a tuple of positive integers, then we define

$$\boldsymbol{d}_{\boldsymbol{a}}(\ell) := \sum_{j=1}^{k} a_j |\ell^{-1}(j)| y_j \in Y_{\geq 0}.$$

We denote $\mathbf{1} = (1, \ldots, 1)$ (k copies), so that $\boldsymbol{d} = \boldsymbol{d}_{\mathbf{1}}$.

Definition 4.3.2. Let $\boldsymbol{E}$ and $\boldsymbol{F}$ be finite sets, $|\boldsymbol{E}| + |\boldsymbol{F}| \geq 3$. Let $\ell : \boldsymbol{F} \to [k]$ be a labelling of $\boldsymbol{F}$. We define

$$\mathcal{R}_0(\boldsymbol{E}, \ell) := \mathcal{R}_0(\boldsymbol{E} \sqcup \boldsymbol{F}).$$

Definition 4.3.3. Given a tuple of objects $\boldsymbol{L}$, finite sets $\boldsymbol{E}, \boldsymbol{F}$ such that $|\boldsymbol{L}| + 2|\boldsymbol{E}| + 2|\boldsymbol{F}| \geq 3$, and a labelling $\ell : \boldsymbol{F} \to [k]$, we define the moduli space

$$\mathcal{R}(\boldsymbol{L}, \boldsymbol{E}, \ell) := \mathcal{R}(\boldsymbol{L}, \boldsymbol{E} \sqcup \boldsymbol{F}).$$

For each pair of objects in the affine Fukaya category $\mathcal{F}(M \setminus \boldsymbol{D})$, we choose a Floer datum (see [3, Section 8e]). We make a universal choice of perturbation data (in the sense of [3, Section 9h]) on each of the moduli spaces $\mathcal{R}_0(\boldsymbol{E}, \ell)$, $\mathcal{R}(\boldsymbol{L}, \boldsymbol{E}, \ell)$, $\mathcal{R}_1(\boldsymbol{L})$ and $\mathcal{R}_2(\boldsymbol{L})$. Note that the choice of perturbation data on the moduli spaces $\mathcal{R}_0$, $\mathcal{R}$ may be different for different labellings ℓ, even though they have the same number of boundary components and internal marked points. Also note that we are choosing Floer and perturbation data which are defined on all of M, not just on $M \setminus \boldsymbol{D}$.

We require that

- the Hamiltonian part of each perturbation datum is 0 on the moduli spaces $\mathcal{R}_0(\boldsymbol{E})$;
- the Hamiltonian part of each perturbation datum vanishes, with its first derivatives, along each divisor D_j;
- the almost-complex structure part of each perturbation datum makes each divisor D_j an almost-complex manifold;
- on the strip-like ends, the perturbation datum agrees with the associated Floer datum;
- the choices of perturbation data are consistent with respect to the Deligne-Mumford compactifications outlined in Section 4.1, in the sense of [3, Section 9i];

- the choices of perturbation data are invariant under shifts of the anchored Lagrangian branes by the covering group action
$$\pi_1(\mathcal{G}(M \setminus \boldsymbol{D})) \cong \widetilde{Y}(M, \boldsymbol{D}),$$
as was the case for the affine Fukaya category (see Section 3.3).

We will use the shorthand
$$u \cdot \boldsymbol{D} := \sum_{j=1}^{k} (u \cdot D_j) y_j \in Y(M, \boldsymbol{D})$$
(see Definition 3.5.3 for the definition of $Y(M, \boldsymbol{D})$), where $u \cdot D_j$ denotes the topological intersection number, for any class $u \in H_2(M)$ or $H_2(M, M \setminus \boldsymbol{D})$. Note that with our choice of perturbation data, any pseudoholomorphic curve u intersects the divisors D_j positively, so
$$u \cdot \boldsymbol{D} \in Y(M, \boldsymbol{D})_{\geq 0}$$
if u is a pseudo-holomorphic disk or sphere that is not contained inside any of the divisors D_j.

Definition 4.3.4. We define an element of the moduli space $\mathcal{M}_0(\boldsymbol{E}, \ell)$ to be a pair (r, u), where r is an element of $\mathcal{R}_0(\boldsymbol{E}, \ell)$ and $u : S_r \to M$ is a smooth map, such that:

- u satisfies the perturbed holomorphic curve equation;
- $u(q_f) \in D_{\ell(f)}$ for each $f \in \boldsymbol{F}$;
- $u \cdot \boldsymbol{D} = \boldsymbol{d}(\ell)$

(see [3, Equations (8.9) and (9.17)] for the perturbed holomorphic curve equation). There is an evaluation map
$$\boldsymbol{ev} : \mathcal{M}_0(\boldsymbol{E}, \ell) \to M^{\boldsymbol{E}}.$$
If $\boldsymbol{E}$ is empty we may omit it from the notation, and we will also write $\mathcal{M}_0(\boldsymbol{E}, \boldsymbol{d})$ for $\mathcal{M}_0(\boldsymbol{E}, \ell)$ where $\boldsymbol{d} = \boldsymbol{d}(\ell)$.

Remark 4.3.5. Suppose that $u \in \mathcal{C}_0(\boldsymbol{E}, \ell)$ is not contained in divisor D_j. Then our assumptions on the perturbation data ensure that u intersects D_j in isolated points with positive multiplicity. Since each marked point q_f with $\ell(f) = j$ contributes at least 1 to $u \cdot D_j$, our requirement that $u \cdot \boldsymbol{D} = \boldsymbol{d}(\ell)$ ensures that u intersects D_j only at the marked points q_f with $\ell(f)$, and each intersection has multiplicity 1.

Definition 4.3.6. Given a tuple $\boldsymbol{L} = (L_0^{\#}, \ldots, L_s^{\#})$ of anchored Lagrangian branes, an **associated set of generators** is a tuple $\boldsymbol{p} = (p_0, \ldots, p_s)$ where p_j is a generator of $CF^*(L_j^{\#}, L_{j-1}^{\#})$ for $j \geq 1$, and p_0 is a generator of $CF^*(L_0^{\#}, L_s^{\#})$.

Definition 4.3.7. Given a tuple of objects $\boldsymbol{L}$ with associated generators $\boldsymbol{p}$, finite sets $\boldsymbol{E}, \boldsymbol{F}$ such that $|\boldsymbol{L}| + 2|\boldsymbol{E}| + 2|\boldsymbol{F}| \geq 3$, and a labelling $\ell : \boldsymbol{F} \to [k]$ as in Definition 4.3.1, we define an element of the moduli space $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell)$ to be a pair (r, u), where r is an element of $\mathcal{R}(\boldsymbol{L}, \boldsymbol{E}, \ell)$ and $u : S_r \to M$ is a smooth map, such that:

- u satisfies the perturbed holomorphic curve equation, with Lagrangian boundary conditions given by the labels $\boldsymbol{L}$;
- u is asymptotic to the generators $\boldsymbol{p}$ along the corresponding strip-like ends;
- $u(q_f) \in D_{\ell(f)}$ for each $f \in \boldsymbol{F}$;

- $u \cdot \boldsymbol{D} = \boldsymbol{d}(\ell)$.

(see [3, Equation (8.10)] for the definition of 'asymptotic to the generators $\boldsymbol{p}$'). There is an evaluation map

$$\boldsymbol{ev} : \mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell) \to M^{\boldsymbol{E}}.$$

If $\boldsymbol{E}$ is empty, we may omit it from the notation, and we will also write $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \boldsymbol{d})$ instead of $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell)$, where $\boldsymbol{d} = \boldsymbol{d}(\ell)$.

By the same reasoning as in Remark 4.3.5, every intersection point of an element $u \in \mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell)$ with a divisor D_j is a marked point (and the intersection has multiplicity 1).

Definition 4.3.8. For a tuple of 2 Lagrangian labels $\boldsymbol{L}$ with associated generators $\boldsymbol{p}$, we define $\mathcal{M}(\boldsymbol{p}, 0)$, the set of holomorphic strips with boundary conditions on $\boldsymbol{L}$, intersection number 0 with the divisors $\boldsymbol{D}$, translation-invariant perturbation coming from the corresponding Floer datum, asymptotic to the generators $\boldsymbol{p}$, modulo translation by $\mathbb{R}$ (see [3, Equation (8.8)]).

Given a holomorphic curve with an internal marked point, we define the notion of 'tangency to a divisor to order k' at the marked point, in accordance with [34]:

Definition 4.3.9. Suppose we are given:

- a Riemann surface S with an internal marked point $q \in S$;
- a perturbed holomorphic curve $u : S \to M$;
- a choice of divisor $D_j \subset M$;
- an integer $k \geq 1$.

We say that u is **tangent to D_j at q to order k** if

- $u(q) \in D_j$;
- all partial derivatives of u at q of order $\leq k$ lie inside the tangent space TD_j.

We remark that this does not depend on the choice of coordinates.

When $k = 0$, this is the same thing as a point constraint $u(q) \in D_j$. For $k \geq 1$, one should think of the curve u having 'ramification of order $k+1$' about the divisor D_j. We can find local holomorphic coordinates $(z_1, \ldots, z_n)$ for M near $u(q)$, such that $u(q)$ corresponds to the origin, and D_j corresponds to $\{z_1 = 0\}$, and a local holomorphic coordinate z for S near q such that q corresponds to the origin. If we assume that the almost-complex structure part of the perturbation datum is equal to the standard complex structure along the cylindrical end associated to the marked point q, and the Hamiltonian part of the perturbation datum vanishes, then u takes the form $(u_1(z), \ldots, u_n(z))$ in these coordinates, where $u_j(z)$ are holomorphic functions, and u_1 has a zero of order $\geq k+1$ at the origin. It follows that the point q contributes at least $k+1$ to the intersection number of u with D_j.

Definition 4.3.10. Suppose that $\boldsymbol{a} = (a_1, \ldots, a_k)$ is a tuple of k positive integers. We define the moduli spaces $\mathcal{M}_0(\boldsymbol{E}, \ell, \boldsymbol{a})$ and $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$ in exactly the same way as we did $\mathcal{M}_0(\boldsymbol{E}, \ell)$ and $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell)$, with the following exceptions:

- For each $f \in \boldsymbol{F}$, we require u to be tangent to $D_{\ell(f)}$ at q_f to order $a_{\ell(f)} - 1$;
- We require that $u \cdot \boldsymbol{D} = d_{\boldsymbol{a}}(\ell)$.

In particular, we have isomorphisms

$$\mathcal{M}_0(\boldsymbol{E}, \ell) \cong \mathcal{M}_0(\boldsymbol{E}, \ell, \mathbf{1})$$

and

$$\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell) \cong \mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell, \mathbf{1}).$$

Note that, because each marked point q_f contributes $\geq a_{\ell(f)}$ to $u \cdot D_{\ell(f)}$, elements $u \in \mathcal{M}_0(\boldsymbol{E}, \ell, \boldsymbol{a})$ do not intersect any of the divisors D_j anywhere other than at marked points q_f, where they intersect with multiplicity $a_{\ell(f)}$, and similarly for $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$.

Definition 4.3.11. Let $\boldsymbol{L}$ be a tuple of Lagrangians with associated generators $\boldsymbol{p}$, a a positive integer, and $j \in [k]$. We define $\boldsymbol{F}$ to be a set with a single element, and ℓ a labelling which assigns j to this element. We let $\boldsymbol{a}$ be any tuple such that $a_j = a$. Then we define

$$\mathcal{M}_1(\boldsymbol{p}, j, a) := \mathcal{M}(\boldsymbol{p}, \ell, \boldsymbol{a}),$$

the moduli space of pseudoholomorphic disks with a single internal marked point, which is tangent to divisor D_j to order $a - 1$. Note that we're not defining anything new; this is just convenient notation for us to have.

Definition 4.3.12. Given a tuple of Lagrangians $\boldsymbol{L}$ with associated generators $\boldsymbol{p}$, a positive integer a, together with a choice of divisor D_j, we define an element of the moduli space $\mathcal{M}_2(\boldsymbol{p}, j, a)$ to consist of the following data:

- a point $r \in \mathcal{R}_2(\boldsymbol{L})$;
- a smooth map $u : S_r \to M$,

such that:

- u satisfies the perturbed holomorphic curve equation;
- u is asymptotic to the generators $\boldsymbol{p}$ along the strip-like ends;
- $u \cdot \boldsymbol{D} = (a + 1)e_j$;
- u is tangent to D_j at q_1 to order $a - 1$, and to D_j at q_2 to order 0.

Note that, as before, elements $u \in \mathcal{M}_2(\boldsymbol{p}, j, a)$ do not intersect the divisors D_i for $i \neq j$, and intersect D_j only at q_1 (with multiplicity a) and q_2 (with multiplicity 1).

4.4. **Indices and orientations.** Each of the moduli spaces of pseudoholomorphic curves we have defined can be defined as the set of zeroes of a smooth section of a Banach bundle over a Banach manifold of maps (more precisely, the moduli space can be covered by such). We follow [3, Chapters 8 and 9] in defining the functional analytic framework, with modifications following [34, Section 6] to take into account the 'orders of tangency' restrictions. This just means we have to use $W^{k+1,p}$ maps rather than $W^{1,p}$, so that we can make sense of 'derivatives of order $\leq k$'.

The linearization of this smooth section defines a Fredholm operator. The moduli spaces are said to be regular when the linearization is surjective everywhere. When they are regular, the moduli spaces are smooth manifolds, with dimension given by the index of the Fredholm operator. In this section, we will outline the calculation of this dimension.

Lemma 4.4.1. *Let $\boldsymbol{L}$ be a tuple of anchored Lagrangian branes with associated generators $\boldsymbol{p}$, and u an element of $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell)$. Let $\tilde{y}_j \in \widetilde{Y}(M, \boldsymbol{D})$ be the degree of p_j. Then the index of the extended linearized operator D_u at u satisfies*

$$f(i(D_u) - 2|\boldsymbol{E}| + 2 - s) = -\tilde{y}_0 + \sum_{j=1}^{s} \tilde{y}_j + p_Y(\boldsymbol{d}(\ell)),$$

and there is a canonical identification of orientation lines

$$o_{p_0} \cong \det(D_u) \otimes o_{p_1} \otimes \ldots \otimes o_{p_s}.$$

Proof. Each path p_j lifts to a path from $L^{\#}_{j-1}$ to $\tilde{y}_j \cdot L^{\#}_j$ in $\widetilde{\mathcal{G}}(M \setminus \boldsymbol{D})$, which we use to define orientation operator D_{p_j} whose index is 0.

We denote the linearized operator of the perturbed holomorphic curve equation at u (with fixed domain S) by $D_{S,u}$. We glue the orientation operators D_{p_j} to $D_{S,u}$ along the strip-like ends to obtain a bundle pair over the closed disk, and hence a Cauchy-Riemann operator $\overline{D}$. The gluing formula implies that

$$\begin{aligned} i(\overline{D}) &= i(D_{S,u}) + i(D_{p_1}) + \ldots + i(D_{p_s}) + (n - i(D_{p_0})) \\ &= i(D_{S,u}) + n, \end{aligned}$$

and there is a canonical isomorphism

$$\det(\overline{D}) \cong \det(D_{S,u}) \otimes o_{p_1} \otimes \ldots \otimes o_{p_s} \otimes o_{p_0}^{\vee}.$$

As in Section 3.3, this bundle pair defining $\overline{D}$ is equivalent to a bundle pair (D^2, u^*TM, ρ), where $\rho : \partial D^2 \to \mathcal{G}(M \setminus \boldsymbol{D})$ which lifts the boundary map ∂u, and hence has index

$$i(\overline{D}) = n + \mu(D^2, u^*TM, \rho).$$

As in Section 3.3, we can compute the homology class

$$[\rho] = -\tilde{y}_0 + \sum_{j=1}^{s} \tilde{y}_j.$$

As in the proof of Lemma 3.5.4, we define a decomposition of the bundle pair (D^2, u^*TM, ρ) into two bundle pairs: (Σ_1, u^*TM, ρ'), consisting of a union of small balls surrounding each of the points q_f for $f \in \boldsymbol{F}$, with boundary conditions given by $\bar{y}_{\ell(f)}$, and $(\Sigma_2, u^*TM, \rho \cup \rho')$ the complement of Σ_1. Then the decomposition property of the boundary Maslov index, together with Lemma 3.1.3, say that

$$\begin{aligned} \mu(D^2, u^*TM, \rho) &= \mu(\Sigma_1, u^*TM, \rho) + \mu(\Sigma_2, u^*TM, \rho \cup \rho') \\ &= \mu(\Sigma_2, u^*TM, \rho \cup \rho') \\ \Rightarrow f(\mu(D^2, u^*TM, \rho)) &= [\rho] + [\rho'] \\ &= -\tilde{y}_0 + \sum_{j=1}^{s} \tilde{y}_j + p_Y(\boldsymbol{d}(\ell)). \end{aligned}$$

The result now follows, as

$$\begin{aligned} i(D_u) &= i(D_{S,u}) + \dim(\mathcal{R}(\boldsymbol{L},\boldsymbol{E})) \\ &= i(\overline{D}) - n + s - 2 + 2|\boldsymbol{E}| \\ &= \mu(D^2, u^*TM, \rho) + s - 2 + 2|\boldsymbol{E}|. \end{aligned}$$

The isomorphism of orientation lines follows after we fix an orientation for $\mathcal{R}(\boldsymbol{L},\boldsymbol{E})$. □

Lemma 4.4.2. *Suppose we have objects $\boldsymbol{L}$ with associated generators $\boldsymbol{p}$, a k-tuple $\boldsymbol{a} = (a_1, \ldots, a_k)$ of positive integers, and an element $u \in \mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$. Let $\tilde{y}_j \in \widetilde{Y}(M, \boldsymbol{D})$ be the degree of p_j. Then the index of the extended linearized operator D_u at u satisfies*

$$f\left(i(D_u) + 2 - s - 2|\boldsymbol{E}| - 2\left|\boldsymbol{d}(\ell) - \boldsymbol{d}_{\boldsymbol{a}}(\ell)\right|\right) = -\tilde{y}_0 + \sum_{j=1}^{s} \tilde{y}_j + p_Y(\boldsymbol{d}_{\boldsymbol{a}}(\ell)).$$

Proof. Follows from Lemma 4.4.1, where we observe that being tangent to $D_{\ell(f)}$ at q_f to order $a_{\ell(f)}$ imposes an additional $2(a_{\ell(f)} - 1)$-dimensional constraint on the disk, leading to the final term on the left-hand side, which is equal to

$$2 \sum_{f \in \boldsymbol{F}} 1 - a_{\ell(f)}.$$

□

We can perform similar calculations for $\mathcal{M}_0$, $\mathcal{M}_1$ and $\mathcal{M}_2$. We obtain:

Lemma 4.4.3. *If D_u is the extended linearized operator at $u \in \mathcal{M}_0(\boldsymbol{E}, \ell, \boldsymbol{a})$, then*

$$f(i(D_u) - 2n + 6 - 2|\boldsymbol{E}| - 2\left|\boldsymbol{d}(\ell) - \boldsymbol{d}_{\boldsymbol{a}}(\ell)\right|) = p_Y(\boldsymbol{d}_{\boldsymbol{a}}(\ell))).$$

Lemma 4.4.4. *If D_u is the extended linearized operator at $u \in \mathcal{M}_2(\boldsymbol{p}, j, a)$, then*

$$f(i(D_u) + 2a + 1 - s) = -\tilde{y}_0 + \sum_{l=1}^{s} \tilde{y}_l + p_Y((a+1)y_j),$$

and there is a canonical isomorphism of orientation lines as before. To clarify, recall that $\tilde{y}_l \in \widetilde{Y}(M, \boldsymbol{D})$ is the degree of p_l, but y_j is the jth generator of $Y(M, \boldsymbol{D})$.

One can prove that these moduli spaces are regular for generic choices of perturbation data, by essentially the same arguments as in [34] and [3, Section 9k]. Namely, for each map u in the Banach manifold of maps, one can choose the perturbation datum essentially arbitrarily on an open subset of the domain, and this is enough to achieve transversality. There are two exceptions: firstly, the moduli space of holomorphic strips that do not intersect the boundary divisors is defined using a translation-invariant perturbation data (see Definition 4.3.8). It is shown in [35, 36] that these moduli spaces are regular for generic choice of Floer data. The second exception is for moduli spaces of holomorphic spheres contained entirely within one of the divisors D_j: our assumptions on the perturbation data along the divisors make it impossible to guarantee regularity in this situation. We will explain in Section 4.5 why this is not a problem for our purposes.

4.5. **Gromov compactness.** We now describe Gromov compactifications of the moduli spaces we have defined. We observe that each moduli space has bounded energy, and standard Gromov compactness shows that any sequence in one of these moduli spaces has a subsequence which converges, up to sphere and disk bubbling (see [31] for spheres and [33] for disks). We would like to show that any sphere or disk that bubbles off must be stable, so that the Gromov compactification of our moduli spaces is regular. First we discuss the case of sphere bubbles.

Any non-constant sphere bubble u must have positive intersection with each divisor D_j. So unless it is contained inside one of the divisors, it has $\ge k \ge 3$ marked points (where k is the number of divisors). In particular, all such sphere bubbles have stable domain, so they are regular for generic choice of perturbation data.

The main difficulty occurs when there are sphere bubbles lying entirely inside one of the divisors D_j, because we restricted our perturbation data so that the Hamiltonian part vanishes along D_j, and the almost-complex structure component makes D_j an almost-complex submanifold. So we cannot guarantee regularity of these sphere bubbles, and we need separate arguments to deal with this case.

Definition 4.5.1. Let $\boldsymbol{E}, \boldsymbol{F}$ be finite sets, and $\ell : \boldsymbol{F} \to [k]$ a labelling. A stratum of $\overline{\mathcal{M}}_0(\boldsymbol{E}, \ell)$ is indexed by a tree T whose semi-infinite edges are indexed by $\boldsymbol{E} \sqcup \boldsymbol{F}$. For each vertex v, we denote by $\boldsymbol{F}_v$ the set of semi-infinite edges incident to v with index in $\boldsymbol{F}$, and by $\ell_v : \boldsymbol{F}_v \to [k]$ the induced labelling of $\boldsymbol{F}_v$. We denote by $\boldsymbol{E}_v$ the set of the remaining edges (finite or semi-infinite) incident to v. The tree is required to be **stable**, in the sense that for each vertex $v \in V(T)$ we have $|\boldsymbol{E}_v| + |\boldsymbol{F}_v| \ge 3$.

Definition 4.5.2. Given such a tree T, we define the corresponding stratum of the Gromov compactification. For each vertex $v \in V(T)$, we define

$$\mathcal{M}_0^T(v) := \mathcal{M}_0(\boldsymbol{E}_v, \ell_v).$$

We then define

$$(\Pi\mathcal{M})_0^T(\boldsymbol{E}, \ell) := \prod_{v \in V(T)} \mathcal{M}_0^T(v).$$

If $\boldsymbol{E}_{\text{int}}$ denotes the set of internal (finite) edges of T, then there is an evaluation map

$$\boldsymbol{ev}^T : (\Pi\mathcal{M})_0^T(\boldsymbol{E}, \ell) \to M^{\boldsymbol{E}_{\text{int}}} \times M^{\boldsymbol{E}_{\text{int}}}.$$

Note that each edge appears twice on the right-hand side, once for each of its endpoints. We define

$$\mathcal{M}_0^T(\boldsymbol{E}, \ell) := \left(\boldsymbol{ev}^T\right)^{-1}\left(\triangle^{\boldsymbol{T}}\right),$$

where

$$\triangle^{\boldsymbol{T}} \subset M^{\boldsymbol{E}_{\text{int}}} \times M^{\boldsymbol{E}_{\text{int}}}$$

denotes the diagonal.

Definition 4.5.3. As a set, we define

$$\overline{\mathcal{M}}_0(\boldsymbol{E}, \ell) := \coprod_T \mathcal{M}_0^T(\boldsymbol{E}, \ell).$$

We equip it with the Gromov topology.

Proposition 4.5.4. *If $|\boldsymbol{E}| \ge 1$, then the space $\overline{\mathcal{M}}_0(\boldsymbol{E}, \ell)$ is compact.*

Proof. Standard Gromov compactness (see [31]) says that any sequence in $\overline{\mathcal{M}}_0(\boldsymbol{E}, \ell)$ has a subsequence which Gromov-converges to a tree of nodal spheres. The space $\overline{\mathcal{M}}_0(\boldsymbol{E}, \ell)$, by definition, contains all **stable** nodal spheres. These are the nodal spheres such that each sphere bubble has ≥ 3 marked points. Thus, to prove the result, we must show that, if a non-constant sphere bubbles off in our moduli space, it must be stable.

Suppose that u is a non-constant pseudoholomorphic sphere bubble appearing in some nodal sphere which is the Gromov limit of a sequence in $\overline{\mathcal{M}}_0(\boldsymbol{E}, \ell)$. If u intersects a divisor D_j in an isolated point, then the point must be a marked point. This is because the intersection point persists in a neighbourhood of the nodal sphere in the Gromov topology, and any isolated intersection point of a sphere in $\mathcal{M}_0(\boldsymbol{E}, \ell)$ with a divisor D_j is a marked point, by Remark 4.3.5. Now u has positive intersection with each of the divisors D_j. Hence, if u has isolated intersection points with the divisors, it has ≥ 3 marked points where it intersects the divisors, and its domain is therefore stable.

If u does not have isolated intersection points with some divisor D_j, it must be contained in D_j by analytic continuation (and our assumptions on the perturbation data near the divisors).

Definition 4.5.5. If $K \subset [k]$, we denote

$$D_K := \bigcap_{j \in K} D_j.$$

Suppose that $u \subset D_K$, but has transverse intersections with all other divisors D_j. The dimension of D_K is $2n - 2|K|$ (by the normal crossings condition), so for the sphere to be non-constant we require that $|K| \leq n-1$. Then, because there are $\geq n+1$ divisors (by definition of a Kähler pair, see Definition 3.5.1), there remain ≥ 2 divisors D_j with which u has isolated intersections. The sphere u must intersect these divisors positively, and the intersections must be marked points. So u has ≥ 2 marked points coming from the corresponding intersections, as well as the marked point corresponding to the node (or to the marked point $q \in \boldsymbol{E}$), hence its domain is stable. $\square$

The virtual codimension of the stratum $\mathcal{M}_0^T(\boldsymbol{E}, \ell)$ in $\overline{\mathcal{M}}(\boldsymbol{E}, \ell)$ is $2(\#\text{ vertices of } T_E) - 1$. If every stratum is regular, then standard gluing theorems show that $\overline{\mathcal{M}}_0(\boldsymbol{E}, \ell)$ has the structure of a compact topological manifold with corners, and the actual codimension of each stratum is equal to the virtual codimension. Regularity of the stratum indexed by the tree T means that the moduli spaces $\mathcal{M}_0^T(v)$ are regular, and the evaluation map $\boldsymbol{ev}^T$ is transverse to the diagonal $\triangle^T$.

If all nodal spheres in the moduli spaces $\mathcal{M}_0^T(v)$ are transverse to the divisors $\boldsymbol{D}$, then the moduli space is regular for generic choice of perturbation data, as we can change the perturbation arbitrarily away from $\boldsymbol{D}$ (compare [31, Section 6.3]). However, it is possible that one of the holomorphic spheres appearing in this moduli space is contained in some divisor D_j. In that case, our assumptions on the perturbation data along the divisor D_j make it impossible to guarantee regularity of this moduli space in M.

However, with some additional assumptions on the perturbation data, one can show that, if the virtual dimension of this stratum of $\mathcal{M}_0^T(\boldsymbol{E}, \ell)$ is negative, then it is actually empty, even if there are sphere bubbles contained inside some of the divisors D_j. We can also show that transversality can still be achieved with these additional assumptions on the perturbation data. This suffices for our purposes of defining algebraic structures out of these moduli spaces.

Remark 4.5.6. We remark that this is the point at which we need the assumption (see Definition 3.5.1) that each divisor D_j is ample.

We now describe these assumptions:

Let $\boldsymbol{E}, \boldsymbol{F}$ be finite sets, and $\ell : \boldsymbol{F} \to [k]$ a labelling of $\boldsymbol{F}$. Given a subset $K \subset [k]$, let

$$\boldsymbol{F}^K := \{f \in \boldsymbol{F} : \ell(f) \notin K\},$$

and

$$\ell^K : \boldsymbol{F}^K \to \overline{K}$$

the restriction of ℓ. There is a forgetful map

$$\mathfrak{f}^K : \mathcal{R}_0(\boldsymbol{E}, \ell) \to \mathcal{R}_0(\boldsymbol{E}, \ell^K).$$

Condition 4.5.7. For each $K \subset [k]$, the perturbation data on $\mathcal{R}_0(\boldsymbol{E}, \ell)$ coincides with the pullback of the perturbation data on $\mathcal{R}_0(\boldsymbol{E}, \ell^K)$ by $\mathfrak{f}^K$, when restricted to D_K.

Observe that even in the presence of Condition 4.5.7, we can still perturb the almost-complex structure arbitrarily away from the divisors D_j, so our previous transversality arguments for pseudoholomorphic spheres which are not contained in a divisor D_j are unaffected by this additional assumption.

Definition 4.5.8. Given a moduli space $\mathcal{M}$ cut out locally by a Fredholm section of a Banach vector bundle, we denote by

$$\text{v.d.}(\mathcal{M})$$

the Fredholm index of the section, where 'v.d.' stands for 'virtual dimension': this is the expected dimension of the moduli space, and the actual dimension if it is regular.

Proposition 4.5.9. *Suppose $|\boldsymbol{E}| \geq 1$. For a generic choice of perturbation data satisfying Condition 4.5.7, any stratum $\mathcal{M}_0^T(\boldsymbol{E}, \ell)$ of $\overline{\mathcal{M}}_0(\boldsymbol{E}, \ell)$ whose virtual dimension is negative, is actually empty.*

Proof. For clarity, we will omit the $(\boldsymbol{E}, \ell)$ from the notation throughout the proof. Let us consider a stratum indexed by a fixed tree T. For the purposes of this proof, we will make T a directed tree (arbitrarily), and denote by $h(e) \in V(T)$ the head of the directed edge $e \in \boldsymbol{E}_{int}$, and by $t(e) \in V(T)$ the tail.

Recall that any holomorphic disk either intersects a divisor transversely, or is contained in it. For each vertex $v \in V(T)$, we define a stratification

$$\mathcal{M}_0^T(v) = \bigcup_{K \subset [k]} \mathcal{M}_0^{T,K}(v),$$

where

$$\mathcal{M}_0^{T,K}(v) := \{u_v \in \mathcal{M}_0^T(v) : u_v \subset D_K, u_v \not\subset D_j \text{ for } j \notin K\}$$

(recall Definition 4.5.5).

We now consider the manifold D_K, with divisors $D_j \cap D_K$ for $j \notin K$ (these are divisors are normal crossings in D_K, because of the normal crossings condition in M). We define the moduli space

$$\widetilde{\mathcal{M}}_0^{T,K}(v) := \left\{ u_v \in \mathcal{M}_0\left(\boldsymbol{E}_v, \ell_v^K\right) : u_v \subset D_K \right\}.$$

Assuming Condition 4.5.7 is satisfied, there is a forgetful map

$$\mathfrak{f}_v^{T,K} : \mathcal{M}_0^{T,K}(v) \to \widetilde{\mathcal{M}}_0^{T,K}(v),$$

obtained by forgetting the marked points with label $j \in K$. Furthermore, the evaluation map

$$\boldsymbol{ev}_v : \mathcal{M}_0^{T,K}(v) \to M^{\boldsymbol{E}_v}$$

factors as follows:

$$\begin{tikzcd}
\mathcal{M}_0^{T,K}(v) \arrow[r, "\boldsymbol{ev}_v"] \arrow[d, "\mathfrak{f}_v^{T,K}"'] & M^{\boldsymbol{E}_v} \\
\widetilde{\mathcal{M}}_0^{T,K}(v) \arrow[r, "\widetilde{\boldsymbol{ev}}_v^{T,K}"] & D_K^{\boldsymbol{E}_v}. \arrow[u, hook]
\end{tikzcd}$$

Lemma 4.5.10. *For generic choice of perturbation data satisfying Condition 4.5.7, $\widetilde{\mathcal{M}}_0^{T,K}(v)$ is a smooth manifold of dimension*

$$\dim\left(\widetilde{\mathcal{M}}_0^{T,K}(v)\right) \le \text{v.d.}_M\left(\mathcal{M}_0^T(v)\right) - 2|K|.$$

Proof. As we saw in the proof of Proposition 4.5.4, $|\boldsymbol{F}_v^K| \ge 2$ and $|\boldsymbol{E}_v| \ge 1$, so the domain of u_v is stable. Because we can perturb the restriction of the almost-complex structure part of the perturbation data to D_K essentially arbitrarily away from the divisors $D_j \cap D_K$, the moduli space $\widetilde{\mathcal{M}}_0^{T,K}(v)$ is regular in D_K for generic choice of perturbation data (observe that this is **not** necessarily the same as being regular in M). Its virtual dimension at u_v (hence its actual dimension, when regular) in D_K is

$$\text{v.d.}_{D_K}\left(\widetilde{\mathcal{M}}_0^{T,K}(v)\right) = 2(n - |K|) + 2|\boldsymbol{E}_v| + 2c_1(TD_K)(u_v) - 6.$$

Denoting the normal bundle of the divisor D_j by ND_j, we have

$$\begin{aligned}
c_1(TM)(u_v) &= c_1\left(TD_K \oplus \bigoplus_{j\in K} ND_j\right)(u_v) \\
&= c_1(TD_K)(u_v) + \sum_{j\in K} c_1(ND_j)(u_v) \\
&= c_1(TD_K)(u_v) + \sum_{j\in K} c_1\left(\mathcal{L}^{\otimes d_j}\Big|_{D_j}\right)(u_v) \\
&= c_1(TD_K)(u_v) + \sum_{j\in K} d_j\omega(u_v) \\
&\ge c_1(TD_K)(u_v),
\end{aligned}$$

because the symplectic area of a pseudoholomorphic sphere is non-negative. It follows that

$$\begin{aligned} \text{v.d.}_M\left(\mathcal{M}_0^T(v)\right) - 2|K| &= 2n + 2|\boldsymbol{E}_v| + 2c_1(TM)(u_v) - 6 - 2|K| \\ &\geq 2(n-|K|) + 2|\boldsymbol{E}_v| + 2c_1(TD_K)(u_v) - 6 \\ &= \text{v.d.}_{D_K}\left(\widetilde{\mathcal{M}}_0^{T,K}(v)\right). \end{aligned}$$

The actual dimension of $\widetilde{\mathcal{M}}_0^{T,K}(v)$ near u_v is generically the same as the virtual dimension in D_K, since it is generically regular in D_K. □

Thus, we have a stratification of

$$(\Pi\mathcal{M})_0^T(\boldsymbol{E},\ell) = \prod_v \mathcal{M}_0^T(v)$$

by smooth manifolds

$$(\Pi\mathcal{M})_0^{T,\boldsymbol{K}} := \prod_v \mathcal{M}_0^{T,K_v}(v),$$

where $\boldsymbol{K}$ denotes the choice of a subset $K_v \subset [k]$ for each $v \in V(T)$. We define

$$\left(\Pi\widetilde{\mathcal{M}}\right)_0^{T,\boldsymbol{K}} := \prod_{v \in V(T)} \widetilde{\mathcal{M}}_0^{T,K_v}(v)$$

and

$$D_{\boldsymbol{K}}^{\boldsymbol{E}_{int}} := \prod_{e \in \boldsymbol{E}_{int}} D_{K_{h(e)}} \times D_{K_{t(e)}},$$

and observe that the evaluation map $\boldsymbol{ev}^{T,\boldsymbol{K}}$ factors as follows:

$$\begin{tikzcd} \Pi\mathcal{M}_0^{T,\boldsymbol{K}} \arrow[r, "\boldsymbol{ev}^T"] \arrow[d, "\mathfrak{f}^{T,\boldsymbol{K}}"'] & M^{\boldsymbol{E}_{int}} \times M^{\boldsymbol{E}_{int}} \\ \Pi\widetilde{\mathcal{M}}_0^{T,\boldsymbol{K}} \arrow[r, "\widetilde{\boldsymbol{ev}}^{T,\boldsymbol{K}}"] & D_{\boldsymbol{K}}^{\boldsymbol{E}_{int}}, \arrow[u, hook] \end{tikzcd}$$

where $\mathfrak{f}^{T,\boldsymbol{K}}$ is a forgetful map.

Now, observe that the map

$$D_{\boldsymbol{K}}^{\boldsymbol{E}_{int}} \to M^{\boldsymbol{E}_{int}} \times M^{\boldsymbol{E}_{int}}$$

is **not** necessarily transverse to Δ^T, and in particular we cannot hope for the map

$$\Pi\widetilde{\mathcal{M}}_0^{T,\boldsymbol{K}} \to M^{\boldsymbol{E}_{int}} \times M^{\boldsymbol{E}_{int}}$$

to be transverse to Δ^T. However, we will explain that

$$\widetilde{\boldsymbol{ev}}^{T,\boldsymbol{K}} : \Pi\widetilde{\mathcal{M}}_0^{T,\boldsymbol{K}} \to D_{\boldsymbol{K}}^{\boldsymbol{E}_{int}}$$

is generically transverse to the diagonal

$$\Delta^{T,\boldsymbol{K}} := \Delta^T \cap D_{\boldsymbol{K}}^{\boldsymbol{E}_{int}}.$$

We have

$$\Delta^{T,\boldsymbol{K}} \cong \prod_{e \in \boldsymbol{E}_{int}} D_{K_{h(e)} \cup K_{t(e)}} \hookrightarrow \prod_{e \in \boldsymbol{E}_{int}} D_{K_{h(e)}} \times D_{K_{t(e)}},$$

where the inclusion is via the 'diagonal map' $p \mapsto (p,p)$. The map $\widetilde{\boldsymbol{ev}}^{T,\boldsymbol{K}}$ is then transverse to $\Delta^{T,\boldsymbol{K}}$ because, for any $e \in \boldsymbol{E}_{int}$, sphere bubbles inside $D_{K_{h(e)}}$ are generically regular, so we can perturb them to make the marked point move in any direction within $TD_{K_{h(e)}}$. Similarly for the sphere bubble in $D_{K_{t(e)}}$.

It follows that the manifold

$$\left(\widetilde{\boldsymbol{ev}}^{T,\boldsymbol{K}}\right)^{-1}\left(\Delta^{T,\boldsymbol{K}}\right)$$

is generically smooth of dimension

$$\begin{aligned}
\dim\left(\left(\widetilde{\boldsymbol{ev}}^{T,\boldsymbol{K}}\right)^{-1}\left(\Delta^{T,\boldsymbol{K}}\right)\right) &= \dim\left(\Pi\widetilde{\mathcal{M}}_0^{T,\boldsymbol{K}}\right) - \operatorname{codim}\left(\Delta^{T,\boldsymbol{K}}\right) \\
&= \sum_v \dim\left(\widetilde{\mathcal{M}}_0^{T,K_v}(v)\right) - \sum_{e\in\boldsymbol{E}_{int}} \left(2n - 2|K_{h(e)} \cap K_{t(e)}|\right) \\
&\leq \sum_{v\in V(T)} \left(\text{v.d.}\left(\mathcal{M}_0^T(v)\right) - 2|K_v|\right) - \sum_{e\in\boldsymbol{E}_{int}} 2n - 2|K_{h(e)} \cap K_{t(e)}| \\
&= \text{v.d.}\left(\Pi\mathcal{M}_0^T\right) - 2n|\boldsymbol{E}_{int}| - 2\sum_{v\in V(T)} |K_v| + 2\sum_{e\in\boldsymbol{E}_{int}} |K_{h(e)} \cap K_{t(e)}| \\
&\leq \text{v.d.}\left(\mathcal{M}_0^T\right) - 2\sum_{e\in\boldsymbol{E}_{int}} |K_{h(e)}| - |K_{h(e)} \cap K_{t(e)}| \\
&\leq \text{v.d.}\left(\mathcal{M}_0^T\right),
\end{aligned}$$

where the second-last line follows as each $v \in V(T)$ is the head of at most one edge $e \in \boldsymbol{E}_{int}(T)$.

In particular, if the expected dimension of $\mathcal{M}_0^T$ is negative, then the manifold is empty. It follows that the intersection of the subspace

$$\boldsymbol{ev}^{-1}\left(\Delta^T\right) \cap \Pi\mathcal{M}_0^{T,\boldsymbol{K}} = \left(\mathfrak{f}^{T,\boldsymbol{K}}\right)^{-1}\left(\left(\widetilde{\boldsymbol{ev}}^{T,\boldsymbol{K}}\right)^{-1}\left(\Delta^{T,\boldsymbol{K}}\right)\right)$$

is empty, for each $\boldsymbol{K}$.

This completes the proof. □

We have an analogous result for the moduli spaces $\overline{\mathcal{M}}_0(\boldsymbol{E}, \ell, \boldsymbol{a})$. Now we describe the Gromov compactification of the moduli spaces $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$.

Definition 4.5.11. Let $\boldsymbol{L}$ be a tuple of elements of the affine Fukaya category, $\boldsymbol{p}$ an associated set of generators, $\boldsymbol{E}, \boldsymbol{F}$ finite sets, $\ell : \boldsymbol{F} \to [k]$ a labelling, such that $|\boldsymbol{L}| + 2|\boldsymbol{E}| + 2|\boldsymbol{F}| \geq 2$, and $\boldsymbol{a}$ a k-tuple of positive integers. A stratum of $\overline{\mathcal{M}}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$ is indexed by an object $\boldsymbol{T}$, where $\boldsymbol{T}$ consists of the following data:

- A tree T, together with a directed planar subtree T_L with labels $\boldsymbol{L}$;
- An indexing of the semi-infinite edges of $T_E := T \setminus T_L$ by $\boldsymbol{E} \sqcup \boldsymbol{F}$;
- For each edge e of T_L, a choice of generator $p_e \in CF^*(L_{r(e)}, L_{l(e)})$, where $L_{r(e)}, L_{l(e)}$ are the Lagrangian labels to the right and left of e respectively, such that the generators are given by $\boldsymbol{p}$ for the external edges.

For each vertex $v \in V(T)$, we denote by $\boldsymbol{F}_v$ the set of semi-infinite edges in T_E that are incident to v and have index in $\boldsymbol{F}$, and by $\ell_v : \boldsymbol{F}_v \to [k]$ the labelling induced by ℓ. We denote by $\boldsymbol{E}_v$ the remaining edges (finite or semi-infinite) in T_E that are incident to v. For each vertex $v \in T_L$, we denote by $\boldsymbol{L}_v$ the tuple of Lagrangians labelling the regions surrounding v, and by $\boldsymbol{p}_v$ the set of chosen generators for the edges adjacent to v. The tree T is required to be **semi-stable**, in the sense that for each vertex $v \in V(T_L)$ we have

$$|\boldsymbol{L}_v| + 2|\boldsymbol{E}_v| + 2|\boldsymbol{F}_v| \geq 2,$$

while for each vertex $v \in V(T_E)$,

$$|\boldsymbol{E}_v| + |\boldsymbol{F}_v| \geq 3.$$

Definition 4.5.12. Given such an object $\boldsymbol{T}$, we define the corresponding stratum of the Gromov compactification. For vertices $v \in T_E$, we define

$$\mathcal{M}^{\boldsymbol{T}}(v) := \mathcal{M}_0(\boldsymbol{E}_v, \ell_v, \boldsymbol{a}).$$

For $v \in T_L$, we define

$$\mathcal{M}^{\boldsymbol{T}}(v) := \mathcal{M}(\boldsymbol{p}_v, \boldsymbol{E}_v, \ell_v, \boldsymbol{a}).$$

Now, letting $\boldsymbol{E}_{\text{int}}$ denote the internal (finite) edges of T_E, we have an obvious evaluation map

$$\boldsymbol{ev}^{\boldsymbol{T}} : \prod_{v \in V(T)} \mathcal{M}^{\boldsymbol{T}}(v) \to M^{\boldsymbol{E}_{\text{int}}} \times M^{\boldsymbol{E}_{\text{int}}}.$$

We define

$$\mathcal{M}^{\boldsymbol{T}}(\boldsymbol{p}, \boldsymbol{E}, \boldsymbol{\ell}, \boldsymbol{a}) := \left(\boldsymbol{ev}^{\boldsymbol{T}}\right)^{-1}\left(\triangle^{\boldsymbol{T}}\right),$$

where

$$\triangle^{\boldsymbol{T}} \subset M^{\boldsymbol{E}_{\text{int}}} \times M^{\boldsymbol{E}_{\text{int}}}$$

denotes the diagonal.

Definition 4.5.13. As a set, we define

$$\overline{\mathcal{M}}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a}) := \coprod_{\boldsymbol{T}} \mathcal{M}^{\boldsymbol{T}}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a}).$$

We equip it with the Gromov topology.

Proposition 4.5.14. *The space $\overline{\mathcal{M}}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$, is compact.*

Proof. Standard Gromov compactness (see [33]) says that any sequence in $\overline{\mathcal{M}}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$ has a subsequence which Gromov-converges to a nodal disk. The space $\overline{\mathcal{M}}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$, by definition, contains all **semi-stable** nodal disks. These are the nodal disks such that each sphere bubble has ≥ 3 marked points, and each disk bubble has

$$\#\text{ boundary punctures} + 2(\#\text{ marked points}) \geq 2.$$

Thus, to prove the result, we must show that, if a non-constant sphere or disk bubbles off in our moduli space, it must be semi-stable. Sphere bubbling was dealt with in Proposition 4.5.4. Any unstable disk has one boundary puncture and no internal marked points; these are constant by exactness of the Lagrangians in $M \setminus \boldsymbol{D}$. Thus there can be no unstable sphere or disk bubbling. $\square$

Lemma 4.5.15. *Suppose that all strata $\mathcal{M}^{\boldsymbol{T}}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$ of $\overline{\mathcal{M}}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$ which contain a sphere bubble have negative virtual dimension. Then, for a generic choice of perturbation data satisfying Condition 4.5.7, $\overline{\mathcal{M}}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$ is a compact topological manifold with corners, and each stratum has the expected dimension.*

Proof. It follows as in Proposition 4.5.9 that any stratum whose virtual dimension is negative is actually empty. Therefore, there are no sphere bubbles. Holomorphic disks intersect the divisors transversely, and are therefore regular. Standard gluing theorems then show that the moduli space is a topological manifold with corners. □

We are particularly interested in moduli spaces of dimension 0 and 1, since we use those to define the Fukaya category. In particular, since sphere bubbles always have codimension ≥ 2, we have the following result:

Corollary 4.5.16. *Suppose we are given a set of Lagrangian boundary conditions $\boldsymbol{L}$ with associated generators $\boldsymbol{p}$, a labelling $\ell : \boldsymbol{F} \rightarrow [k]$, and a tuple $\boldsymbol{a}$. Then, for a generic choice of perturbation data,*

- *If the virtual dimension of $\overline{\mathcal{M}}(\boldsymbol{p}, \ell, \boldsymbol{a})$ is 0, then it is a compact zero-dimensional manifold;*
- *If the virtual dimension of $\overline{\mathcal{M}}(\boldsymbol{p}, \ell, \boldsymbol{a})$ is 1, then it is a compact one-dimensional manifold, with boundary consisting of the 0-dimensional moduli spaces $\mathcal{M}^{\boldsymbol{T}}(\boldsymbol{p}, \ell, \boldsymbol{a})$, where $\boldsymbol{T}$ has two vertices, both contained in T_L.*

Now let $\boldsymbol{L}$ be a set of Lagrangian labels, $\boldsymbol{p}$ an associated set of generators, a be a positive integer, and D_j be one of the divisors. We define the three types of boundary strata in the Gromov compactification of $\mathcal{M}_2(\boldsymbol{p}, j, a)$ (compare Figure 4.2.2). We observe that any spheres bubbling off from a sequence in $\mathcal{M}_2(\boldsymbol{p}, j, a)$ are necessarily constant, because they do not intersect the divisors D_i for $i \neq j$; thus we need only consider strata consisting of disk bubbles.

Definition 4.5.17. Let $\boldsymbol{T}$ consist of the following data:

- A directed planar tree T with Lagrangian labels $\boldsymbol{L}$, and a distinguished vertex v_1;
- For each edge e of T, a generator $p_e \in CF^*(L_{r(e)}, L_{l(e)})$,

such that all vertices are semi-stable with the possible exception of v_1. We define

$$\mathcal{M}_2^{1;\boldsymbol{T}}(\boldsymbol{p}, j, k) := \mathcal{M}_2(\boldsymbol{p}_{v_1}, j, a) \times \prod_{v \neq v_1} \mathcal{M}(\boldsymbol{p}_v, \ell_v)$$

(note that for $v \neq v_1$, $\boldsymbol{F}_v = \phi$, so the ℓ_v is irrelevant but we include it in the notation for consistency).

The second stratum corresponds to $t \rightarrow 0$, so the marked points z_1 and z_2 come together and bubble off a pseudo-holomorphic sphere. This sphere has intersection number 0 with all the divisors other than D_j, hence it must be constant. Thus, the holomorphic disk attached to the sphere has intersection number $(a+1)$ with the divisor D_j, and 0 with the other divisors, and only intersects D_j at the nodal point z where it is attached to the constant sphere. It follow that the disk is tangent to D_j at z to order $a+1$. In other words, it is an element of $\mathcal{M}_1(\boldsymbol{p}, j, a+1)$ (i.e., we can choose our perturbation data to make this so).

Definition 4.5.18. Let $\boldsymbol{T}$ consist of the following data:

- A directed planar tree T with Lagrangian labels $\boldsymbol{L}$, and a distinguished vertex v_1;

- For each edge e of T, a generator $p_e \in CF^*(L_{r(e)}, L_{l(e)})$;

such that all vertices are semi-stable with the possible exception of v_1. We define

$$\mathcal{M}_2^{2,\boldsymbol{T}}(\boldsymbol{p}, j, a) := \mathcal{M}(\boldsymbol{p}_{v_1}, j, a+1) \times \prod_{v \neq v_1} \mathcal{M}(\boldsymbol{p}_v, \ell_v).$$

The third stratum corresponds to $t \to 1$, so the marked points z_1 and z_2 move to the boundary and bubble off disks at the boundary.

Definition 4.5.19. Let $\boldsymbol{T}$ consist of the following data:

- A directed planar tree T with Lagrangian labels $\boldsymbol{L}$, and two distinguished vertices v_1 and v_2;
- For each edge e of T, a generator $p_e \in CF^*(L_{r(e)}, L_{l(e)})$,

such that all vertices are semi-stable with the possible exception of v_1 and v_2, and the branch of T containing v_1 lies strictly to the left of the branch containing v_2. We define

$$\mathcal{M}_2^{3,\boldsymbol{T}}(\boldsymbol{p}, j, a) := \mathcal{M}_1(\boldsymbol{p}_{v_1}, j, a) \times \mathcal{M}_1(\boldsymbol{p}_{v_2}, j, 1) \times \prod_{v \neq v_1, v_2} \mathcal{M}(\boldsymbol{p}_v, \ell_v).$$

Definition 4.5.20. We define the moduli space

$$\overline{\mathcal{M}}_2(\boldsymbol{p}, j, a) := \left(\coprod_{\boldsymbol{T}} \mathcal{M}_2^{1,\boldsymbol{T}}(\boldsymbol{p}, j, a)\right) \coprod \left(\coprod_{\boldsymbol{T}} \mathcal{M}_2^{2,\boldsymbol{T}}(\boldsymbol{p}, j, a)\right) \coprod \left(\coprod_{\boldsymbol{T}} \mathcal{M}_2^{3,\boldsymbol{T}}(\boldsymbol{p}, j, a)\right)$$

as a set.

Lemma 4.5.21. *For generic choice of perturbation data, $\overline{\mathcal{M}}_2(\boldsymbol{p}, j, a)$ has the structure of a compact manifold with corners, of the expected dimension (see Lemma 4.4.4). Furthermore,*

- *The stratum $\mathcal{M}_2^{1,\boldsymbol{T}}(\boldsymbol{p}, j, k)$ has codimension $|V(T)| - 1$,*
- *The stratum $\mathcal{M}_2^{2,\boldsymbol{T}}(\boldsymbol{p}, j, k)$ has codimension $|V(T)|$, and*
- *The stratum $\mathcal{M}_2^{3,\boldsymbol{T}}(\boldsymbol{p}, j, k)$ has codimension $|V(T)| - 2$.*

4.6. **Branched covers.** Let $\phi : (N, \boldsymbol{D}') \to (M, \boldsymbol{D})$ be an $\boldsymbol{a}$-branched cover of Kähler pairs (see Definition 3.5.9).

Let $\boldsymbol{L}$ be a tuple of anchored Lagrangian branes in $N \setminus \boldsymbol{D}'$, $\boldsymbol{p}$ an associated set of generators, $\boldsymbol{E}$ a finite set, and ℓ a labelling. Denote by $\phi(\boldsymbol{L})$ the image of these branes in $M \setminus \boldsymbol{D}$, and by $\phi(\boldsymbol{p})$ the associated set of generators. We would like to related the moduli space $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell)$ of disks in N and the moduli space $\mathcal{M}(\phi(\boldsymbol{p}), \boldsymbol{E}, \ell, \boldsymbol{a})$ of disks in M.

Let us choose perturbation data for the moduli space $\mathcal{M}(\phi(\boldsymbol{p}), \boldsymbol{E}, \ell, \boldsymbol{a})$ in M.

Condition 4.6.1. In a neighbourhood of the divisors $\boldsymbol{D} \subset M$, the almost-complex structure part of the perturbation datum is equal to the standard (integrable) complex structure.

Under Condition 4.6.1, the pullback of the perturbation data in M by ϕ to N is a valid choice of perturbation data in N. Note that this is not true for generic perturbation data: the pullback of a generic almost-complex structure by ϕ may be singular along the divisors $\boldsymbol{D}$.

Lemma 4.6.2. *If our perturbation data in M satisfy Condition 4.6.1, and we use the pulled-back perturbation data to define the moduli space in N, then there is an isomorphism of moduli spaces:*

$$\begin{array}{ccc} \mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell) & \xrightarrow{\cong} & \mathcal{M}(\phi(\boldsymbol{p}), \boldsymbol{E}, \ell, \boldsymbol{a}), \\ u & \mapsto & \phi \circ u. \end{array}$$

Proof. It is clear that this map is well-defined and injective. It is also surjective: suppose we are given $u \in \mathcal{M}(\phi(\boldsymbol{p}), \boldsymbol{E}, \ell, \boldsymbol{a})$. It is clear that, locally, u lifts to a pseudoholomorphic curve in $N \setminus \boldsymbol{D}'$, away from the marked points q_f. At a marked point q_f, u is tangent to the divisor $D_{\ell(f)}$ to order $a_{\ell(f)} - 1$, and it follows that a loop around q_f gets mapped to a loop going $a_{\ell(f)}$ times around divisor $D_{\ell(f)}$. Therefore, a punctured neighbourhood of q_f lifts to $N \setminus \boldsymbol{D}'$. By the removable singularity theorem, the point q_f also lifts, so u lifts locally on a neighbourhood of the marked points q_f. Therefore, since the disk is contractible, u lifts to N, and the lift is clearly an element of $\mathcal{M}(\boldsymbol{p}, \boldsymbol{E}, \ell, \boldsymbol{a})$. □

We observe that it is possible to achieve regularity of the moduli spaces $\mathcal{M}(\phi(\boldsymbol{p}), \boldsymbol{E}, \ell)$ if we require our perturbation data to satisfy Condition 4.6.1, because we can still perturb the Hamiltonian part of the perturbation data essentially arbitrarily away from the strip-like ends and marked points (the transversality argument follows [3, Section 9k]). However, it is **not** possible to achieve regularity of the moduli spaces $\mathcal{M}_0(\boldsymbol{E}, \ell)$: in particular, on a sphere bubble contained inside one of the divisors, the perturbation datum is required to be equal to the standard complex structure, which may not be regular. Therefore we can in general **not** guarantee regularity of the Gromov compactification

$$\overline{\mathcal{M}}(\phi(\boldsymbol{p}), \boldsymbol{E}, \ell),$$

since its strata involve sphere bubbles.

However, we recall that spheres can not bubble off from the moduli spaces $\mathcal{M}_1(\boldsymbol{p}, j, a)$ and $\mathcal{M}_2(\boldsymbol{p}, j, a)$, because a non-constant sphere bubble must intersect all of the divisors. Therefore, we **can** achieve regularity of the Gromov compactifications

$$\overline{\mathcal{M}}_1(\phi(\boldsymbol{p}), j, a) \text{ and } \overline{\mathcal{M}}_2(\phi(\boldsymbol{p}), j, a)$$

with perturbation data satisfying Condition 4.6.1. This allows us to prove:

Lemma 4.6.3. *There exist choices of perturbation data in M and N such that Lemma 4.5.15 remains true in both M and N, and Lemma 4.5.21 remains true in M, and such that there are furthermore isomorphisms of moduli spaces*

$$\begin{array}{ccc} \mathcal{M}_1(\boldsymbol{p}, j, 1) & \overset{\cong}{\to} & \mathcal{M}_1(\phi(\boldsymbol{p}), j, a_j) \\ u & \mapsto & \phi \circ u \end{array}$$

for all $\boldsymbol{p}, j$.

Proof. First, we choose perturbation data on the moduli spaces $\mathcal{M}_1(\phi(\boldsymbol{p}), j, a_j)$ satisfying Condition 4.6.1. We define the perturbation data on $\mathcal{M}_1(\boldsymbol{p}, j, 1)$ to be the pullback of this perturbation data under ϕ. We can then extend these choices to consistent choices of perturbation data for **all** of the moduli spaces $\mathcal{M}_0$, $\mathcal{M}$, $\mathcal{M}_1$, $\mathcal{M}_2$, separately in M and N, such that the moduli spaces and their Gromov compactifications are regular. □

5. THE RELATIVE FUKAYA CATEGORY

In this section, we give our definition of the **relative Fukaya category** of a Kähler pair $(M, \boldsymbol{D})$, which we denote $\mathcal{F}(M, \boldsymbol{D})$. It is a (possibly curved) $\boldsymbol{G}(M, \boldsymbol{D})$-graded deformation of the affine Fukaya category, in the sense of Definition 2.4.3. We also define an orbifold version of the relative Fukaya category. In Section 5.2, we describe the behaviour of the relative Fukaya category with respect to branched covers.

5.1. The definition.

Suppose that $(M, \boldsymbol{D})$ is a Kähler pair, and $\boldsymbol{a} = (a_1, \ldots, a_k)$ a tuple of k positive integers (where k is the number of divisors in $\boldsymbol{D}$).

Definition 5.1.1. We define the ring

$$R_{\boldsymbol{a}} := \mathbb{C}[[r_1, \ldots, r_k]],$$

and equip it with the $\boldsymbol{G}(M, \boldsymbol{D})$-grading, where r_j has grading

$$(2(1 - a_j), a_j y_j) \in (\mathbb{Z} \oplus Y)/Z$$

(see Corollary 3.5.5).

Example 5.1.2. Suppose that $(M, \boldsymbol{D}) = (M_1^n, \boldsymbol{D})$ as in Example 3.5.2, and $\boldsymbol{a} = (a, \ldots, a)$. Then we have

$$R_{\boldsymbol{a}} \cong R_a^n,$$

where R_a^n is the $\boldsymbol{G}(M, \boldsymbol{D})$-graded ring introduced in Definition 2.2.13.

We give a definition of the **smooth orbifold relative Fukaya category** $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$, based on the definition of the relative Fukaya category given in [1]. It is a (possibly curved) $\boldsymbol{G}(M, \boldsymbol{D})$-graded deformation of the affine Fukaya category over $R_{\boldsymbol{a}}$. We denote

$$\mathcal{F}(M, \boldsymbol{D}) := \mathcal{F}(M, \boldsymbol{D}, \boldsymbol{1}),$$

and call it the **relative Fukaya category**.

The objects of $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$ are the same as in the affine Fukaya category: anchored Lagrangian branes in $M \setminus \boldsymbol{D}$. Given a tuple of anchored Lagrangian branes $\boldsymbol{L}$ with associated generators $\boldsymbol{p}$, and an element $\boldsymbol{d} \in \mathbb{Z}_{\geq 0}^k$, we choose a labelling ℓ such that $\boldsymbol{d}(\ell) = \boldsymbol{d}$, and define the coefficient of $r^{\boldsymbol{d}} p_0$ in $\mu^s(p_s, \ldots, p_1)$ to be

$$\frac{\#\left(\mathcal{M}(\boldsymbol{p}, \ell, \boldsymbol{a})\right)}{\boldsymbol{d}!},$$

where we denote

$$\boldsymbol{d}! := d_1! d_2! \ldots d_k!,$$

and $\#$ denotes a signed count of the zero-dimensional part of the moduli space, with signs defined according to the canonical isomorphism of orientation spaces given by Lemma 4.4.2.

We observe that this is a finite sum, by Corollary 4.5.16. Note that, while the affine Fukaya category is not curved (any pseudoholomorphic disk with boundary on an exact Lagrangian is constant, since it has zero energy), the relative Fukaya category may be curved.

It follows from the index computation in Lemma 4.4.2 that the structure maps μ^s define a $\boldsymbol{G}(M,\boldsymbol{D})$-graded A_∞ deformation of the affine Fukaya category over $R_{\boldsymbol{a}}$. Observe that the order-0 component of μ^s counts disks that completely avoid the divisors $\boldsymbol{D}$, and therefore coincides with the definition of the structure maps in the affine Fukaya category.

The fact that $\mu\circ\mu=0$ follows also from Corollary 4.5.16, since the signed count of boundary points of a compact 1-dimensional manifold with boundary is 0. The sign computation follows directly from that of [3, Section 12g], essentially because the fibres of the forgetful maps $\mathcal{R}(\boldsymbol{L},\boldsymbol{E})\to\mathcal{R}(\boldsymbol{L},\phi)$ have a complex structure, hence are canonically oriented.

Observe that we needed the factor $(\boldsymbol{d}!)^{-1}$ in the definition of the structure coefficients μ^s. This is because, if $\boldsymbol{d}=\boldsymbol{d}_1+\boldsymbol{d}_2$, then given a labelling $\ell:\boldsymbol{F}\to[k]$ with $\boldsymbol{d}(\ell)=\boldsymbol{d}$, there are $\boldsymbol{d}!/(\boldsymbol{d}_1!\boldsymbol{d}_2!)$ ways of choosing a partition $\boldsymbol{F}=\boldsymbol{F}_1\sqcup\boldsymbol{F}_2$ such that the restricted labellings ℓ_1, ℓ_2 on $\boldsymbol{F}_1$ and $\boldsymbol{F}_2$ satisfy $\boldsymbol{d}(\ell_1)=\boldsymbol{d}_1$ and $\boldsymbol{d}(\ell_2)=\boldsymbol{d}_2$. So, in the boundary of the one-dimensional component of the moduli space $\mathcal{M}(\boldsymbol{p},\ell)$, which consists of nodal disks with two components, there are $\boldsymbol{d}!/(\boldsymbol{d}_1!\boldsymbol{d}_2!)$ ways for the marked points q_f to be distributed between the two components.

It is important to consider in what sense the smooth orbifold relative Fukaya category is dependent on the choices (of Floer and perturbation data) involved in its construction. We recall the argument of [3, Section 10a]: Let I denote a set of possible choices of Floer and perturbation data. For each $i\in I$, we denote by $\mathcal{F}(M,\boldsymbol{D},\boldsymbol{a})^i$ the smooth orbifold relative Fukaya category defined using those choices. We define a new A_∞ category, the **total** category $\mathcal{F}(M,\boldsymbol{D},\boldsymbol{a})^{\mathrm{tot}}$, as follows:

- Objects are pairs (L,i) where L is an object of $\mathcal{F}(M,\boldsymbol{D},\boldsymbol{a})$ and $i\in I$;
- For each pair of objects we choose a Floer datum, and for each set of labels of objects we choose a perturbation datum;
- We require that, for a pair $(L_0,i),(L_1,i)$, the Floer datum is that given by i;
- We require that, for a set of labels $(L_0,i),\dots,(L_k,i)$, the perturbation data are those given by the index i;
- The rest of the Floer and perturbation data we choose arbitrarily.

The rest of the construction (of morphism spaces and composition maps) follows that of the smooth orbifold relative Fukaya category. It follows that for each $i\in I$, there is a full embedding

$$\mathcal{F}(M,\boldsymbol{D},\boldsymbol{a})^i\hookrightarrow\mathcal{F}(M,\boldsymbol{D},\boldsymbol{a})^{\mathrm{tot}}$$

which is given, on the level of objects, by

$$L\mapsto(L,i).$$

When restricting to the affine Fukaya category ($r_j=0$), it follows from the PSS isomorphism that these embeddings are quasi-equivalences, and hence invertible, and therefore that the affine Fukaya category does not depend on the choice of data $i\in I$, up to quasi-equivalence (see [3, Section 10a].

This need no longer be the case for the relative Fukaya category (these embeddings need not be quasi-equivalences nor invertible), and in general the relative Fukaya may depend on the data used to define it (compare [2, Section 8f]). However, in this paper we are only interested in a certain full subcategory of the relative Fukaya category which is necessarily **minimal** (for grading reasons). Therefore, rather than quasi-equivalence (which we recall is not necessarily a well-behaved notion over the power series ring R), we can use the simpler notion of formal diffeomorphism.

Let $\mathcal{L}$ be a set of objects of $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$, and I be some set of possible choices of Floer and perturbation data for the full subcategory $\mathcal{C} \subset \mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$ with objects $\mathcal{L}$. Let us form the total category $\mathcal{C}^{\mathrm{tot}}$, as above.

Lemma 5.1.3. *Suppose that we can choose Floer data and perturbation data for $\mathcal{C}^{\mathrm{tot}}$ so that it is minimal. Then, for any $i, j \in I$, there is a $\boldsymbol{G}$-graded quasi-equivalence of minimal A_∞ categories over R,*

$$\mathcal{C}^i \cong \mathcal{C}^j.$$

Proof. First, note that we have A_∞ embeddings of minimal A_∞ categories

$$\mathcal{C}^i \hookrightarrow \mathcal{C}^{\mathrm{tot}} \hookleftarrow \mathcal{C}^j,$$

as above. Now note that, because the objects (L, i) and (L, j) are quasi-isomorphic when restricted to the affine Fukaya category, and $\mathcal{C}^{\mathrm{tot}}$ is minimal, these objects are quasi-isomorphic in $\mathcal{C}^{\mathrm{tot}}$. It follows that the above embeddings are quasi-equivalences. By Lemma 2.3.34, quasi-equivalences of minimal A_∞ categories can be inverted over R. It follows that there is a quasi-equivalence $\mathcal{C}^i \cong \mathcal{C}^j$, as required. □

In other words, if we can choose the category $\mathcal{C}^{\mathrm{tot}}$ to be minimal (e.g., for grading reasons), then $\mathcal{C}$ is independent of the choice of perturbation data $i \in I$ made in its construction, up to formal diffeomorphism.

Remark 5.1.4. If M is Calabi-Yau, then by Remark 3.5.7, there is a canonical morphism of grading data

$$\boldsymbol{p} : \boldsymbol{G}(M, \boldsymbol{D}) \to \boldsymbol{G}_{\mathbb{Z}}.$$

Thus, we obtain a canonical $\mathbb{Z}$-grading on the category

$$\mathcal{F}(M, \boldsymbol{D}) \cong \boldsymbol{p}_* \mathcal{F}(M, \boldsymbol{D}),$$

such that the $\mathbb{Z}$-grading of R is zero.

5.2. **Behaviour with respect to ramified covers.** Suppose that $\phi : (N, \boldsymbol{D}') \to (M, \boldsymbol{D})$ is an $\boldsymbol{a}$-branched cover of Kähler pairs, where $\boldsymbol{a} = (a_1, \ldots, a_k)$. In this section, we will examine the relationship between the following three categories:

$$\mathcal{F}(M, \boldsymbol{D}),\ \mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a}), \text{ and } \mathcal{F}(N, \boldsymbol{D}').$$

We first recall the behaviour of the affine Fukaya category under covers. Note that $\phi : N \setminus \boldsymbol{D}' \to M \setminus \boldsymbol{D}$ is a finite (unramified) cover. Thus, it induces an injective morphism of grading data,

$$\boldsymbol{p} : \boldsymbol{G}(N, \boldsymbol{D}') \to \boldsymbol{G}(M, \boldsymbol{D}).$$

We recall from Proposition 3.4.1 that there is a fully faithful embedding

$$\boldsymbol{p}^*\mathcal{F}(M \setminus \boldsymbol{D}) \to \mathcal{F}(N \setminus \boldsymbol{D'}).$$

We observe that, if it were possible to choose perturbation data in M such that Condition 4.6.1 were satisfied, then Lemma 4.6.2 would imply, by a similar argument, that there is a fully faithful embedding

$$\boldsymbol{p}^*\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a}) \to \mathcal{F}(N, \boldsymbol{D'}).$$

Remark 5.2.1. We observe that $\boldsymbol{p}^* R_{\boldsymbol{a}}$ is exactly the $\boldsymbol{G}(N, \boldsymbol{D'})$-graded coefficient ring over which $\mathcal{F}(N, \boldsymbol{D'})$ is defined; this follows immediately from the definition of $R_{\boldsymbol{a}}$ (Definition 5.1.1) and Lemma 3.5.11.

However, we recall (see discussion in Section 4.6) that it is **not** possible to guarantee sufficient regularity of all of our moduli spaces under Condition 4.6.1, so we can not quite make this statement. Ideally, we would find a better version of Condition 4.6.1 that would allow us to guarantee lifting and regularity. However, we have been unable to do this, and instead circumvent this problem by a rather ugly and ad-hoc method, which we now describe.

We recall, from the discussion at the end of Section 4.6, that it **is** possible to obtain regularity for moduli spaces with only a single marked point, under Condition 4.6.1. Therefore, we can make sure that the result is true 'to first order' (see Section 2.6). Recall (Section 2.6) that we denote by $\mathfrak{m} \subset R$ the maximal ideal. If $\mathcal{F}$ is an R-linear A_∞ category, then $\mathcal{F}/\mathfrak{m}^2$ is an $R/\mathfrak{m}^2$-linear category, which retains only the information about the first-order part of $\mathcal{F}$. Now we prove the result:

Proposition 5.2.2. *Given an $\boldsymbol{a}$-branched cover of Kähler pairs $\phi : (N, \boldsymbol{D'}) \to (M, \boldsymbol{D})$, there exists a $\boldsymbol{G}(M, \boldsymbol{D})$-graded A_∞ category $\mathcal{F}(\phi)$ over $R_{\boldsymbol{a}}$, such that there exist A_∞ functors*

$$\mathcal{G}_1 : \boldsymbol{p}^*\mathcal{F}(\phi) \to \mathcal{F}(N, \boldsymbol{D'}),$$

and

$$\mathcal{G}_2 : \mathcal{F}(\phi)/\mathfrak{m}^2 \to \mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})/\mathfrak{m}^2,$$

and both $\mathcal{G}_1/\mathfrak{m}$ and $\mathcal{G}_2/\mathfrak{m}$ are quasi-equivalences of the zeroth-order categories.

Proof. We would like to choose perturbation data in N which are invariant with respect to the action of $Y(M, \boldsymbol{D})$, but this will be problematic, because the geometric action of $Y(M, \boldsymbol{D})$ on N is non-trivial, so we need to employ a trick to bypass this problem (compare [2, Section 8b]).

The covering map ϕ induces an injective homomorphism

$$p : Y(N, \boldsymbol{D'}) \to Y(M, \boldsymbol{D}).$$

We denote

$$\Gamma := Y(M, \boldsymbol{D})/Y(N, \boldsymbol{D'}).$$

As part of the construction of $\mathcal{F}(\phi)$, we choose a function

$$q : \Gamma \to Y(M, \boldsymbol{D})$$

(**not** necessarily a group homomorphism) that splits the map $Y(M, \boldsymbol{D}) \to \Gamma$.

Objects of $\mathcal{F}(\phi)$ are pairs (L, γ), where L is an anchored Lagrangian brane in $M \setminus \boldsymbol{D}$ and $\gamma \in \Gamma$. We define $Y(M, \boldsymbol{D})$ to act on our objects by

$$y \cdot (L, \gamma) := ((y - q(y)) \cdot L, \gamma + y),$$

and Γ to act by

$$\gamma \cdot (L, \gamma') := (L, \gamma + \gamma').$$

On the level of objects, the A_∞ morphisms are given by

$$(L, \gamma) \mapsto q(\gamma) \cdot L$$

(recall that there is a correspondence between anchored Lagrangian branes in $M \setminus \boldsymbol{D}$ and anchored Lagrangian branes in $N \setminus \boldsymbol{D'}$).

We now choose Floer data for our anchored Lagrangian branes in M, satisfying Condition 4.6.1, invariant under the action of $Y(M, \boldsymbol{D}) \oplus \Gamma$. This is possible because the action of this group on the underlying geometric Lagrangians is trivial. We define the morphism spaces as usual, with grading defined so that

$$CF^*_{\mathcal{F}(\phi)}((L_0, \gamma_0), (L_1, \gamma_1)) \cong CF^*_{\mathcal{F}(M \setminus \boldsymbol{D})}(q(\gamma_0) \cdot L_0, q(\gamma_1) \cdot L_1)$$

as $\boldsymbol{G}$-graded vector spaces.

This makes $\mathcal{F}(\phi)$ into a $\boldsymbol{G}(M, \boldsymbol{D})$-graded pre-category. For labellings ℓ with $|\boldsymbol{d}(\ell)| \le 1$, we choose perturbation data in M for the moduli spaces $\mathcal{M}(\boldsymbol{p}, \ell, \boldsymbol{a})$, also satisfying Condition 4.6.1 and invariant under the action of $Y(M, \boldsymbol{D}) \oplus \Gamma$. This allows us to define $\mathcal{F}(\phi)$ to first order, by counting holomorphic disks in M intersecting only a single divisor, exactly by analogy with $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})/\mathfrak{m}^2$. Thus, we in fact have an isomorphism of $\boldsymbol{G}(M, \boldsymbol{D})$-graded categories

$$\mathcal{F}(\phi)/\mathfrak{m}^2 \cong \mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})/\mathfrak{m}^2 \otimes \mathbb{C}[\Gamma],$$

where $\gamma \in \Gamma$ has degree $q(\gamma)$. In particular, the categories are quasi-equivalent.

Furthermore, we observe that the above-defined map on the level of objects,

$$\begin{array}{rcl} \boldsymbol{p}^*\mathcal{F}(\phi) & \to & \mathcal{F}(N, \boldsymbol{D'}) \\ (L, \gamma) & \mapsto & q(\gamma) \cdot L, \end{array}$$

defines an equivalence of $\boldsymbol{G}(N, \boldsymbol{D'})$-graded pre-categories. I.e., a generator p of $CF^*((L_0, \gamma_0), (L_1, \gamma_1))$ only lifts to a generator of $CF^*(q(\gamma_0) \cdot L_0, q(\gamma_1) \cdot L_1)$ if its degree $y + q(\gamma_0 - \gamma_1) \in Y(M, \boldsymbol{D})$ lies in the image of $Y(N, \boldsymbol{D'})$.

Now we extend the definition of $\mathcal{F}(\phi)$ to higher-order terms. Given objects $\boldsymbol{L} := ((L_0, \gamma_0), \ldots, (L_s, \gamma_s))$, with associated generators $\boldsymbol{p} = (p_0, \ldots, p_s)$ of degrees $(y_0, \ldots, y_s)$ in $Y(M, \boldsymbol{D})$, and ℓ a labelling, such that

$$y_0 = f(2 - s - 2|\boldsymbol{d}(\ell) - \boldsymbol{d}_{\boldsymbol{a}}(\ell)|) + p_Y(\boldsymbol{d}_{\boldsymbol{a}}(\ell)) + \sum_{j=1}^{s} y_j$$

(recall that this condition must be satisfied if the coefficient of $r^{\boldsymbol{d}(\ell)} p_0$ in $\mu^s(p_s, \ldots, p_1)$ is to be non-zero), we define $\phi^*(\boldsymbol{L}, \boldsymbol{p})$ to be the tuple of anchored Lagrangian branes in $N \setminus \boldsymbol{D'}$,

$$(q(\gamma_0) \cdot L_0, (y_1 + q(\gamma_1)) \cdot L_1, \ldots, (y_1 + \ldots + y_s + q(\gamma_s)) \cdot L_s),$$

with the associated generators which are the lifts of the p_j (note that p_0 does lift, by the equation we imposed on the y_j). We define perturbation data for the moduli spaces $\mathcal{M}(\phi^*(\boldsymbol{L}, \boldsymbol{p}), \ell)$ in N, such that:

- They are given by the pullback under ϕ of the perturbation data on $\mathcal{M}(\boldsymbol{p}, \ell, \boldsymbol{a})$ in M, for $|\boldsymbol{d}(\ell)| \le 1$ (recalling that these perturbation data satisfy Condition 4.6.1, and hence can be pulled back under ϕ);
- They are invariant with respect to the action of $Y(M, \boldsymbol{D})$ on objects.

Note that, we can still achieve transversality in the presence of this final assumption. To see why, observe that the action of $Y(M, \boldsymbol{D})$ may involve a non-trivial geometric action of the covering group Γ. However, under the action

$$y \cdot (L, \gamma) = ((y - q(y)) \cdot L, \gamma + y),$$

the element in the first factor $y - q(y)$ acts trivially on N. Thus, we can choose perturbation data by considering only the first factors of our objects, on which the action of $Y(M, \boldsymbol{D})$ is trivial, then push these perturbation data forward using the geometric action of the second factors via q.

We now explain why the Gromov compactifications $\overline{\mathcal{M}}(\phi^*(\boldsymbol{L}, \boldsymbol{p}), \ell)$ of zero- and one-dimensional moduli spaces are generically regular. We recall the discussion of Section 4.6: the only obstruction to regularity is the appearance of sphere bubbles contained entirely within one of the divisors $\boldsymbol{D}$. If we imposed Condition 4.6.1 on all of our moduli spaces, we could not guarantee this, because the almost-complex structure on each divisor would be required to be the standard (integrable) complex structure, and hence not necessarily regular. However, we have only imposed this condition on moduli spaces $\mathcal{M}(\phi^*(\boldsymbol{L}, \boldsymbol{p}), \ell)$ where $|\boldsymbol{d}(\ell)| \le 1$, and any sphere bubbling off from such a moduli space is necessarily constant, because it does not intersect some divisor D_i. Therefore, these moduli spaces are generically regular. For the remaining moduli spaces with $|\boldsymbol{d}(\ell)| \ge 2$, we can perturb the almost-complex structure arbitrarily on the divisors, and therefore we can apply the argument of Proposition 4.5.9 to prove regularity. Thus, our moduli spaces satisfy the analogue of Corollary 4.5.16.

We now define the coefficient of $r^{\boldsymbol{d}} \mu^s(p_s, \ldots, p_1)$ to be the signed count of points in the zero-dimensional part of the moduli space $\mathcal{M}(\phi^*(\boldsymbol{L}, \boldsymbol{p}), \ell)$, where $\boldsymbol{d}(\ell) = \boldsymbol{d}$. It follows as in the definition of the relative Fukaya category that μ^s define a $\boldsymbol{G}(M, \boldsymbol{D})$-graded A_∞ deformation of $\mathcal{F}(M \setminus \boldsymbol{D})$ over $R_{\boldsymbol{a}}$. It follows immediately from the definition that there is an A_∞ quasi-equivalence

$$\boldsymbol{p}^* \mathcal{F}(\phi) \to \mathcal{F}(N, \boldsymbol{D}'),$$

sending $(L, \gamma) \mapsto \gamma \cdot L$ on the level of objects. This completes the proof. □

Remark 5.2.3. We recall that the notion of quasi-equivalence of A_∞ categories over R is not necessarily well-behaved; however, in the situation in which we will apply Proposition 5.2.2, the parts of the categories $\mathcal{F}(\phi)$, $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$ and $\mathcal{F}(N, \boldsymbol{D}')$ will necessarily be minimal, so quasi-equivalences are well-behaved by Lemma 2.3.34.

Now we would like to relate $\mathcal{F}(M, \boldsymbol{D})$ to $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$. Again, we will only relate the first-order part of the deformations. We recall from Section 2.6 that a first-order deformation of the A_∞ category $\mathcal{F} := \mathcal{F}(M \setminus \boldsymbol{D})$ over R consists of A_∞ structure maps

$$\mu^* = \mu^*_0 + \mu^*_1,$$

where μ^*_0 gives the structure maps of $\mathcal{F}$, and the A_∞ relations say that μ^*_1 defines a class

$$[\mu_1] \in HH^2(\mathcal{F}, \mathcal{F} \otimes R^1_{\boldsymbol{1}}).$$

Theorem 4. *Let* $\mathcal{F} := \mathcal{F}(M \setminus \boldsymbol{D})$, $\boldsymbol{G} := \boldsymbol{G}(M, \boldsymbol{D})$, *and let*

$$[\mu_{1,\mathbf{1}}] := \sum_{j=1}^{k} r_j \alpha_j \in HH^2_{\boldsymbol{G}}(\mathcal{F}, \mathcal{F} \otimes R^1_{\mathbf{1}})$$

be the first-order deformation class of $\mathcal{F}(M, \boldsymbol{D}, \mathbf{1}) \cong \mathcal{F}(M, \boldsymbol{D})$. *Then the first-order deformation class of* $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$ *is given by*

$$[\mu_{1,\boldsymbol{a}}] = \sum_{j=1}^{k} r_j \alpha_j^{a_j} \in HH^2_{\boldsymbol{G}}(\mathcal{F}, \mathcal{F} \otimes R^1_{\boldsymbol{a}}),$$

where the power is taken with respect to the Yoneda product on $HH^*(\mathcal{F})$.

Proof. We define elements

$$\beta_j(b) \in CC^*(\mathcal{F}),$$

for $b \geq 1$, as follows: Let $\boldsymbol{L}$ be a tuple of anchored Lagrangian branes in $M \setminus \boldsymbol{D}$, with associated generators $\boldsymbol{p}$. Then the coefficient of p_0 in $\beta_j(b)^s(p_s, \ldots, p_1)$ is given by the count of rigid elements in the moduli space $\mathcal{M}_1(\boldsymbol{p}, j, b)$. It follows from the fact that the signed count of points in the boundary of the one-dimensional component of the moduli space $\overline{\mathcal{M}}_1(\boldsymbol{p}, j, b)$ is 0, that each $\beta_j(b)$ is a Hochschild cocycle. Furthermore, by definition we have

$$[\beta_j(1)] = \alpha_j$$

and

$$\mu_{1,\boldsymbol{a}} = \sum_{j=1}^{k} r_j \beta_j(a_j).$$

We also define elements

$$H_j(b) \in CC^*(\mathcal{F})$$

by counting rigid elements in $\mathcal{M}_2(\boldsymbol{p}, j, b)$.

Lemma 5.2.4. *We have*

$$\beta_j(b+1) = \beta_j(b) \bullet \beta_j(1) \pm \partial(H_j(b))$$

in $CC^*(\mathcal{F})$, *where* $\bullet$ *denotes the Yoneda product and* ∂ *denotes the Hochschild differential.*

Proof. The result follows from the fact that the signed count of points in the boundary of the one-dimensional component of the moduli space $\overline{\mathcal{M}}_2(\boldsymbol{p}, j, b)$ is 0. See Lemma 4.5.21 for the description of the boundary components. The boundary points $\mathcal{M}^{1,\boldsymbol{T}}(\boldsymbol{p}, j, b)$ contribute the term $\partial H_j(b)$ to the sum, the boundary points $\mathcal{M}^{2,\boldsymbol{T}}(\boldsymbol{p}, j, b)$ contribute the term $\beta_j(b+1)$, and the boundary points $\mathcal{M}^{3,\boldsymbol{T}}(\boldsymbol{p}, j, b)$ contribute the term $\beta_j(b) \bullet \beta_j(1)$. □

It follows that, on the level of Hochschild cohomology,

$$[\beta_j(b+1)] = [\beta_j(b)] \bullet \alpha_j,$$

and hence, by induction, that

$$[\beta_j(b)] = \alpha_j^b.$$

The result follows immediately. □

6. MORSE-BOTT COMPUTATIONS IN THE FUKAYA CATEGORY

In this section, we consider the Kähler pair $(M, \boldsymbol{D}) = (M_1^n, \boldsymbol{D})$ of Example 3.5.2, and the tuple

$$\boldsymbol{a} := (n, \ldots, n)$$

(n copies) associated to the branched cover of Kähler pairs

$$\phi : (M_n^n, \boldsymbol{D}) \to (M_1^n, \boldsymbol{D})$$

(see Example 3.5.10). We consider the grading datum $\boldsymbol{G} := \boldsymbol{G}(M, \boldsymbol{D})$. We define the $\boldsymbol{G}$-graded rings

$$R := R_1^n \cong \mathbb{C}[[r_1, \ldots, r_n]]$$

and

$$R_n := R_n^n \cong \mathbb{C}[[r_1, \ldots, r_n]]$$

from Definition 2.2.13, and recall that they are the coefficient rings of $\mathcal{F}(M, \boldsymbol{D})$ and $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$ respectively (see Example 5.1.2). Throughout this section, we denote by

$$Y := \mathbb{Z}\langle y_1, \ldots, y_n \rangle$$

the abelian group appearing in the pseudo-grading datum $\boldsymbol{H}(M, \boldsymbol{D})$.

We recall that

$$M := \left\{ \sum_j z_j = 0 \right\} \subset \mathbb{CP}^{n-1},$$

with the divisors $D_j := \{z_j = 0\}$. Thus $M \cong \mathbb{CP}^{n-2}$, and $\boldsymbol{D}$ consists of n hyperplanes with normal crossings. $M \setminus \boldsymbol{D}$ is called the (generalized) **pair of pants**.

We construct an immersed Lagrangian sphere $L^n : S^{n-2} \to M \setminus \boldsymbol{D}$ in the pair of pants (summarising the construction in [16]). The main result of this section (Corollary 6.5.6) is that the endomorphism algebra $CF^*(L^n, L^n)$, computed in $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$, is of type A (see Definition 2.5.2).

To do this, we first give a Morse-Bott description of the endomorphism algebra $CF^*(L^n, L^n)$ in the relative Fukaya category, $\mathcal{F}(M, \boldsymbol{D})$, to first order. Structure coefficients in this description are given by counts of 'holomorphic flipping pearly trees' rather than holomorphic disks. A holomorphic flipping pearly tree is a Morse-Bott version of a holomorphic disk, made out of holomorphic disks and Morse flowlines. We introduce them because it is often possible to explicitly identify moduli spaces of flipping pearly trees, and therefore to make explicit computations of the structure coefficients in the Fukaya category.

The construction is based on [16] (note that the original idea comes from [26]). The extra content here is that, whereas [16] describes the endomorphism algebra in the affine Fukaya category only, we will describe the endomorphism algebra in the first-order relative Fukaya category.

There are some transversality issues in the definition of moduli spaces of flipping pearly trees, involving the possibility of unstable disk and sphere bubbles. In the moduli spaces of holomorphic disks that we used to define the relative Fukaya category, we avoided this problem by introducing extra internal marked points where they intersected the divisors $\boldsymbol{D}$.

This approach is no longer possible for flipping pearly trees: they can intersect the divisors on their boundary. For this reason, we can not guarantee transversality and therefore can not give a complete Morse-Bott description of $CF^*(L^n, L^n)$ in the relative Fukaya category.

However, for moduli spaces with very few intersections with the divisors (in particular, those which intersect only a single divisor), we can rule out any unstable disk or sphere bubbles in an ad hoc way. Thus, we are able to give a Morse-Bott description of $CF^*(L^n, L^n)$ to first order, and in particular to identify the first-order deformation class in $\mathcal{F}(M, \boldsymbol{D})$. This allows us, via Theorem 4, to determine the first-order deformation class in $\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})$. The first-order deformation class is all we need to determine that the algebra is of type A, so the failure of transversality in our Morse-Bott model at higher order does not concern us (to clarify: $CF^*(L^n, L^n)$ is perfectly well-defined at all orders, but our Morse-Bott model for it is well-defined only to first order).

After describing the construction of the Lagrangian L^n in Section 6.1, the structure of this section follows that of Section 4: first we introduce the moduli space of pearly trees (possible domains for a flipping pearly tree), then we describe our choice of perturbation data, then we describe the moduli space of flipping pearly trees (pseudoholomorphic maps into M), explain why transversality holds, then describe Gromov compactness.

6.1. **The Lagrangian immersion** $L^n : S^{n-2} \to M \setminus \boldsymbol{D}$.

In [16], we introduced a one-parameter family of Lagrangian immersions

$$L^n_\epsilon : S^{n-2} \to M \setminus \boldsymbol{D},$$

for $\epsilon > 0$ sufficiently small (actually we called these L^{n-2}; apologies for the change in notation, but it makes many formulae cleaner). We briefly recall the construction of L^n.

We consider the Lagrangian immersion $L' : S^{n-2} \to M$ which is the double cover of the real locus $\mathbb{RP}^{n-2}$ of M. If we think of

$$S^{n-2} := \left\{ \sum_{j=1}^n x_j = 0, \sum_{j=1}^n x_j^2 = 1 \right\} \subset \mathbb{R}^n,$$

then the immersion is given by

$$(x_1, \ldots, x_n) \mapsto [x_1 : \ldots : x_n].$$

We construct the immersion L^n by perturbing the immersion L'.

Namely, by the Weinstein Lagrangian neighbourhood theorem, L' can be extended to an immersion of the radius-η cotangent disk bundle

$$D^*_\epsilon S^{n-2} \to M,$$

which is J_0-holomorphic along the zero section, and such that complex conjugation acts by -1 on the covector.

We construct a function $f : S^{n-2} \to \mathbb{R}$ by setting

$$f(x_1, \ldots, x_n) = \sum_{j=1}^n g(x_j),$$

where $g : \mathbb{R} \to \mathbb{R}$ has the properties

(1) $g'(x) > 0$;
(2) $g(-x) = -g(x)$;
(3) $g(x) = x$ for $|x| < \delta$;
(4) $g'(x)$ is a strictly decreasing function of $|x|$ for $|x| > \delta$;
(5) $g'(x) < \delta$ for $|x| > 2\delta$,

for some small $\delta > 0$.

We then define $L^n_\epsilon : S^{n-2} \to M$ to be the image of the graph of the exact one-form ϵdf in $D^*_\eta S^{n-2}$, under the immersion into M, so that $L' = L^n_0$. The fact that ∇f is transverse to the hypersurfaces $\{x_j = 0\}$ implies that the image L^n_ϵ avoids the divisors $\{z_j = 0\}$ for $\epsilon > 0$, so we obtain a Lagrangian immersion $L^n_\epsilon : S^{n-2} \to M \setminus \boldsymbol{D}$. It has self-intersections at the critical points of f (where it intersects the other branch of the double cover). We observe that, for $n \geq 4$, L^n automatically lifts to $\widetilde{\mathcal{G}}(M \setminus \boldsymbol{D})$, because $\pi_1(S^{n-2}) = 0$. We choose such a lift, and hence define an anchored brane structure on the Lagrangian L^n.

The flowlines of ∇f are illustrated in Figure 6.1.1, in the case $n = 4$. The hypersurfaces $\{x_j = 0\}$ split S^{n-2} into $2^n - 2$ regions, indexed by the proper non-empty sets $K \subset [n]$. Namely, K corresponds to the region where coordinates x_j are negative for $j \in K$ and positive for $x_j \notin K$. Each region contains a unique critical point p_K of f.

The Floer endomorphism algebra $CF^*(L^n, L^n)$ can be defined, despite L^n being immersed (see [16, Section 3.1]), and is generated by the self-intersection points of L^n (which are the points p_K indexed by proper non-empty sets $K \subset [n]$), together with the Morse cohomology of S^n (which we choose to have generators p_ϕ and $p_{[n]}$, corresponding to the identity and top class respectively). Thus, $CF^*(L^n, L^n)$ has generators p_K indexed by subsets $K \subset [n]$.

It follows from [16, Proposition 3.3] and [16, Proposition 3.7] that there is an isomorphism

$$CF^*(L^n, L^n) \cong A$$

as $\boldsymbol{G}$-graded vector spaces, where $A \cong A_n$ is the $\boldsymbol{G}$-graded exterior algebra of Definition 2.2.12.

We now observe that we can define the endomorphism algebra $CF^*(L^n, L^n)$ in $\mathcal{F}(M, \boldsymbol{D})/\mathfrak{m}^2$. Most of the rest of this section is concerned with computing $CF^*(L^n, L^n)$ in $\mathcal{F}(M, \boldsymbol{D})/\mathfrak{m}^2$, using ‘flipping pearly trees’.

Remark 6.1.1. Note that, in [16, Section 3.1], well-definedness of $CF^*(L^n, L^n)$ in the affine Fukaya category was proved by passing to the cover $M^n \setminus \boldsymbol{D}$ of $M \setminus \boldsymbol{D}$, to bypass proving Gromov compactness for immersed Lagrangians. One may worry that this will no longer be valid when we consider disks passing through the divisors $\boldsymbol{D}$, about which the cover $M^n \to M$ has some branching. However, as we saw in the proof of Proposition 5.2.2, for the first-order relative Fukaya category it is possible to choose perturbation data that lift to the branched cover; and therefore we can apply the same trick to rigorously define $CF^*(L^n, L^n)$ in $\mathcal{F}(M, \boldsymbol{D})/\mathfrak{m}^2$.

The main result we will prove is:

Proposition 6.1.2. *There exists a $\boldsymbol{G}$-graded A_∞ category $\mathcal{F}'$ over $R/\mathfrak{m}^2$, such that;*



FIGURE 6.1.1. The case $n = 4$. The dashed circles represent the hypersurfaces $D_j^{\mathbb{R}} = D_j \cap \mathbb{RP}^2$ as labeled. Each region is labeled with the list of coordinates that are negative in that region (e.g., the label '124' means that $x_1 < 0, x_2 < 0, x_3 > 0, x_4 < 0$ in that region). The arrows represent the index-1 Morse flow lines of ∇f. The dots represent critical points of f. The picture really lives on a sphere, and the three points labeled '4' should be identified (at infinity).

- *$\mathcal{F}'$ has two objects: L and L';*
- *L and L' are quasi-isomorphic in the zeroth-order category $\mathcal{F}'/\mathfrak{m}$;*
- *The endomorphism algebra of L in $\mathcal{F}'$ coincides with the endomorphism algebra of L^n in $\mathcal{F}(M, \boldsymbol{D})/\mathfrak{m}^2$;*

Furthermore, the endomorphism algebra of L' in $\mathcal{F}'$ satisfies:

$$CF^*_{\mathcal{F}'}(L', L') \cong (A, \mu^*),$$

where:

- *A is the $\boldsymbol{G}$-graded vector space of Definition 2.2.12;*

- (A, μ_0^2) *coincides with the exterior algebra multiplication on* A*;*
- *We have*

$$\Phi(\mu^*) = u_1 \dots u_n + \sum_{j=1}^n r_j u_j \in HH^2_{\boldsymbol{G}}(A, A \otimes R/\mathfrak{m}^2)$$

  where Φ *is the HKR map (see Definition 2.5.1).*

Remark 6.1.3. The main result of [16] was the zeroth-order part of Proposition 6.1.2. It remains to prove that flipping pearly trees can be made to work to first order, and that the first-order deformation classes are as claimed.

6.2. **Flipping Pearly trees.** For the purposes of this section, $\boldsymbol{L}$ will denote a tuple of objects of $\mathcal{F}'$ (i.e., it consists only of two types of entries: L (representing the Lagrangian immersion $L^n_\epsilon : S^{n-2} \to M$ for some $\epsilon > 0$) or L' (representing the Lagrangian immersion $L' : S^{n-2} \to M$).

Definition 6.2.1. If T is a semi-stable directed planar tree with labels $\boldsymbol{L}$, we introduce the following notation:

- $V(T)$ is the set of vertices of T;
- $E(T)$ is the set of edges of T;
- $E'(T) \subset E(T)$ is the subset of edges with both sides labeled L';
- $F'(T)$ is the set of flags (v, e) of T such that $e \in E'(T)$;
- $C(T)$ is the set of 'segments' between consecutive edges around a vertex (these are indexed by pairs of consecutive flags around a vertex);
- If $C \in C(T)$, then $L_C \in \boldsymbol{L}$ is the label associated to C.

Definition 6.2.2. Let $\boldsymbol{L}$ be a tuple. We denote by $\mathcal{R}_3(\boldsymbol{L})$ the moduli space of **flipping pearly trees**, where a flipping pearly tree $r \in \mathcal{R}_3(\boldsymbol{L})$ consists of the following data:

- A semi-stable directed planar tree T_r with labels $\boldsymbol{L}$, such that all internal edges have both sides labeled L' (i.e., all internal edges are contained in $E'(T_r)$);
- A designation of each edge $e \in E'(T_r)$ as either **flipping** or **non-flipping**;
- For each stable vertex $v \in V(T_r)$, a point $r_v \in \mathcal{R}(\boldsymbol{L}_v, \phi)$;
- For each internal edge e, a length parameter $l_e \in [0, \infty)$.

See Figure 6.2.1 for a picture of a flipping pearly tree. We also allow one special case: if $\boldsymbol{L} = (L', L')$, then we permit T_r to have no vertices, just a single edge with both sides labeled L'.

Now, given a flipping pearly tree $r \in \mathcal{R}_3(\boldsymbol{L})$, we define an associated topological space S_r. There are a few special cases first:

Definition 6.2.3. If $\boldsymbol{L} = (L', L')$, and T_r is the tree with a single edge, then we define $S_r := \mathbb{R}$. If $|\boldsymbol{L}| = 2$ but $\boldsymbol{L} \neq (L', L')$, then $S_r := \mathbb{R} \times [0, 1]$.

Now we define S_r in the remaining cases:

Definition 6.2.4. Given a flipping pearly tree $r \in \mathcal{R}_3(\boldsymbol{L})$, we define a topological space S_r as follows:



FIGURE 6.2.1. A flipping pearly tree S. Observe that all edges have label L' on either side.

- For each semi-stable vertex $v \in V(T_r)$ with both sides labeled L', we define S_v to be a disk with two boundary marked points, corresponding to the edges incident to v.
- For each stable vertex $v \in V(T_r)$, we define S_v to be the boundary-marked disk with modulus r_v, with all marked points punctured except for those corresponding to edges in $E'(T_r)$ (they remain as marked points). These are the '**pearls**'.
- We define

$$S^p := \coprod_{v \in V(T_r)} S_v.$$

- For each internal edge e, we define $S_e := [0, l_e]$. For each external edge e in $E'(T_r)$, we define $S_e := \mathbb{R}^{\pm}$, with the $+$ or $-$ depending on the orientation of the edge.
- We define

$$S^e := \coprod_{e \in E(r)} S_e.$$

- For each flag $f = (v, e) \in F'(T_r)$, there is a corresponding marked boundary point $m(f) \in S_v$ and boundary point $b(f) \in S_e$.
- We define

$$S_r := (S^p \sqcup S^e)/\sim$$

  where

$$m(f) \sim b(f) \text{ for all } f \in F'(T_r).$$

Definition 6.2.5. Given $r \in \mathcal{R}_3(\boldsymbol{L})$, we also define a 'boundary' $(\partial S)_r$ and a continuous map

$$(\partial S)_r \to S_r,$$

as follows:

- For each segment $C \in C(T_r)$ adjacent to vertex $v \in V(T_r)$, we define $(\partial S)_C$ to be the corresponding component of the boundary of S_v. Thus, $(\partial S)_C$ is an interval, and its two ends correspond to consecutive marked points on the disk with modulus r_v. If the marked point is punctured in S_v (i.e., if its sides are not both labeled L'), then that end of the interval $(\partial S)_C$ is open, and if the marked point remains in S_v (i.e., if its sides are both labeled L'), then that end of the interval $(\partial S)_C$ is closed.
- We define
$$\begin{aligned} (\partial S)^p &:= \coprod_{C \in C(T_r)} (\partial S)_C, \text{ with the obvious map} \\ (\partial S)^p &\to S^p. \end{aligned}$$
- For each edge e, we define $(\partial S)_e := S_e \times \{0,1\}$ (two copies of S_e).
- We define
$$\begin{aligned} (\partial S)^e &:= \coprod_{e \in E(T_r)} (\partial S)^e, \text{ with the obvious map} \\ (\partial S)^e &\to S^e. \end{aligned}$$
- For each flag $f = (v,e) \in F'(T_r)$, there are points $\tilde{m}_j(f) \in (\partial S)_v$ for $j = 0,1$, from the boundary components to the right and left of $m(f)$ respectively, and points $\tilde{b}_j(f) = (b(f), j) \in (\partial S)_e$, for $j = 0,1$.
- We define
$$(\partial S)_r := ((\partial S)^p \sqcup (\partial S)^e)/\sim,$$
where
$$\tilde{m}_j(f) \sim \tilde{b}_j(f) \text{ for all } f \in F'(r), \text{ and } j = 0,1$$
(see Figure 6.2.2).
- It is clear that there is a continuous map
$$(\partial S)_r \to S_r.$$

Definition 6.2.6. An **automorphism** of a flipping pearly tree is a map $S_r \to S_r$ such that each pearl gets sent to itself by a biholomorphism which preserves the marked points, and each edge gets sent to itself by a translation preserving marked points (in particular, edges are fixed by any automorphism, unless they are infinite).

In particular, the possible non-trivial automorphisms of a flipping pearly tree are:

- If $\boldsymbol{L} = (L', L')$ and $S_r = \mathbb{R}$, then automorphisms are translations of $\mathbb{R}$;
- If $|\boldsymbol{L}| = 2$ but $\boldsymbol{L} \neq (L', L')$, then $S_r = \mathbb{R} \times [0,1]$ and automorphisms are translations in the $\mathbb{R}$-direction;
- If v is a semi-stable vertex of $T(r)$, then S_v is a disk with two marked boundary points, and there is an $\mathbb{R}$ family of automorphisms (translations) of S_v preserving the marked boundary points.



FIGURE 6.2.2. Defining the boundary $\partial S \to S$ of a flipping pearly tree S, and attaching a strip (shaded in grey) along it.

Definition 6.2.7. From our universal choice of strip-like ends for the moduli spaces $\mathcal{R}(\boldsymbol{L}, \phi)$, we can define a subset

$$S^p_{\text{thin}} \subset S^p,$$

called the **thin** region, which consists of the images of strip-like ends under gluing maps (see [3, Remark 9.1]). To clarify: the thin region includes a neighbourhood of each boundary marked point of a pearl, and also **all** of any semi-stable pearl S_v (see Figure 6.2.3). We define the corresponding **thick** region

$$S^p_{\text{thick}} := S^p \setminus S^p_{\text{thin}}.$$

Definition 6.2.8. We define the region

$$S^e_{\text{thin}} \subset S^e$$

to be the set of points on edges which are distance > 1 from the boundary of the edge, and

$$S^e_{\text{thick}} := S^e \setminus S^e_{\text{thin}}$$

(see Figure 6.2.3).

As in [16, Section 4.1], we can define a topology on the moduli space $\mathcal{R}_3(\boldsymbol{L})$. The important point is that thin regions with opposite sides labelled L' can stretch until they 'break', then



Figure 6.2.3. The thick and thin regions of the same flipping pearly tree S illustrated in Figure 6.2.2. The thick regions are shown in light grey, and the thin regions in dark grey. Note that the unstable disk (with two marked points) is entirely thin, and also that it is possible for an internal edge to be entirely thick (when it has length ≤ 2).

become an internal edge (see Figure 6.2.4). The difference from [16] is that we now allow semi-stable vertices, and this means that $\mathcal{R}_3(\boldsymbol{L})$ no longer has the structure of a manifold with boundary. All we can say is that it is stratified by manifolds – however we will see later that the space of holomorphic maps of pearly trees into our manifold **is** a manifold with boundary, which is what we need to define our algebraic structures.

The strata of $\overline{\mathcal{R}}_3(\boldsymbol{L})$ are indexed by semi-stable directed planar trees T with labels $\boldsymbol{L}$. Note there is no requirement that internal edges have opposite sides labeled L' here. The tree T corresponds to the codimension-$(|V(T)|-1)$ stratum

$$\mathcal{R}_3^T(\boldsymbol{L}) \cong \prod_{v \in V(T)} \mathcal{R}_3(\boldsymbol{L}_v).$$

Points in this stratum correspond to flipping pearly trees, where the pearls are allowed to be nodal and the edges are allowed to have infinite length.



FIGURE 6.2.4. A thin region with both sides labelled L' (upper left) can stretch until it becomes a thin region (upper right), then break, becoming a thick internal edge (lower right), which then stretches until it has a thin region in its interior (lower left).

Remark 6.2.9. The moduli space $\overline{\mathcal{R}}_3(\boldsymbol{L})$ is not compact, because our flipping pearly trees can have arbitrarily many semi-stable vertices. In practice (see the proof of Proposition 6.4.1), we have an a priori upper bound N on the number of semi-stable vertices that we need consider. We consider the subspace

$$\overline{\mathcal{R}}_3(\boldsymbol{L}, N) \subset \overline{\mathcal{R}}_3(\boldsymbol{L}),$$

consisting of stable flipping pearly trees with $\leq N$ semi-stable vertices. This subspace **is** compact.

6.3. **Floer and perturbation data.**

Definition 6.3.1. We define

$$\mathcal{H}^e := C^\infty(S^n, \mathbb{R})$$

(think of this as the space of Morse functions on S^n), and

$$\mathcal{H}^p := \{H \in C^\infty(M, \mathbb{R}) : H \text{ vanishes, with its first derivatives, along each divisor } D_j\}$$

(think of this as the space of Hamiltonians on M), and $\mathcal{J}$, the space of smooth almost-complex structures on M which are compatible with ω, and make the divisors D_j almost-complex submanifolds.

Definition 6.3.2. Let $\boldsymbol{L} = (L_0, L_1)$ be a 2-element tuple. For each such tuple, we choose a Floer datum $(H_{\boldsymbol{L}}, J_{\boldsymbol{L}})$ consisting of

$$H_{\boldsymbol{L}} \in C^\infty([0,1], \mathcal{H}^p) \text{ and } J_{\boldsymbol{L}} \in C^\infty([0,1], \mathcal{J})$$

such that:

- $H_{\boldsymbol{L}} = 0$ unless $\boldsymbol{L} = (L, L)$;
- the time-1 Hamiltonian flow of $H_{(L,L)}$ makes L^n transverse to itself;
- $J_{(L',L')} = J_0$ is constant, equal to the standard integrable complex structure.

Now, if $(L_0, L_1) \neq (L', L')$, then we define a **generator** of $CF^*_{\mathcal{F}'}(L_0, L_1)$ to be a path $p : [0,1] \to M$ which is a flowline of the Hamiltonian vector field of $H_{\boldsymbol{L}}$, such that $p(0) \in L_0$ and $p(1) \in L_1$. One defines $CF^*_{\mathcal{F}'}(L_0, L_1)$ to be the $R/\mathfrak{m}^2$-module generated by its generators. It is $\boldsymbol{G}$-graded.

Remark 6.3.3. The $\boldsymbol{G}$-grading is defined exactly as in the affine Fukaya category (Section 3.3). One might worry that L' intersects the divisors D_j, hence doesn't lie in $M \setminus \boldsymbol{D}$. However, we simply push L' off itself using ∇f (as in the definition of L^n_ϵ), and use the pushed-off version of L' for all grading (and index) computations – see [16, Proof of Proposition 5.3]. The point is that the grading of the relative Fukaya category arises from index computations of the relevant Fredholm operators, which are computed purely topologically.

Definition 6.3.4. If $(L_0, L_1) = (L', L')$, then we define the Floer datum to contain additional information, namely:

- the Morse function $f : S^n \to \mathbb{R}$;
- another Morse function $h : S^n \to \mathbb{R}$, with exactly two critical points.

One defines a **generator** of $CF^*_{\mathcal{F}'}(L', L')$ to be a critical point of one of the Morse functions f or h, and $CF^*_{\mathcal{F}'}(L', L')$ to be the $R/\mathfrak{m}^2$-module generated by these critical points. We identify the critical points of f as p_K for $K \subset [n]$ proper and non-empty, and the critical points of h as p_ϕ and $p_{[n]}$. Then $CF^*_{\mathcal{F}'}(L', L')$ is $\boldsymbol{G}$-graded, where p_K has the same grading as the corresponding generator θ^K of A, where A is the $\boldsymbol{G}$-graded algebra of Definition 2.2.12.

Lemma 6.3.5. *The $\boldsymbol{G}$-graded morphism spaces in $\mathcal{F}'$ are as follows:*

$$\begin{aligned} CF^*_{\mathcal{F}'}(L, L) &\cong CF^*_{\mathcal{F}'}(L', L') \\ &\cong A \otimes R/\mathfrak{m}^2 \\ &\cong \bigoplus_{K \subset [n]} R/\mathfrak{m}^2 \cdot p_K, \end{aligned}$$

and

$$\begin{aligned} CF^*_{\mathcal{F}'}(L, L') &\cong CF^*_{\mathcal{F}'}(L', L) \\ &\cong A \otimes R/\mathfrak{m}^2 \oplus CM^*(f) \otimes R/\mathfrak{m}^2 \\ &\cong \bigoplus_{K \subset [n]} R/\mathfrak{m}^2 \cdot p_K \oplus \bigoplus_{K \subset [n], K \neq \phi, [n]} R/\mathfrak{m}^2 \cdot q_K, \end{aligned}$$

where $CM^*(f)$ is the Morse complex of the function f. The $\boldsymbol{G}$-grading of generators labelled by $p_K \in A$ is as in Definition 2.2.12. The $\boldsymbol{G}$-grading of generators labelled by q_K is $(|K|-1, 0)$ (where $|K|-1$ is the Morse index of q_K).

Proof. See [16, Proof of Proposition 5.5]. □

Definition 6.3.6. In each morphism space, we call the generators labelled p_K, where $K \subset [n]$ is not equal to ϕ or $[n]$, the **flipping generators**, and the others (labelled q_K, p_ϕ or $p_{[n]}$) the **non-flipping generators**. The terminology comes from the definition of L^n as a perturbation of the double cover $S^n \to \mathbb{RP}^n \hookrightarrow \mathbb{CP}^n$. Flipping generators correspond to paths p from one sheet of the cover to the opposite sheet; non-flipping generators correspond to paths from one sheet to the same sheet.

Definition 6.3.7. A **perturbation datum** for a fixed flipping pearly tree $r \in \mathcal{R}_3(\boldsymbol{L})$ consists of the data (K^e, K^p, J), where:

- $K^e \in C^\infty(S^e, \mathcal{H}^e)$;
- $K^p \in \Omega^1(S^p, \mathcal{H}^p)$;
- $J \in C^\infty(S^p, \mathcal{J})$,

such that, for each boundary component C of a pearl in S with Lagrangian label L_C,

$$K^p(\xi)|_{L_C} = 0 \text{ for all } \xi \in TC \subset T(\partial S).$$

Definition 6.3.8. We say that a perturbation datum is **compatible with the Floer data** if, on each component of the thin regions of S^p and S^e, the perturbation datum agrees with the corresponding (translation-invariant) Floer datum. Explicitly, this means that:

- On each strip-like end of a pearl, (K^p, J) is given by the translation-invariant extension of the Floer datum $(H_{\boldsymbol{L}}, J_{\boldsymbol{L}})$;
- In a neighbourhood of each boundary marked point of a pearl, and also on all of each semi-stable pearl with both sides labeled L', we have $(K^p, J) = (0, J_0)$;
- On each thin region of a flipping edge, $K^e = f$;
- On each thin region of a non-flipping edge, $K^e = h$.

Remark 6.3.9. Note that, if our perturbation datum is compatible with the Floer data, then it is preserved by any automorphism of the flipping pearly tree.

Definition 6.3.10. We define the notion of a **compatible universal choice of perturbation data** for the moduli spaces $\mathcal{R}(\boldsymbol{L})$, by analogy with [3, Section 9i].

Definition 6.3.11. Let $\boldsymbol{L}$ be a tuple of objects of $\mathcal{F}'$, and $\boldsymbol{p}$ an associated set of generators. A **holomorphic flipping pearly tree $\boldsymbol{u}$** with ends on $\boldsymbol{p}$ consists of the following data:

- A flipping pearly tree $r \in \mathcal{R}_3(\boldsymbol{L})$;
- For each vertex $v \in V(T(r))$, a smooth map $u_v : S_v \to M$;
- For each edge $e \in E(r)$ with both sides labeled L', a smooth map $u_e : S_e \to S^n$;
- A continuous map $\tilde{u} : \partial S_r \to S^n$.

We impose the following requirements on these maps:

- $\boldsymbol{u}$ is asymptotic to the generators $\boldsymbol{p}$ along the strip-like ends and external edges;
- For each semi-stable vertex v, the map u_v satisfies the perturbed holomorphic curve equation
$$(Du_v - Y)^{0,1} = 0,$$
where, for $\xi \in TS$, $Y(\xi)$ is the Hamiltonian vector field of the function $K^p(\xi)$;
- The maps u_e satisfy the Morse flow equation
$$Du_e - \nabla K^e \quad = \quad 0;$$
- For each boundary component C of a pearl S_v,
$$L_C \circ \tilde{u}|_C = u_v|_C;$$

- For each edge e with both sides labeled L',
  $$\tilde{u}|_{S_e \times \{0\}} = u_e,$$
  and
  $$\tilde{u}|_{S_e \times \{1\}} = \begin{cases} u_e & \text{if } e \text{ is non-flipping} \\ a \circ u_e & \text{if } e \text{ is flipping} \end{cases}$$
  where we recall that $a : S^n \to S^n$ is the antipodal map.
- If $v \in V(T(r))$ is semi-stable, then the map u_v is non-constant.

Definition 6.3.12. Two holomorphic flipping pearly trees are **equivalent** if they are related by an automorphism of the domain (recall from Remark 6.3.9 that any automorphism of the domain preserves the perturbation datum and hence acts on the space of holomorphic flipping pearly trees).

Definition 6.3.13. Given a flipping holomorphic pearly tree $\boldsymbol{u}$ as defined above, one obtains a well-defined homology class $[\boldsymbol{u}] \in H_2(M, L^n)$ as follows (see Figure 6.2.2):

- Start with the continuous map $\boldsymbol{u} : S \to \mathbb{CP}^n$ associated with the flipping holomorphic pearly tree.
- Glue a thin strip along the boundary ∂S of the flipping pearly tree;
- If the boundary component or edge has label L, then it already gets mapped to L^n, so we map the strip into $\mathbb{CP}^n$ by making it constant along its width.
- If the boundary component or edge has label L', then by construction, there is a continuous lift $\tilde{u}$ of the boundary of the strip to S^n.
- Thus, we can map the strip into $\mathbb{CP}^n$ by letting it interpolate between the zero section and the graph of ϵdf in the Weinstein neighbourhood $D^*_\eta S^n$ used in the construction of L^n. Thus, boundary components of the strip with label L' now lie on L^n.

We now define the intersection number $\boldsymbol{u} \cdot D_j$ to be the topological intersection number of this class $[\boldsymbol{u}] \in H_2(\mathbb{CP}^n, L^n)$ with $D_j \in H_{2n-2}(\mathbb{CP}^n)$, and
$$\boldsymbol{u} \cdot \boldsymbol{D} := \sum_j ([\boldsymbol{u}] \cdot D_j)\, y_j \in Y.$$

Definition 6.3.14. Let $\boldsymbol{u}$ be a holomorphic flipping pearly tree. For each $v \in V(T(r))$, the map u_v defines a homology class in $H_2(\mathbb{CP}^n, \mathbb{RP}^n) \cong \mathbb{Z}$, because its boundary gets mapped to a Weinstein neighbourhood of $\mathbb{RP}^n$. We denote this homology class by $d_v \in \mathbb{Z}_{\geq 0}$ (it is non-negative because holomorphic disks have non-negative area). We denote the sum of all homology classes d_v by $d_{\boldsymbol{u}} \in \mathbb{Z}_{\geq 0}$.

Now we explain how to compute these intersection numbers in a simple way. It helps if the holomorphic flipping pearly trees are in general position, in the following sense:

Definition 6.3.15. We say that a holomorphic flipping pearly tree $\boldsymbol{u}$ is **in general position** if:

- Each boundary component C with label L' is transverse to the real hypersurfaces $D_j^{\mathbb{R}} \subset S^n$;
- No flipping marked points lie on the hypersurfaces $D_j^{\mathbb{R}}$.

Lemma 6.3.16. *Given a holomorphic flipping pearly tree $\boldsymbol{u}$, we can perturb the defining equations of the divisors D_j so that*

- *The intersection numbers $\boldsymbol{u} \cdot D_j$ do not change;*
- *$\boldsymbol{u}$ is in general position with respect to the perturbed divisors.*

Proof. See the proof of [16, Proposition 5.1]. □

If $\boldsymbol{u}$ is in general position, then we can split the surface defining our homology class $[\boldsymbol{u}]$ into regions $[u_v]$ corresponding to the pearls v, and $[u_e]$ corresponding to the edges e of the pearly tree, in such a way that the boundary of each such region does not intersect the divisors D_j. We cut the pearls off from the strips in the obvious way – since they are joined at boundary marked points, which don't lie on the hypersurfaces $D_j^{\mathbb{R}}$, the cuts we introduce do not intersect the divisors D_j.

Thus, each region defines a class in $H_2(M, M \setminus \boldsymbol{D})$, and $[\boldsymbol{u}] \cdot \boldsymbol{D}$ is equal to the sum of $[u_v] \cdot \boldsymbol{D}$ and $[u_e] \cdot \boldsymbol{D}$ over all pearls v and edges e of the pearly tree.

Lemma 6.3.17. *Let $\boldsymbol{u}$ be a holomorphic flipping pearly tree in general position. Then we have:*

- *For each non-flipping edge e, $[u_e] \cdot D_j = 0$;*
- *For each flipping edge e, $[u_e] \cdot D_j$ is equal to the topological intersection number of the edge $u_e : [0, l_e] \to S^n$ with the hypersurface $D_j^{\mathbb{R}}$ (this is non-negative because the gradient of the function f crosses all hypersurfaces $D_j^{\mathbb{R}}$ positively);*
- *For each pearl v, $[u_v] \cdot D_j$ is equal to the sum of the number of internal intersections of u_v with D_j (these are counted positively by positivity of intersections), together with $+1$ for each time a boundary lift $\tilde{u}|_C$ with label L' crosses $D_j^{\mathbb{R}}$ in the negative direction (and 0 if the lift crosses in the positive direction).*

Proof. See [16, Proposition 5.1]. □

Corollary 6.3.18. *If $\boldsymbol{u}$ is a flipping holomorphic pearly tree, then the intersection numbers $\boldsymbol{u} \cdot D_j$ are non-negative.*

Definition 6.3.19. Let $\boldsymbol{L}$ be a tuple, $\boldsymbol{p}$ an associated set of generators, and $\boldsymbol{d} \in Y_{\geq 0}$. We define $\mathcal{M}_3(\boldsymbol{p}, \boldsymbol{d})$ to be the moduli space of holomorphic flipping pearly trees $\boldsymbol{u}$ with labels $\boldsymbol{L}$ and ends on $\boldsymbol{p}$, and such that

$$\boldsymbol{u} \cdot \boldsymbol{D} = \boldsymbol{d},$$

modulo equivalence.

Proposition 6.3.20. *Let $\boldsymbol{L}$ be a tuple, $\boldsymbol{p}$ an associated set of generators. Suppose that $(p_{K_1}, \ldots, p_{K_k})$ is the tuple obtained from $\boldsymbol{p}$ by keeping only the flipping generators (see Definition 6.3.6). Then we have*

$$[\boldsymbol{u}] \cdot \boldsymbol{D} = q y_{[n]} + \sum_{j=1}^{k} y_{K_j}$$

for some $q \in \mathbb{Z}$, and the homology class of $[\boldsymbol{u}] \in H_2(\mathbb{CP}^n, \mathbb{RP}^n) \cong \mathbb{Z}$ is given by the formula

$$[\boldsymbol{u}] = 2q + k.$$



FIGURE 6.3.1. An example of a legal labeling of a flipping holomorphic pearly tree, which might contribute to the coefficient of p_ϕ in the A_∞ product $\mu^7(p_{\{1\}}, \ldots, p_{\{7\}})$. We have illustrated a simple case, in which all external flowlines are constant because the points $p_{\{j\}}$ are maxima of the Morse function f. The external label '1' means the set $\{1\}$, while '$\bar{1}$' means the complement $\{2, 3, 4, 5, 6, 7\}$. The big label '1' in the middle of a pearl means that the pearl has degree 1.

Proof. Follows from [16, Lemma 5.8], and a slight modification of [16, Proposition 5.10]. □

Definition 6.3.21. Let $\boldsymbol{u}$ be a holomorphic flipping pearly tree in general position, all of whose boundary components are labelled L'. For each flag f corresponding to a flipping edge of the tree, we have corresponding marked points in the boundary $(\partial S)_r$, namely

$$\tilde{m}_0(f) \sim \tilde{b}_0(f) \text{ and } \tilde{m}_1(f) \sim \tilde{b}_1(f).$$

The boundary map $\tilde{u}$ sends these points to antipodal regions $S^n_K, S^n_{\bar{K}}$ respectively, for some $K \subset [n]$ (recall that S^n_K is defined to be the region where $x_j < 0$ for $j \in K$ and $x_j > 0$ for $j \notin K$). We attach the labels K and $\bar{K}$ to the marked points $\tilde{m}_0(f)$ and $\tilde{m}_1(f)$, respectively. Figure 6.3.1 shows a possible labeling of a flipping holomorphic pearly tree.

Lemma 6.3.22. *Let $\boldsymbol{u}$ be a holomorphic flipping pearly tree with all sides labelled L', in general position, and equipped with labels as above. We decompose $[\boldsymbol{u}]$ into regions $[u_e]$, corresponding to edges, and $[u_v]$, corresponding to pearls, as before. Then we have:*

- *For each flipping edge e, the label K_0 at the start of the edge contains the label K_1 at the end of the edge, and*
$$[u_e] \cdot \boldsymbol{D} = y_{K_1} - y_{K_0};$$
- *For each pearl v such that:*
  - *The homology class is $d_v \in H_2(\mathbb{CP}^n, \mathbb{RP}^n) \cong \mathbb{Z}$;*
  - *The points immediately after the flipping marked points of u_v have labels $K_1, \ldots, K_{k_v}$, if we traverse the boundary in positive direction (in other words, the points $\tilde{m}_1(f)$ for all outgoing flags, and $\tilde{m}_0(f)$ for the incoming flag);*

  then we have:
  - *For some $q_v \in \mathbb{Z}$,*
$$[\boldsymbol{u}] \cdot \boldsymbol{D} = q_v y_{[n]} + \sum_{j=1}^{k_v} y_{K_j};$$

- *The homology class* d_v *satisfies*

$$d_v = 2q_v + k_v.$$

We recall that non-flipping edges u_e *do not contribute to the intersection numbers.*

Proof. For edges, the statement follows from the fact that ∇f only crosses divisors positively, and $[u_e]$ picks up an intersection point with D_j each time u_e crosses D_j positively (compare [16, Figure 9(b)]). For pearls, the statement follows from Proposition 6.3.20. $\square$

6.4. **Transversality and compactness.**

Proposition 6.4.1. *For generic choice of Floer and perturbation data, the components of* $\mathcal{M}_3(\boldsymbol{p}, \boldsymbol{d})$ *with* $|\boldsymbol{d}| \le 1$ *are regular, and have the structure of topological manifolds of the expected dimension.*

Proof. The moduli spaces $\mathcal{M}_3(\boldsymbol{p}, \boldsymbol{d})$ are constructed by gluing together pieces corresponding to the different possible underlying trees (see [16, Section 4.4]). If we are to obtain a topological manifold, we need each piece to be cut out transversely, and also for the 'seams' along which the pieces are glued (corresponding to holomorphic flipping pearly trees with nodal pearls, where a Morse flowline is about to form as in Figure 6.2.4) to be regular.

This amounts to checking the following:

- The Cauchy-Riemann operator $(Du^p - Y)^{0,1}$ on each stable pearl is surjective;
- For the semi-stable pearls, we require that the moduli space of J_0-holomorphic disks with two boundary marked points, modulo translation, is regular;
- The Morse flow operator $(Du^e - \nabla K^e)$ on each edge is surjective;
- For moduli spaces consisting of a single Morse edge, we require that the moduli space of Morse flowlines of f or h is regular;
- The restriction that marked points on pearls coincide with ends of edges is cut out transversely;
- The restriction that marked points on pearls coincide at a node (with both sides labelled L') is cut out transversely (this is the requirement that the 'seams' are regular).

The Cauchy-Riemann operators on pearls corresponding to stable vertices are generically surjective, by perturbing K^p (as in [3, Section 9k]). The moduli spaces of semi-stable pearls with sides labelled anything other than (L', L') are regular for generic choice of Floer data, by the arguments of [35, 36]. The semi-stable pearls with opposite sides labelled L' are different – we have required that $K^p = 0$ and $J = J_0$ on these pearls, which is not a generic condition. However, the moduli space of such pearls is regular by the 'automatic regularity' result of [31, Proposition 7.4.3] (the automatic regularity result deals with holomorphic spheres in $\mathbb{CP}^n = M$, and our moduli space of holomorphic disks is the real locus of this moduli space, hence also regular).

The Morse flow operators are surjective for generic choice of Floer and perturbation data.

The intersections of marked points on pearls with the endpoints of edges are generically transverse, by perturbing K^e near the end of the edge.

Now we deal with intersections of marked points at a node connecting two pearls, which are necessary to make the 'seams' along which we glue different parts of our moduli space. If one of the pearls involved is stable, then the intersection is transverse, because we can perturb K^p on the stable pearl (see [16, Section 4.6]) to move the marked point in any direction we please. A problem arises if both pearls are semi-stable (with both sides labelled L'). However, this situation does not arise in the moduli spaces $\mathcal{M}_3(\boldsymbol{p},\boldsymbol{d})$ with $|\boldsymbol{d}| \le 1$: any semi-stable pearl must contribute at least 1 to one of the intersection numbers $\boldsymbol{u} \cdot D_j$.

This follows from Lemma 6.3.22: if $[\boldsymbol{u}] \cdot \boldsymbol{D} = 0$ for a pearl with two boundary marked points with labels K_0 and K_1, then

$$qy_{[n]} + y_{K_0} + y_{K_1} = 0.$$

If $K_0 = \phi$ or $[n]$, then Lemma 6.3.22 shows that $d_v = 0$ so the unstable pearl u_v has zero energy and must be constant, which is not allowed. If $K_0 \neq \phi$ or $[n]$, then necessarily $q = -1$, and Lemma 6.3.22 shows that

$$d_v = 2q_v + k_v = 0,$$

so again u_v is constant. □

Proposition 6.4.2. *For generic choice of Floer and perturbation data, the components of $\mathcal{M}_3(\boldsymbol{p},\boldsymbol{d})$ of virtual dimension ≤ 1, and with $|\boldsymbol{d}| \le 1$, have the structure of compact topological manifolds with boundary, of the expected dimension. The boundary strata of the one-dimensional moduli spaces correspond to nodal holomorphic flipping pearly trees (see* [16, Definition 4.33, 4.34]*).*

Proof. This essentially follows from Gromov compactness, as outlined in [16, Proposition 4.6]. However, we must do a bit more work in this case: we must check that the number of semi-stable pearls is bounded, so that we are only gluing together finitely many pieces to make our moduli space (see Remark 6.2.9). This is true because the homology class $d_{\boldsymbol{u}}$ of any $\boldsymbol{u} \in \mathcal{M}_3(\boldsymbol{p},\boldsymbol{d})$ is fixed, by Proposition 6.3.20, and any semi-stable pearl v contributes at least 1 to $d_{\boldsymbol{u}}$, because it is required to be non-constant.

Furthermore, we must rule out the possibility of sphere and disk bubbling in our moduli spaces. Sphere bubbling is easy to rule out: any non-constant sphere bubble must intersect each divisor D_j at least once, hence contribute at least n to $\boldsymbol{d}$. So sphere bubbling does not happen in moduli spaces with $|\boldsymbol{d}| \le 1$.

Disk bubbling needs a little more work. Suppose a holomorphic disk bubbles off some pearl in a holomorphic flipping pearly tree. Let us denote by u_1 the disk, and by u_2 the rest of the holomorphic flipping pearly tree. We will assume that u_1 has boundary on L' (the case with boundary on L^n is not very different). We can regard this configuration as a holomorphic flipping pearly tree with a vertex v of valence 1, connected by a Morse edge of length 0 to the rest of the holomorphic flipping pearly tree. We will show that, if u_1 is non-constant, then its virtual dimension (by which we mean the virtual dimension of the moduli space in which it lies) is $\ge n+1$. This is the same as showing that its Maslov index is at least 3. This will show that disk bubbling generically does not happen, because the rest of the holomorphic flipping pearly tree has virtual dimension < 0.

Firstly, by Lemma 6.3.22 (which works exactly the same if there are pearls of valence 1), if the edge is non-flipping then the disk has zero energy, and hence is constant.

So let us assume that the edge is flipping, and the label attached to it is $K \subset [n]$ (as in Definition 6.3.21). By Lemma 6.3.22, we have

$$[u_1] \cdot \boldsymbol{D} = q y_{[n]} + y_K.$$

Because $|\boldsymbol{d}| \leq 1$, it must be that $[u_1] \cdot \boldsymbol{D} = y_j$, $q = 0$, and $K = \{j\}$ for some $j \in [n]$. It follows, by Lemma 6.3.22, that the homology class of u_1 is $d_1 = 1$. Thus, the virtual dimension of u_1 is (by the real analogue of Lemma 4.4.3)

$$\text{v.d.}(u_1) = n + d_1(n+1) - 3 + 1 = 2n - 1 \geq n + 1$$

(because $n \geq 2$).

The connect sum formula now says that the virtual dimension of $\boldsymbol{u}$ is

$$\text{v.d.}(\boldsymbol{u}) = \text{v.d.}(u_1) + \text{v.d.}(u_2) + 1 - n \leq 1$$

(since we are considering moduli spaces of virtual dimension ≤ 1), and hence

$$\text{v.d.}(u_2) \leq n - \text{v.d.}(u_1) \leq -1.$$

We saw in Proposition 6.4.1 that moduli spaces of holomorphic flipping pearly trees are generically regular, hence a moduli space of virtual dimension < 0 is generically empty. So we may conclude that no disk bubbling occurs for generic choices of Floer and perturbation data. $\square$

6.5. **Morse-Bott model for the first-order Fukaya category.** This section contains the proof of Proposition 6.1.2.

We define a $\boldsymbol{G}$-graded A_∞ category $\mathcal{F}'$ over $R/\mathfrak{m}^2$, with two objects, L and L'. The $\boldsymbol{G}$-graded morphism spaces $CF^*(L_0, L_1)$ are the $R/\mathfrak{m}^2$-module freely generated by their generators, and are described explicitly in Lemma 6.3.5. Given a tuple $\boldsymbol{L}$ with associated generators $\boldsymbol{p} = (p_0, p_1, \ldots, p_s)$, the coefficient of $r^{\boldsymbol{d}} p_0$ in the A_∞ product $\mu^s(p_s, \ldots, p_1)$ is given by the signed count of holomorphic flipping pearly trees in the zero-dimensional component of the moduli space $\mathcal{M}_3(\boldsymbol{p}, \boldsymbol{d})$ (by analogy with the usual definition of the Fukaya category). By the standard argument, the composition maps μ^s satisfy the A_∞ relations.

Furthermore, they are $\boldsymbol{G}$-graded, by the same argument as for the relative Fukaya category (keeping in mind Remark 6.3.3). To see why, recall that the grading depended on the index theory of a Cauchy-Riemann operator coming from a bundle pair (D^2, u^*TM, F), where $u : D^2 \to M \setminus \boldsymbol{D}$ was some smooth map, and F a lift of ∂u to $\mathcal{G}(M \setminus \boldsymbol{D})$. Given a holomorphic flipping pearly tree $\boldsymbol{u}$, we can construct (by a modification of the construction of the smooth surface representing the homology class $[\boldsymbol{u}]$) a smooth boundary-punctured disk $\tilde{\boldsymbol{u}}$ mapping to $M \setminus \boldsymbol{D}$, with boundary on L^n, such that the index of the associated Cauchy-Riemann operator is equal to the index of the Fredholm operator defining the moduli space of holomorphic flipping pearly trees near $\boldsymbol{u}$. Note that $\tilde{\boldsymbol{u}}$ will **not** be holomorphic itself: we are simply using it to compute the index topologically.

Lemma 6.5.1. *For small enough $\epsilon > 0$, the objects L^n_ϵ and L' are quasi-isomorphic in $\mathcal{F}'/\mathfrak{m}$.*

Proof. See [16, Proposition 5.5]. $\square$

Lemma 6.5.2. *For appropriate choices of perturbation data, there is a strict isomorphism*

$$CF^*_{\mathcal{F}(M,\boldsymbol{D})/\mathfrak{m}^2}(L^n, L^n) \cong CF^*_{\mathcal{F}'}(L, L).$$

Proof. Suppose that $\boldsymbol{L}$ is a tuple, all of whose entries are L^n, $|\boldsymbol{L}| \geq 3$, and ℓ is a labelling such that $\boldsymbol{d}(\ell) = y_j$ for some j. Then there is a forgetful map $\mathcal{R}(\boldsymbol{L}, \ell) \to \mathcal{R}_3(\boldsymbol{L})$, obtained by simply forgetting the internal marked point. Suppose that we choose Floer and perturbation data on the moduli space $\mathcal{R}(\boldsymbol{L}, \ell)$ by pulling back the corresponding data from $\mathcal{R}_3(\boldsymbol{L})$.

With these assumptions, if $\boldsymbol{p}$ is a set of generators associated with the tuple $\boldsymbol{L}$, then there is a forgetful map $\mathcal{M}(\boldsymbol{p}, \ell) \to \mathcal{M}_3(\boldsymbol{p}, e_j)$. This map is clearly bijective: given a holomorphic disk with topological intersection number 1 with the divisor D_j, there exists a unique internal point which gets mapped to D_j (uniqueness follows from positivity of intersections).

If $|\boldsymbol{L}| = 2$, we choose the perturbation data on the moduli space $\mathcal{R}(\boldsymbol{L}, \ell)$ to be translation-invariant, like for $\mathcal{R}_3(\boldsymbol{L}, \ell)$. The definitions of the two moduli spaces now are different: $\mathcal{M}_3(\boldsymbol{p}, e_j)$ is a quotient of a moduli space of holomorphic strips by translation, whereas $\mathcal{M}(\boldsymbol{p}, \ell)$ is a moduli space of holomorphic strips with an internal marked point getting mapped to D_j. However, this map is still bijective: given a strip with topological intersection number 1 with divisor D_j, there is a unique internal point which gets mapped to D_j. Introducing this as a marked point defines a class in $\mathcal{M}(\boldsymbol{p}, \ell)$, and the result is independent of translation of the domain in the original strip.

Observe that, because the moduli spaces $\mathcal{M}_3(\boldsymbol{p}, e_j)$ are regular, the moduli spaces $\mathcal{M}(\boldsymbol{p}, \ell)$ are regular, even though we have made a non-generic choice of perturbation data. So it is legitimate to compute coefficients in the relative Fukaya category using the moduli spaces $\mathcal{M}(\boldsymbol{p}, \ell)$ defined using these Floer and perturbation data. □

Now we recall (Lemma 6.3.5) that the underlying $\boldsymbol{G}$-graded vector space of $CF^*_{\mathcal{F}'}(L', L')$ is A. So $CF^*_{\mathcal{F}'}(L', L')$ has the form (A, μ^*).

Lemma 6.5.3. *As a $\boldsymbol{G}$-graded algebra, we have*

$$(A, \mu_0^2) \cong (A, \wedge),$$

where '$\wedge$' denotes the usual product on the exterior algebra.

Proof. See [16, Theorem 5.12]. □

Now recall from Section 2.5 that there is a map

$$\Phi : CC^*_{\boldsymbol{G}}(A, A \otimes R) \to p^* \left(\mathbb{C}[[U]] \otimes A \otimes R\right).$$

Lemma 6.5.4. *We have*

$$\Phi\left(\mu^*\right) = u_1 \ldots u_n + \sum_{j=1}^{n} r_j u_j + \mathfrak{m}^2$$

(up to some signs).

Proof. See [16, Proposition 5.15] for the proof that

$$\Phi(\mu_0^*) = \pm u_1 \dots u_n.$$

Our aim now is to calculate the first-order terms in $\Phi(\mu^*)$. By Lemma 2.5.11, we know that the degree-2 part of

$$HH^*_{\boldsymbol{G}}(A, A \otimes R^1) \cong p^* \left(\mathbb{C}[[U]] \otimes A \otimes R^1\right)$$

is generated by the elements $r_j u_j$. Thus, the first-order part of $\Phi(\mu^*)$ can be written as

$$\sum_{j=1}^{n} c_j r_j u_j$$

for some numbers $c_j \in \mathbb{C}$.

The number c_j is given by the count of holomorphic flipping pearly trees $\boldsymbol{u}$ in the moduli space $\mathcal{M}_3((p_\phi, p_{\{j\}}), e_j)$. Such a holomorphic flipping pearly tree must be a chain of semi-stable J_0-holomorphic pearls (see Definition 6.3.8).

By Proposition 6.3.20, the homology class of such $\boldsymbol{u}$ is 1. Because semi-stable pearls are not allowed to be constant, this means there can only be a single semi-stable pearl. This pearl is a J_0-holomorphic disk with boundary on $\mathbb{RP}^n$,

$$u_v : (D, \partial D) \to (\mathbb{CP}^n, \mathbb{RP}^n),$$

together with two marked boundary points, considered up to translation. The homology class $[u_v] \in H_2(\mathbb{CP}^n, \mathbb{RP}^n)$ is 1. Thus, u_v is one half of a J_0-holomorphic sphere of degree 1 in $\mathbb{CP}^n$. That is, it is one half of a complex line in $\mathbb{CP}^n$, and its boundary is a real line in $\mathbb{RP}^n$.

The Morse index of the input $p_{\{j\}}$ is n (see [16, Corollary 2.11]), so the corresponding Morse edge must be constant: this means we just have a point constraint that our real line must pass through the point $L'(p_{\{j\}})$. Similarly, the Morse index of p_ϕ is 0, so the corresponding Morse edge must be constant: this means we have a point constraint that our real line must pass through the point $L'(p_\phi)$. There is a unique real line through the points $L'(p_{\{j\}})$ and $L'(p_\phi)$, so the pearl must be one half of the corresponding complex line.

Furthermore, the pearl must admit a lift $\tilde{u}_v$ of the boundary to S^n, which changes sheets at $p_{\{j\}}$ but not at p_ϕ. Since u_v is one half of a complex line, this lift $\tilde{u}_v$ must be one half of a great circle, from $p_{\{j\}}$ to its antipode $p_{\overline{\{j\}}}$, and passing through the point p_ϕ. Clearly, there is exactly one such half-great circle. Furthermore, the orientation of this half-great circle determines uniquely which half of the complex line we must take. Thus, we have uniquely determined our holomorphic flipping pearly tree $\boldsymbol{u}$.

It follows from Proposition 6.3.20 that $[\boldsymbol{u}] \cdot \boldsymbol{D} = y_j$.

We have shown that the moduli space $\mathcal{M}((p_{\{j\}}, p_\phi), e_j)$ contains a unique element $\boldsymbol{u}$. Thus, each coefficient c_j must be ± 1. □

Remark 6.5.5. We could also have calculated $[\boldsymbol{u}] \cdot \boldsymbol{D}$ using Lemma 6.3.17, and the exercise helps us to get a picture of $\boldsymbol{u}$: the edges u_e are constant, hence do not contribute to $[\boldsymbol{u}] \cdot \boldsymbol{D}$. The pearl u_v does not intersect any divisor D_i in its interior (it is half of a complex line, hence intersects the divisor D_i exactly once, and that intersection is on the real locus $\mathbb{RP}^n$). So we get no contributions to $[\boldsymbol{u}] \cdot \boldsymbol{D}$ from interior intersection points. The boundary lift $\tilde{u}_v$ moves along a great circle from $p_{\{j\}}$ to $p_{\overline{\{j\}}}$, hence crosses the divisors D_i positively for $i \neq j$, and the divisor D_j once negatively. Therefore, we have $[\boldsymbol{u}] \cdot \boldsymbol{D} = y_j$. See Figure 3(b) for the picture in the one-dimensional case.

This completes the proof of Proposition 6.1.2.

Corollary 6.5.6. *Let us denote*

$$\mathcal{A} := CF^*_{\mathcal{F}(\phi)}(L^n, L^n),$$

where $\mathcal{F}(\phi)$ is the category of Proposition 5.2.2 and $\phi : (M^n, \boldsymbol{D}) \to (M, \boldsymbol{D})$ is the branched cover of Example 3.5.10. Then $\mathcal{A}$ satisfies all of the conditions required to be of type A, in the sense of Definition 2.5.2, except it may not be strictly H-equivariant.

Remark 6.5.7. If one is willing to accept that $CF^*_{\mathcal{F}(M,\boldsymbol{D},\boldsymbol{a})}(L^n, L^n)$ can be defined to all orders, despite L^n being immersed (compare Remark 6.1.1), then we could substitute it for $CF^*_{\mathcal{F}(\phi)}(L^n, L^n)$ in the statement of Corollary 6.5.6. We prefer to work with $\mathcal{F}(\phi)$ because it does not require us to assume a general statement of Gromov compactness for immersed Lagrangians.

Proof. It follows from Proposition 6.1.2 and Proposition 5.2.2 that there is a quasi-isomorphism

$$\mathcal{A}/\mathfrak{m} \cong CF^*_{\mathcal{F}'/\mathfrak{m}}(L', L'),$$

because

$$\mathcal{F}(M, \boldsymbol{D}, \boldsymbol{a})/\mathfrak{m} \cong \mathcal{F}(M, \boldsymbol{D})/\mathfrak{m}.$$

It follows that the underlying $\boldsymbol{G}$-graded vector space of $\mathcal{A}$ is A, and the product μ^2_0 is the exterior product. Furthermore, if $\mathcal{A} = (A, \mu^*)$, it follows that

$$\Phi(\mu^*) = u_1 \dots u_n + \mathfrak{m}$$

(see Definition 2.5.1).

Now let $\mathcal{A} := \mathcal{A}/\mathfrak{m}$. We recall from Lemma 2.5.8 that the spectral sequence induced by the length filtration on $CC^*(\mathcal{A})$ has E_2 page

$$E_2^{*,*} \cong HH^*_c(A),$$

and converges to $HH^*(\mathcal{A})$. It follows from Proposition 6.1.2 that the first-order deformation class of

$$CF^*_{\mathcal{F}(M,\boldsymbol{D})}(L^n, L^n)$$

is given by

$$[\mu_1] = \sum_{j=1}^{n} r_j u_j + (\text{higher-order in length filtration}).$$

It follows from Theorem 4, and the fact that the spectral sequence induced by the length filtration respects the multiplication given by the Yoneda product, that the first-order deformation class of

$$CF^*_{\mathcal{F}(M,\boldsymbol{D},\boldsymbol{a})}(L^n, L^n)$$

is given by

$$[\mu_{1,\boldsymbol{a}}] = \sum_{j=1}^{n} r_j u_j^n + (\text{higher-order in length filtration}).$$

It follows that

$$\Phi(\mu^*) = u_1 \dots u_n + \sum_{j=1}^{n} r_j u_j^n + \mathfrak{m}^2,$$

from which the result follows, by Proposition 5.2.2. ☐

We now have to deal with the fact that $\mathcal{A}$ may not be strictly H-equivariant. It is clear that H acts on $(M, \boldsymbol{D})$, preserving the anchored Lagrangian brane L^n, because all of our constructions of $(M, \boldsymbol{D})$ and L^n have been symmetric with respect to permuting the coordinates. However, it may not be possible to choose our perturbation data H-equivariantly and still achieve transversality, so we may only have H-equivariance 'up to homotopy'. We can fix this using the arguments of Appendix A, which says that we can replace $\mathcal{A}$ by a quasi-equivalent algebra which is strictly H-equivariant: essentially, we just apply the proof of Theorem 3 to this strictly H-equivariant replacement.

Corollary 6.5.8. *Suppose that $\mathcal{B}$ is a $\boldsymbol{G}$-graded A_∞ algebra over R of type A. Then there exists $\psi \in \mathrm{Aut}(R)$, and an A_∞ quasi-isomorphism*

$$\mathcal{B} \cong \psi \cdot \mathcal{A}.$$

Proof. First, observe that Corollary 2.5.7 does not require strict H-equivariance, so there is a quasi-isomorphism

$$\mathcal{A}/\mathfrak{m} \cong \mathcal{B}/\mathfrak{m} =: \mathcal{A}.$$

Now, by a version of Proposition A.0.6 the subcategory of $\mathcal{F}(\phi)$ with object L^n embeds, H-equivariantly, into a strictly H-equivariant A_∞ category, in such a way that the order-0 component of the embedding is a quasi-equivalence. Now recall that $\mathcal{A}$ is necessarily minimal (by Lemma 2.5.9), and it follows easily from the proof of Proposition A.0.6 that this strictly H-equivariant A_∞ category can be chosen to be minimal too. Any A_∞ functor between minimal A_∞ categories, whose order-0 component is a quasi-equivalence, is necessarily a quasi-equivalence. Using the fact that quasi-equivalences of minimal A_∞ categories can be inverted (Lemma 2.3.34), we can apply [2, Lemma 4.3] to prove that $\mathcal{A}$ is quasi-equivalent to a strictly H-equivariant A_∞ algebra of type A. The result now follows from Theorem 3. ☐

Definition 6.5.9. We denote

$$\widetilde{\mathcal{A}} \subset \mathcal{F}(M^n, \boldsymbol{D})$$

the full $\boldsymbol{G}_n^n$-graded subcategory whose objects are the lifts of L^n.

Corollary 6.5.10. *If $\mathcal{B}$ is any A_∞ algebra of type A, then there exists $\psi[[T]] \in \mathrm{Aut}(R)$ and an A_∞ quasi-isomorphism of $\boldsymbol{G}_n^n$-graded R-linear A_∞ categories,*

$$\widetilde{\mathcal{A}} \cong \psi \cdot \boldsymbol{p}_1^* \underline{\mathcal{B}}.$$

Proof. Consider the branched cover

$$\phi : (M_n^n, \boldsymbol{D}) \to (M_1^n, \boldsymbol{D})$$

of Example 3.5.10. By Proposition 5.2.2, there is a fully faithful embedding

$$\boldsymbol{p}_1^* \mathcal{F}(\phi) \to \mathcal{F}(M^n, \boldsymbol{D}),$$

and in particular we have a quasi-equivalence

$$\boldsymbol{p}_1^* \underline{\mathcal{A}} \cong \widetilde{\mathcal{A}}.$$

The result now follows from Corollary 6.5.8. □

7. The B-model

The aim of this section is to prove Theorem 2.

7.1. Homological perturbation lemma.

We will use a version of the homological perturbation lemma which is not quite the usual one (for which see, for example, [37], [38]), but rather the slightly modified version used in [4], so we feel it is as well to state it.

Suppose we are given:

- An A_∞ algebra (B, μ^*) (over a $\mathbb{C}$-algebra R);
- A map $\partial : B \to B$ that is a Maurer-Cartan element for (B, μ^*), in the sense that $\tilde{\mu}^* := (\mu^1 + \partial, \mu^2, \ldots)$ is an A_∞ structure on B;
- A chain complex (C, d_C);
- Chain maps
$$(C, d_C) \underset{p}{\overset{i}{\rightleftarrows}} (B, \mu^1);$$
- A map
$$h : B \to B,$$

such that

- $pi = \mathrm{id}$;
- h defines a homotopy between ip and the identity, which just means that
$$ip = \mathrm{id} - [\mu^1, h];$$
- the **side conditions**
$$\begin{aligned} h^2 &= 0, \\ hi &= 0, \text{ and} \\ ph &= 0. \end{aligned}$$
are satisfied;
- there exists some integer N such that $(\partial h)^N = 0$.

Then we construct:

- An A_∞ structure ν^* on C;
- An A_∞ morphism I^* from $(B, \tilde{\mu}^*)$ to (C, ν^*);
- An A_∞ morphism P^* from (C, ν^*) to $(B, \tilde{\mu}^*)$,

such that I^* and P^* are mutually inverse A_∞ quasi-isomorphisms. In fact, we can show that $P^1 \circ I^1 = \mathrm{id}$, and we can construct an A_∞ homotopy H^* such that

$$I^* \circ P^* = \mathrm{id} - [\tilde{\mu}^*, H^*].$$

This result is proved in the case $\partial = 0$ in [37]. The operations ν^*, I^*, P^*, H^* are defined by certain counts over stable directed planar trees (we use the opposite orientation convention from Definition 4.2.1, so trees have s incoming edges and a single outgoing edge). We attach the operation μ^k to each vertex of arity k (arity = number of incoming edges). If $\partial \neq 0$, then we make exactly the same construction, but sum instead over **semistable** directed planar trees, and attach the operation ∂ to each vertex of arity 1. The assumption that $(\partial h)^N = 0$ ensures that we need only sum over a finite number of trees, because a tree with a sufficiently long chain of vertices of arity 1 does not contribute to the sum.

For example, to define ν^s, we sum over semistable directed planar trees with s incoming edges. We attach operations to each vertex and edge of such a tree, as follows (omitting signs):

- to each vertex of arity 1, attach ∂;
- to each vertex of arity $k \geq 2$, attach μ^k;
- to each internal edge, attach h;
- to each incoming edge, attach i;
- to each outgoing edge, attach p.

Composing the operations as prescribed by the tree determines a map $C^{\otimes s} \to C$. Summing these maps, over all such trees, defines ν^s.

The modifications in the definitions of I^*, P^*, H^*, and the proofs that ν^* is an A_∞ structure, that I^* and P^* are A_∞ morphisms, and that H^* defines an A_∞ homotopy from $I^* \circ P^*$ to id, should all be clear from [37]. The fact that $P^1 \circ I^1 = \mathrm{id}$ follows easily from the side conditions.

7.2. **Matrix factorization computations.** Let k be a field of characteristic 0, and R a commutative k-algebra. Consider the polynomial R-algebra $S := R[x_1, \ldots, x_n]$. Suppose we are given $w \in S$. We consider the differential $\mathbb{Z}_2$-graded category of matrix factorizations $MF(S, w)$. Objects are finitely-generated free $\mathbb{Z}_2$-graded S-modules K, equipped with a 'differential' $\delta_K : K \to K$ of odd degree such that $\delta_K^2 = w \cdot \mathrm{id}$. Morphisms are S-module homomorphisms, with the standard differential and compositions.

By [39, Theorem 3.9], there is an exact equivalence between $\mathrm{Ho}(MF(S,w))$ (where 'Ho' denotes the homotopy category) and Orlov's 'derived category of singularities' $D^b_{\mathrm{Sing}}(w^{-1}(0))$. We consider a matrix factorization (B, δ_B) which corresponds to the ideal

$$\mathcal{O}_0 := (x_1, \ldots x_n)$$

under this equivalence. Following the method described in [40, Section 2.3], we take B to be the free finitely-generated $\mathbb{Z}_2$-graded algebra generated by odd supercommuting variables $\theta_1, \ldots, \theta_n$ (with S in even degree). That is,

$$B := S[\theta_1, \ldots, \theta_n].$$

We define the differential on B to be

$$\delta := \delta_0 + \delta_1,$$

where

$$\begin{aligned} \delta_0 &= \sum_{j=1}^{n} x_j \frac{\partial}{\partial \theta_j}, \\ \delta_1 &= \sum_{j=1}^{n} w_j \theta_j, \end{aligned}$$

where $w_j \in S$ are elements chosen such that

$$w = \sum_{j=1}^{n} w_j x_j.$$

Observe that

$$\begin{aligned} \delta_0^2 &= 0, \\ \delta_1^2 &= 0, \text{ and} \\ [\delta_0, \delta_1] &= w. \end{aligned}$$

It follows that $\delta^2 = w \cdot \mathrm{id}$ as required.

Now consider the differential $\mathbb{Z}_2$-graded algebra

$$\mathcal{B} := \mathrm{Hom}^*_{MF(S,w)}(B, B).$$

Again following [40, Section 2.3], we take the underlying vector space to be the algebra of differential operators

$$\mathcal{B} := S\left[\theta_1, \ldots, \theta_n, \frac{\partial}{\partial \theta_1}, \ldots, \frac{\partial}{\partial \theta_n}\right],$$

acting on B in the obvious way, with the natural multiplication and the differential $d := d_0 + d_1$, where $d_j := [\delta_j, -]$. $\mathcal{B}$ is freely generated, as an R-module, by generators $x^{\boldsymbol{b}} \theta^J \partial^K$, where $\boldsymbol{b} = (b_1, \ldots, b_n)$ is a multi-index, $J \subset [n]$, $K \subset [n]$. We will use the shorthand ∂_j for $\partial / \partial \theta_j$.

Now we use the homological perturbation lemma to construct a minimal A_∞ model for $\mathcal{B}$.

To put ourselves in the situation of Section 7.1, let us consider $\mathcal{B}$ with the A_∞ (in fact, differential graded) structure given by the differential $\mu^1 = d_0$ and standard multiplication μ^2, and let $\partial := d_1$. Then $(\mathcal{B}, \tilde{\mu}^*) = \mathcal{B}$. Furthermore, let

$$\begin{aligned} C &:= R[\partial_1, \ldots, \partial_n], \\ d_C &= 0, \\ i : (C, d_C) &\rightarrow (\mathcal{B}, d_0) \text{ the obvious inclusion,} \\ p : (\mathcal{B}, d_0) &\rightarrow (C, d_C) \text{ the projection defined by} \\ p(x^{\boldsymbol{b}} \theta^J \partial^K) &= \begin{cases} \partial^K & \text{if } \boldsymbol{b} = 0 \text{ and } J = \phi, \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

so that $pi = \mathrm{id}$. We define

$$\begin{aligned} \tilde{h} : B &\to B, \\ \tilde{h}(f\theta^J\partial^K) &:= \left(\sum_{j=1}^{n} \frac{\partial f}{\partial x_j}\theta_j\right)\theta^J\partial^K. \end{aligned}$$

One can check that

$$[d_0, \tilde{h}]\left(x^{\boldsymbol{b}}\theta^J\partial^K\right) = (|\boldsymbol{b}| + |J|)\left(x^{\boldsymbol{b}}\theta^J\partial^K\right).$$

Therefore, if we define

$$h\left(x^{\boldsymbol{b}}\theta^J\partial^K\right) := \begin{cases} 0 & \text{if } \boldsymbol{b} = 0 \text{ and } J = \phi, \\ \frac{1}{|\boldsymbol{b}|+|J|}\tilde{h}\left(x^{\boldsymbol{b}}\theta^J\partial^K\right) & \text{otherwise,} \end{cases}$$

then we have

$$ip = \mathrm{id} - [d_0, h].$$

Furthermore, we can check that the side conditions are satisfied:

$$\begin{aligned} h^2 &= 0 \text{ (for the same reason the exterior derivative squares to 0)}, \\ hi &= 0, \text{ and} \\ ph &= 0. \end{aligned}$$

Finally, observe that $h\partial$ decreases the grading $|x^{\boldsymbol{b}}\theta^J\partial^K| := |K|$ by 1, so $(h\partial)^{n+1} = 0$.

Thus, we can apply Section 7.1 to construct an A_∞ structure ν^* on C, which is quasi-isomorphic to $\mathcal{B}$. If w has degree ≥ 3, then the differential $\nu^1 = 0$ and the product ν^2 is the standard (exterior algebra) product on C. Thus, $\nu^{\geq 3}$ defines a Maurer-Cartan element in the Hochschild cochain complex $CC^*(C)$.

Now we recall, from Definition 2.5.1, the Hochschild-Kostant-Rosenberg map from the Hochschild cohain complex to the space of polyvector fields,

$$\Phi : CC^*(C) \to S[\partial_1, \ldots, \partial_n],$$

given by

$$\Phi(\alpha) := \sum_{s \geq 0} \alpha^s(\boldsymbol{x}, \ldots, \boldsymbol{x}),$$

where we denote

$$\boldsymbol{x} := \sum_{j=1}^{n} x_j \partial_j.$$

Proposition 7.2.1. *(see* [24, Proposition 7.1]*) The image of the Maurer-Cartan element $\nu^{\geq 3}$ under Φ is exactly the superpotential w.*

Proof. We recall the construction of the maps ν^k from Section 7.1, by summing over trees. The only trees that give a non-zero contribution to a product $\nu^k(\partial_{i_1}, \ldots, \partial_{i_k})$ are those depicted in Figure 7.2.1.

Given such a tree, with inputs $\partial_{i_1}, \ldots, \partial_{i_k}$, one easily determines the output: it is the constant term of

$$\partial_{i_k} \ldots \partial_{i_2} w_{i_1},$$



FIGURE 7.2.1. The only trees contributing to the homological perturbation lemma computation.

divided by $(k-1)!$ (coming from the terms in the denominator of h). Given a monomial $x^{\boldsymbol{b}}$ in w_j, there are $|\boldsymbol{b}|!/\boldsymbol{b}!$ ways of choosing the order of the inputs ∂_{i_m} we take from the term

$$\nu^{|\boldsymbol{b}|+1}(\boldsymbol{x}, \ldots, \boldsymbol{x})$$

of $\Phi(\nu^{\geq 3})$, in such a way that $j = i_1$ comes first, and the constant term of

$$\partial_{i_k} \ldots \partial_{i_2} w_j$$

is non-zero. Thus, each monomial $x^{\boldsymbol{b}}$ of w_j contributes a term

$$x_j \frac{1}{|\boldsymbol{b}|!} \frac{|\boldsymbol{b}|!}{\boldsymbol{b}!} \left(\partial^{\boldsymbol{b}} x^{\boldsymbol{b}}\right) x^{\boldsymbol{b}} = x_j x^{\boldsymbol{b}}$$

to $\Phi(\nu^{\geq 3})$, and the result follows. □

7.3. $\boldsymbol{G}$-Graded matrix factorizations.

For the purposes of this section, let $\boldsymbol{G}$ be a grading datum:

$$\mathbb{Z} \xrightarrow{f} Y \to X \to 0,$$

with sign morphism $\boldsymbol{\sigma}$.

Let S be a $\boldsymbol{G}$-graded algebra such that $\boldsymbol{\sigma}_* S$ is concentrated in degree $0 \in \mathbb{Z}_2$, and let $w \in S$ be an element of degree $f(2) \in Y$.

Definition 7.3.1. A ***G*-graded matrix factorization** of $w \in S$ is a $\boldsymbol{G}$-graded finitely-generated free S-module K, together with a homomorphism

$$\delta_K \in \mathrm{Hom}_S(K,K)$$

of degree $f(1) \in Y$, such that

$$\delta_K^2 = w \cdot \mathrm{id}.$$

Definition 7.3.2. We define the **differential *G*-graded category of matrix factorizations**, $MF^{\boldsymbol{G}}(S,w)$:

- Objects are $\boldsymbol{G}$-graded matrix factorizations of w;
- Morphisms are S-module homomorphisms:
$$\mathrm{Hom}((K,\delta_K),(L,\delta_L)) := \mathrm{Hom}_S(K,L);$$
- Differential on morphism spaces is as usual:
$$\partial(F) := \delta_L \circ F - (-1)^{\sigma(F)} F \circ \delta_K;$$
- Composition is composition of S-module homomorphisms.

We note that the morphism spaces are naturally $\boldsymbol{G}$-graded S-modules, the differential and composition maps have degrees $f(1)$ and $f(0) \in Y$ respectively, and they satisfy the Leibniz rule. It follows that $MF^{\boldsymbol{G}}(S,w)$ is a $\boldsymbol{G}$-graded A_∞ category over S (see Remark 2.3.12). In fact it is a differential $\boldsymbol{G}$-graded category, since all $\mu^{\ge 3}$ are zero.

We observe that ordinary matrix factorizations are nothing more than $\boldsymbol{G}_\sigma$-graded matrix factorizations. It follows that there is a fully faithful embedding

$$\boldsymbol{\sigma}_* MF^{\boldsymbol{G}}(S,w) \to MF(S,w).$$

Now let us introduce our main example. We will introduce graded matrix factorizations mirror to the smooth orbifold relative Fukaya category (compare Section 6.1). Let $\boldsymbol{G} := \boldsymbol{G}_1^n$ be the grading datum introduced in Example 2.1.13. Let

$$R := R_n^n \cong \mathbb{C}[[r_1,\ldots,r_n]]$$

be the $\boldsymbol{G}$-graded power series ring introduced in Definition 2.2.13.

Let U be the $\boldsymbol{G}$-graded vector space of Example 2.2.9. We consider the $\boldsymbol{G}$-graded algebra

$$S := R[U] \cong \mathbb{C}[[r_1,\ldots,r_n]][u_1,\ldots,u_n].$$

We consider the element

$$w = u_1 \ldots u_n + \sum_{j=1}^n r_j u_j^n \in S,$$

which has degree $f(2) \in Y$, hence we can define the $\boldsymbol{G}$-graded category of matrix factorizations $MF^{\boldsymbol{G}}(S,w)$.

We consider the $\boldsymbol{G}$-graded S-module

$$K := R[U] \otimes \Lambda\left(U^\vee\right) \cong \mathbb{C}[[r_1,\ldots,r_n]][u_1,\ldots,u_n][\theta_1,\ldots,\theta_n],$$

where the variables θ_j anti-commute. We introduce the differential

$$\begin{aligned} \delta_K : K &\rightarrow K \\ \delta_K &= \sum_{j=1}^{n} u_j \frac{\partial}{\partial \theta_j} + w_j \theta_j, \end{aligned}$$

where

$$w_j = \frac{u_1 \dots u_n}{n u_j} + r_j u_j^{n-1}.$$

We observe that δ_K has degree $f(1) \in Y$, and that

$$\delta_K^2 = w \cdot \mathrm{id}.$$

Thus, (K, δ_K) is a $\boldsymbol{G}$-graded matrix factorization of w. We denote it by $\mathcal{O}_0$. Finally, we observe that δ_K is H-invariant, where H is the symmetric group acting in the obvious way.

Corollary 7.3.3. *Let us define the $\boldsymbol{G}$-graded A_∞ algebra over R:*

$$\mathcal{B} := \mathrm{Hom}_{MF^{\boldsymbol{G}}(S,w)}(\mathcal{O}_0, \mathcal{O}_0).$$

Then $\mathcal{B}$ is of type A, in the sense of Definition 2.5.2.

Proof. Follows from Proposition 7.2.1. We observe that it is necessary to check that the $\boldsymbol{G}$-grading and H-equivariance interact appropriately with the homological perturbation lemma construction, but this is clear. □

Corollary 7.3.4. *Let*

$$\mathcal{A} := CF^*_{\mathcal{F}(\phi)}(L^n, L^n),$$

where $\mathcal{F}(\phi)$ is the category of Proposition 5.2.2 and $\phi : (M^n, \boldsymbol{D}) \rightarrow (M, \boldsymbol{D})$ is the branched cover of Example 3.5.10. Let

$$\mathcal{B} := \mathrm{Hom}_{MF^{\boldsymbol{G}}(S,w)}(\mathcal{O}_0, \mathcal{O}_0)$$

as above. Both $\mathcal{A}$ and $\mathcal{B}$ are $\boldsymbol{G}$-graded A_∞ algebras over R, and there exists a power series $\psi \in \mathbb{C}[[T]]$, with $\psi(0) = 1$ (recalling $T = r_1 \dots r_n$), and an A_∞ quasi-isomorphism

$$\mathcal{A} \cong \psi \cdot \mathcal{B}.$$

Proof. Follows from Corollary 7.3.3 and Corollary 6.5.8. □

7.4. **Equivariant matrix factorizations.**

Suppose that $\boldsymbol{p} : \boldsymbol{G}' \rightarrow \boldsymbol{G}$ is an injective morphism of grading data. Then we consider the $\boldsymbol{G}'$-graded category

$$\boldsymbol{p}^* MF^{\boldsymbol{G}}(S, w).$$

Remark 7.4.1. Objects of $\boldsymbol{p}^* MF^{\boldsymbol{G}}(S, w)$ are again $\boldsymbol{G}$-graded matrix factorizations, but the morphism spaces are just the parts whose Y-grading lies in the image of $p : Y' \rightarrow Y$. Thus $\boldsymbol{p}^* MF^{\boldsymbol{G}}(S, w)$ embeds, fully faithfully, in the category of $\mathrm{coker}(p)^*$-equivariant matrix factorizations.

For example, let $\boldsymbol{p}_1 : \boldsymbol{G}^n_n \rightarrow \boldsymbol{G}^n_1$ be the morphism defined in Lemma 2.1.14.

Definition 7.4.2. We denote

$$\widetilde{\mathcal{B}} := \boldsymbol{p}_1^* \underline{\mathcal{B}} \subset \boldsymbol{p}_1^* MF^{\boldsymbol{G}}(S,w).$$

It is the full subcategory whose objects are $\mathcal{O}_0$ and its shifts by elements $y \in Y$.

We obtain the following:

Corollary 7.4.3. *There exists a power series* $\psi \in \mathbb{C}[[T]]$, $\psi(0) = 1$, *and an* A_∞ *quasi-isomorphism of* $\boldsymbol{G}_n^n$*-graded* R*-linear* A_∞ *categories,*

$$\widetilde{\mathcal{A}} \cong \psi \cdot \widetilde{\mathcal{B}}.$$

Proof. Follows from Corollary 7.3.3 and Corollary 6.5.10. ☐

7.5. **Coherent sheaves.** In this section, we will explain how to relate equivariant categories of coherent sheaves on a projective variety to equivariant categories of graded matrix factorizations. We state a result closely related to [22, Proposition 1.2.2] (see also [23, Section 2]), in the language of $\boldsymbol{G}$-graded matrix factorizations. First, we recall Orlov's **category of graded matrix factorizations** (see [21, Section 3.1]).

Let S be a $\mathbb{Z}$-graded ring, and $w \in S$ homogeneous of degree d. Recall from Example 2.1.11 the grading datum $\boldsymbol{G}_{MF(d)}$. We define

$$\begin{aligned} q : \mathbb{Z} &\to \mathbb{Z} \oplus \mathbb{Z}/(2,-d), \\ q(j) &= (0,j), \end{aligned}$$

so we can equip $S \cong q_* S$ with a $\boldsymbol{G}_{MF(d)}$-grading. Then w has degree $(0,d) \sim (2,0) = f(2) \in Y$, so we can define the category of $\boldsymbol{G}_{MF(d)}$-graded matrix factorizations. Now there is a unique injective morphism of grading data, $\boldsymbol{p} : \boldsymbol{G}_{\mathbb{Z}} \to \boldsymbol{G}_{MF(d)}$. We define

$$\mathrm{GrMF}(S,w) := \boldsymbol{p}^* MF^{\boldsymbol{G}_{MF(d)}}(S,w).$$

It is not hard to see that this definition coincides with Orlov's category of graded matrix factorizations of w (actually Orlov defines graded matrix factorizations to be the homotopy category of this differential graded category). Namely, if we denote

$$\pi : \mathbb{Z} \oplus \mathbb{Z} \to \mathbb{Z} \oplus \mathbb{Z}/(2,-d),$$

then given a $\boldsymbol{G}_{MF(d)}$-graded S-module K, $\pi^* K$ is a quasi-periodic complex of graded S-modules, as in the usual definition of the category of graded matrix factorizations.

We recall the relationship of GrMF to coherent sheaves. Suppose that k is a field, $S = k[u_1, \ldots, u_n]$ is the $\mathbb{Z}$-graded polynomial ring, and $w \in S$ is homogeneous of degree n. Suppose that the variety

$$X := \{w = 0\} \subset \mathbb{P}^{n-1}_k$$

is smooth. Then, because X is Calabi-Yau, [21, Theorem 3.11] says that there is an equivalence of triangulated categories,

$$D^b Coh\,(X) \cong \mathrm{Ho}(\mathrm{GrMF}(S,w)).$$

Remark 7.5.1. We observe that the matrix factorization $\mathcal{O}_0$ is always $\boldsymbol{G}_{MF(n)}$-graded. If $\mathcal{O}_0[j]$ denotes the shift of $\mathcal{O}_0$ by the integer j (i.e., by $f(j)$), then we can arrange that the object

$$\bigoplus_{j=0}^{n-1} \mathcal{O}_0[j] \in Ob(\mathrm{GrMF}(S,w))$$

corresponds, under Orlov's equivalence, to the restriction of the Beilinson exceptional collection

$$\bigoplus_{j=0}^{n-1} i^*\Omega^j(j),$$

where $i : X \hookrightarrow \mathbb{P}^{n-1}_k$ denotes the inclusion. See [41, Remark 5.20], also [42, Section IV.A].

Now let us consider the situation of Section 7.4. Recall the commutative square of grading data of Lemma 2.1.14. We have

$$R := \mathbb{C}[[r_1, \ldots, r_n]]$$

and

$$S := R[u_1, \ldots, u_n]$$

are $\boldsymbol{G}_1^n$-graded rings, and $w \in S$ has degree $f(2) \in Y$. Recall that the Novikov field Λ is an R-algebra, via the map

$$\begin{array}{rcl} R & \to & \Lambda, \\ r_j & \mapsto & r. \end{array}$$

We define

$$S_{nov} := S \otimes_R \Lambda \cong \Lambda[u_1, \ldots, u_n],$$

with the object

$$w_{nov} := w \otimes 1 \in S_{nov}.$$

Equip S_{nov} with the standard $\mathbb{Z}$-grading (where Λ is concentrated in degree 0, and each u_j has degree 1). Consider the variety

$$\widetilde{N}^n_{nov} := \{w_{nov} = 0\} \subset \mathrm{Proj}(S_{nov}) \cong \mathbb{P}^{n-1}_{\Lambda}.$$

Define the group Γ_n to be the kernel of the map

$$(\mathbb{Z}_n)^n / y_{[n]} \to \mathbb{Z}_n,$$

and equip $\widetilde{N}^n_{nov}$ with the action of the character group Γ_n^*, acting by multiplying the coordinates u_j by nth roots of unity. We denote

$$N^n_{nov} := \widetilde{N}^n_{nov} / \Gamma^*_n.$$

We observe that these definitions coincide with those given in the Introduction.

Lemma 7.5.2. *There is a fully faithful embedding of triangulated categories,*

$$\mathrm{Ho}\left(\boldsymbol{q}_{1*}\boldsymbol{p}_1^* MF^{\boldsymbol{G}}(S,w)\right) \otimes_R \Lambda \to D^b Coh\left(N^n_{nov}\right).$$

Proof. First, observe that $\boldsymbol{q}_{2*}S$ is a $\boldsymbol{G}_{MF(n)}$-graded ring. Furthermore, it is easy to check that $\boldsymbol{q}_{2*}R$ is concentrated in degree 0, and u_i has degree $(0,1)$. This coincides with the $\boldsymbol{G}_{MF(n)}$-grading of S induced by the standard $\mathbb{Z}$-grading.

There is a fully faithful embedding

$$\boldsymbol{q}_{2*}MF^{\boldsymbol{G}_1^n}(S,w) \to MF^{\boldsymbol{G}_{MF(n)}}(\boldsymbol{q}_{2*}S,w),$$

which sends

$$(K,\delta_K) \mapsto (\boldsymbol{q}_{2*}K,\delta_K).$$

Therefore, there is a fully faithful embedding

$$\boldsymbol{p}_2^*\boldsymbol{q}_{2*}MF^{\boldsymbol{G}}(S,w) \to \mathrm{GrMF}(S,w).$$

Now note that $\boldsymbol{p}_2^*\boldsymbol{q}_{2*}R \cong R$ is concentrated in degree $0 \in \mathbb{Z}$, so the morphism $R \to \Lambda$ respects the $\mathbb{Z}$-grading, and we obtain a fully faithful embedding of Λ-linear differential graded categories

$$\left(\boldsymbol{p}_2^*\boldsymbol{q}_{2*}MF^{\boldsymbol{G}}(S,w)\right) \otimes_R \Lambda \to \mathrm{GrMF}(S,w) \otimes_R \Lambda.$$

There is a fully faithful embedding of Λ-linear differential graded categories

$$\mathrm{GrMF}(S,w) \otimes_R \Lambda \to \mathrm{GrMF}(S_{nov},w_{nov}),$$

which sends

$$K \mapsto K \otimes_R \Lambda$$

on the level of objects (recall that K is by definition a **free** S-module). By Orlov's theorem [21, Theorem 3.11], there is an equivalence of Λ-linear triangulated categories,

$$\mathrm{Ho}\left(\mathrm{GrMF}(S_{nov},w_{nov})\right) \cong D^bCoh\left(\widetilde{N}^n_{nov}\right).$$

It follows that there is an equivalence of triangulated categories

$$\mathrm{Ho}\left(\mathrm{GrMF}(S_{nov},w_{nov})^{\Gamma_n^*}\right) \cong D^bCoh^{\Gamma_n^*}\left(\widetilde{N}^n_{nov}\right).$$

Hence, by the argument above, there is a fully faithful embedding

$$\mathrm{Ho}\left(\boldsymbol{p}_2^*\boldsymbol{q}_{2*}MF^{\boldsymbol{G}}(S,w)^{\Gamma_n^*} \otimes_R \Lambda\right) \to D^bCoh^{\Gamma_n^*}\left(\widetilde{N}^n_{nov}\right) \cong D^bCoh(N^n_{nov}).$$

Now we recall Lemma 2.2.14. It shows that there is an isomorphism

$$\boldsymbol{p}_2^*\boldsymbol{q}_{2*}MF^{\boldsymbol{G}}(S,w)^{\Gamma_n^*} \cong \boldsymbol{q}_{1*}\boldsymbol{p}_1^*MF^{\boldsymbol{G}}(S,w),$$

where

$$\Gamma_n := \ker(q_{2,X})/\mathrm{im}(p_{1,X}).$$

In this case, an examination of Lemma 2.1.14 shows that Γ_n is indeed the kernel of the map

$$(\mathbb{Z}_n)^n/y_{[n]} \to \mathbb{Z}_n$$

given by summing the coordinates, and the Γ_n-gradings of morphism spaces correspond under Orlov's equivalence. The result follows. □

Corollary 7.5.3. *There is an equivalence of* Λ*-linear triangulated categories,*

$$\left(\boldsymbol{q}_{1*}\widetilde{\mathcal{B}}\right) \otimes_R \Lambda \cong D^bCoh\left(N^n_{nov}\right).$$

Proof. It follows from Lemma 7.5.2 that there is a fully faithful embedding of the left-hand side into the right-hand side. In fact, the embedding is essentially surjective. This follows from Remark 7.5.1, together with (the Γ_n^*-equivariant version of) [2, Lemma 5.4], together with (the Γ_n^*-equivariant version of) Beilinson's generation result [43]. □

Definition 7.5.4. We denote

$$\widetilde{\mathcal{A}}_{nov} := \boldsymbol{q}_{1*}\widetilde{\mathcal{A}} \otimes_R \Lambda.$$

It is a full subcategory of $\boldsymbol{q}_{1*}\mathcal{F}(M^n, \boldsymbol{D}) \otimes_R \Lambda$.

Corollary 7.5.5. *There exists $\psi \in \mathbb{C}[[r^n]]$, and an equivalence of Λ-linear triangulated categories,*

$$\psi \cdot D^b Coh(N^n_{nov}) \cong \mathrm{Ho}\left(Tw\left(\widetilde{\mathcal{A}}_{nov}\right)\right),$$

where 'Tw' denotes forming the category of twisted complexes.

Proof. Follows from Corollary 7.5.3 and Corollary 7.4.3. □

8. The full Fukaya category

In this Section, we consider the full Fukaya category $\mathcal{F}(M^n)$. We explain why there is an embedding

$$\mathcal{F}(M^n, \boldsymbol{D}) \otimes_R \Lambda \to \mathcal{F}(M^n),$$

and prove that the full subcategory

$$\widetilde{\mathcal{A}} \otimes_R \Lambda \subset \mathcal{F}(M^n)$$

(see Definition 6.5.9) split-generates, using the criterion of [5]. This allows us to complete the proof of Theorem 1.

8.1. Relating the full and relative Fukaya categories.

Let (M, ω) be a compact symplectic manifold, satisfying $c_1(M) = 0$. The Fukaya category $\mathcal{F}(M)$ is defined in [5]. It depends on the choice of a bulk class $\mathfrak{b} \in H^{even}(M; \Lambda_0)$, and a background class $st \in H^2(M; \mathbb{Z}_2)$. We choose both of these to be 0. The objects of $\mathcal{F}(M)$ are (L, b), where $L \subset M$ is a graded spin Lagrangian submanifold, and $b \in H^1(L; \Lambda_0)$ is a weak bounding cochain. Composition maps are defined by counting holomorphic disks. $\mathcal{F}(M)$ is a $\mathbb{Z}$-graded A_∞ category.

We would like to relate $\mathcal{F}(M, \boldsymbol{D})$ to $\mathcal{F}(M)$. First, we observe that there is a canonical morphism of grading data,

$$\boldsymbol{q} : \boldsymbol{G}(M, \boldsymbol{D}) \to \boldsymbol{G}_{\mathbb{Z}},$$

by Remark 3.5.7. Thus we can define a $\boldsymbol{G}_{\mathbb{Z}}$-graded (i.e., $\mathbb{Z}$-graded) A_∞ category

$$\boldsymbol{q}_*\mathcal{F}(M, \boldsymbol{D}).$$

$\mathcal{F}(M, \boldsymbol{D})$ is defined over the coefficient ring $R \cong \mathbb{C}[[r_1, \ldots, r_k]]$. It is a simple task to show that the coefficient ring $\boldsymbol{q}_*R$ has degree $0 \in \mathbb{Z}$. Therefore, the ring homomorphism

$$\begin{array}{rcl} R & \to & \Lambda, \\ r_j & \mapsto & r^{d_j}, \end{array}$$

respects the $\mathbb{Z}$-grading, because both have degree $0 \in \mathbb{Z}$. This makes Λ into an R-module, and means that we can define the $\mathbb{Z}$-graded, Λ-linear A_∞ category

$$\boldsymbol{q}_*\mathcal{F}(M, \boldsymbol{D}) \otimes_R \Lambda.$$

We remark that, up until this point, we have given a complete definition of the relative Fukaya category $\mathcal{F}(M, \boldsymbol{D})$ of a Kähler pair (satisfying assumptions as in Definition 3.5.1), using explicit domain-dependent perturbations of the holomorphic curve equation. To define the full Fukaya category $\mathcal{F}(M)$ however, we need virtual perturbations as in [20, 5]. These two categories should be related as follows:

Assumption 8.1.1. There is a fully faithful embedding

$$\boldsymbol{q}_* \mathcal{F}(M, \boldsymbol{D}) \otimes_R \Lambda \to \mathcal{F}(M)$$

of $\mathbb{Z}$-graded, Λ-linear A_∞ categories.

To prove this, we would have to relate the two different perturbation schemes (explicit domain-dependent perturbations versus perturbations of Kuranishi structures), which would take us beyond the scope of this paper. So we leave it as an assumption. We do, however, provide the following justification:

Remark 8.1.2. We observe that there is an obvious map on the level of unobstructed objects (Lagrangians with $\mu^0 = 0$):

$$\begin{array}{rcl} Ob(\boldsymbol{q}_* \mathcal{F}(M, \boldsymbol{D}))_{unob} & \to & Ob(\mathcal{F}(M)), \\ L & \mapsto & (L, 0), \end{array}$$

and anchored Lagrangian branes automatically come with a grading (recall that taking $\boldsymbol{q}_*$ of a category involves identifying certain objects; in this case, this exactly means that we identify all anchored Lagrangian branes which have the same grading). However, there is no map in the other direction: objects of $\mathcal{F}(M)$ may intersect the divisors $\boldsymbol{D}$. Suppose now that:

- $\boldsymbol{L}$ is a tuple of exact, transversely-intersecting anchored Lagrangian branes in $M \setminus \boldsymbol{D}$;
- $\boldsymbol{p}$ is an associated set of generators (intersection points) of $\boldsymbol{L}$;
- the moduli space of rigid, boundary-punctured holomorphic disks in M with boundary on $\boldsymbol{L}$, asymptotic to $\boldsymbol{p}$, is regular.

We show that a rigid holomorphic disk in this moduli space contributes the same term to an A_∞ structure map μ^s in $\mathcal{F}(M, \boldsymbol{D}) \otimes_R \Lambda$ and in $\mathcal{F}(M)$. We have

$$\omega(u) = -\alpha(p_0) + \sum_{j=1}^{s} \alpha(p_j) + |u \cdot \boldsymbol{D}|$$

by Stokes' theorem, where $\alpha(p)$ denotes the symplectic action of generator p. In particular, if we define the map

$$\begin{array}{rcl} CF^*_{\mathcal{F}(M,\boldsymbol{D}) \otimes_R \Lambda}(L_0, L_1) & \to & CF^*_{\mathcal{F}(M)}(L_0, L_1), \\ p & \mapsto & r^{\alpha(p)} p, \end{array}$$

then the holomorphic disk u contributes the same term $\pm r^{\omega(u)}$ to μ^s in both categories. Note that, in $\mathcal{F}(M, \boldsymbol{D})$, we have $(u \cdot \boldsymbol{D})!$ choices for the labelling of the marked points mapping to the divisors $\boldsymbol{D}$, so this disk in fact contributes $(u \cdot \boldsymbol{D})!$ identical terms to μ^s, each of which is $\pm r^{\omega(u)}/(u \cdot \boldsymbol{D})!$ (see the definition in Section 5.1). Thus its total contribution is exactly $\pm r^{\omega(u)}$.

8.2. **Split-generation.** We recall the subcategory

$$\widetilde{\mathcal{A}}_{nov} \subset \boldsymbol{q}_{1*}\mathcal{F}(M^n, \boldsymbol{D}) \otimes_R \Lambda$$

from Definition 7.5.4. By abuse of notation, we will identify this category with its image under the embedding of Assumption 8.1.1. Our aim in this section is to prove that $\widetilde{\mathcal{A}}_{nov}$ split-generates $D^\pi \mathcal{F}(M^n)$.

Definition 8.2.1. We define the **closed-open string map** from the (small) quantum cohomology ring of M (with coefficients in Λ) to the Hochschild cohomology of $\mathcal{F}(M)$,

$$\mathcal{CO} : QH^*(M) \to HH^*(\mathcal{F}(M)),$$

as follows: let $\alpha \in H^j(M;\mathbb{C})$ be Poincaré dual to a smooth cycle $A \subset M$. Let $\boldsymbol{L}$ be a tuple of objects with associated generators $\boldsymbol{p}$. We consider the moduli space $\mathcal{M}_4(\boldsymbol{p}, A)$, whose objects consist of pairs (r, u), where $r \in \mathcal{R}_1(\boldsymbol{L})$ and $u : S_r \to M$ is a smooth map, such that

- u satisfies the (perturbed) holomorphic curve equation, with Lagrangian boundary conditions given by the labels $\boldsymbol{L}$;
- u is asymptotic to the generators $\boldsymbol{p}$ at the boundary punctures;
- $u(q) \in A$, where $q \in S_r$ is the internal marked point.

Then each rigid disk $u \in \mathcal{M}_4(\boldsymbol{p}, A)$ contributes a term $\pm r^{\omega(u)} p_0$ to

$$\mathcal{CO}(\alpha)(p_s, \ldots, p_1).$$

We remark that

- When we say 'perturbed' holomorphic curve equation, it really means we must define a Kuranishi structure on $\mathcal{M}_4(\boldsymbol{p}, A)$ and introduce virtual perturbations thereof (see [5]);
- The map $\mathcal{CO}$ is a homomorphism of $\mathbb{Z}$-graded Λ-algebras, where the product on $QH^*(M)$ is quantum cup product $*$, and the product on $HH^*(\mathcal{F}(M))$ is the Yoneda product, and the $\mathbb{Z}$-gradings are the standard ones.

We now aim to apply the following result, which is due to [5]:

Theorem 5. *If (M, ω) is a compact $2d$-dimensional Calabi-Yau symplectic manifold, $\mathcal{L}$ a full subcategory of $\mathcal{F}(M)$ with some finite set of objects, and the map*

$$\mathcal{CO}^{2d} : QH^{2d}(M) \to HH^{2d}(\mathcal{L})$$

is non-zero, then $\mathcal{L}$ split-generates $\mathcal{F}(M)$.

We consider the subcategory $\widetilde{\mathcal{A}}_{nov} \subset \mathcal{F}(M^n)$ (actually this is not a finite collection of Lagrangians, because we include all shifts, but it will suffice to choose one representative of each geometric lift of L^n to M^n). We aim to understand the map

$$\mathcal{CO}^{2(n-2)} : QH^{2(n-2)}(M^n) \to HH^{2(n-2)}\left(\widetilde{\mathcal{A}}_{nov}\right)$$

by first understanding the degree-2 part of the map, $\mathcal{CO}^2$, then using the fact that $\mathcal{CO}$ is a Λ-algebra homomorphism. It is expected that the image of the class of the symplectic form,

$$\mathcal{CO}([\omega]) \in HH^2(\mathcal{F}(M)),$$

should be the class corresponding to the deformation of $\mathcal{F}(M)$ given by scaling the Novikov parameter r. In fact, in our relative setting, we can make a statement with a cleaner proof:

Lemma 8.2.2. *Let $(M, \boldsymbol{D})$ be a Kähler pair. Consider the full subcategory*

$$\boldsymbol{q}_* \mathcal{F}(M, \boldsymbol{D}) \otimes_R \Lambda \subset \mathcal{F}(M)$$

of Assumption 8.1.1. Then $\mathcal{CO}([\omega])$ is the image of the class

$$\frac{1}{d_j} \left(r_j \frac{\partial \mu^*}{\partial r_j} \right) \otimes 1 \in HH^*(\mathcal{F}(M, \boldsymbol{D})) \otimes_R \Lambda$$

in $HH^*(\mathcal{F}(M, \boldsymbol{D}) \otimes_R \Lambda)$, for any j.

To clarify: $\mu^* \in CC^*(\mathcal{F}(M, \boldsymbol{D}))$ is the A_∞ structure map, and

$$r_j \frac{\partial \mu^*}{\partial r_j} \in CC^*(\mathcal{F}(M, \boldsymbol{D}))$$

is a Hochschild cochain, as can be seen by applying $r_j \partial / \partial r_j$ to the A_∞ associativity equation $\mu^* \circ \mu^* = 0$. Thus, it defines a class in $HH^*(\mathcal{F}(M, \boldsymbol{D})$, and we consider the image of this class under the map

$$HH^*(\mathcal{F}(M, \boldsymbol{D})) \otimes_R \Lambda \to HH^*(\mathcal{F}(M, \boldsymbol{D}) \otimes_R \Lambda).$$

Proof. Given Lagrangian branes $\boldsymbol{L}$ with associated generators $\boldsymbol{p}$, each rigid holomorphic disk u with boundary on $\boldsymbol{L}$, asymptotic to $\boldsymbol{p}$, contributes a term

$$\pm r_1^{u \cdot D_1} \ldots r_k^{u \cdot D_k}$$

to the coefficient of p_0 in $\mu^s(p_s, \ldots, p_1)$, and hence a term

$$\pm (u \cdot D_j) r_1^{u \cdot D_1} \ldots r_k^{u \cdot D_k}$$

to the corresponding coefficient of $r_j \partial \mu^s / \partial r_j$, and hence a term

$$\pm (u \cdot D_j) r^{\omega(u)}$$

to the corresponding coefficient of $(r_j \partial \mu^s / \partial r_j) \otimes_R 1$ (see Remark 8.1.2).

On the other hand, recall that D_j is Poincaré dual to $d_j \omega$ by definition of a Kähler pair. So, by definition of the map $\mathcal{CO}$, each such holomorphic disk u together with an internal marked point q mapping to D_j, contributes a term $\pm r^{\omega(u)}$ to the corresponding coefficient of $\mathcal{CO}(d_j \omega)$. There are $u \cdot D_j$ choices for the internal marked point q, so the total contribution of each such holomorphic disk u is $\pm (u \cdot D_j) r^{\omega(u)}$.

Therefore,

$$\mathcal{CO}(d_j \omega) = \left(r_j \frac{\partial \mu^*}{\partial r_j} \right) \otimes_R 1$$

as required. □

Remark 8.2.3. Again, Lemma 8.2.2 should perhaps be thought of as an assumption, for the same reason that we make Assumption 8.1.1.

Proposition 8.2.4. *Let M^n be the Calabi-Yau Fermat hypersurface of Example 3.5.2, and recall the full subcategory $\widetilde{\mathcal{A}}_{nov} \subset \mathcal{F}(M^n)$. The map*

$$\mathcal{CO}^{2(n-2)} : QH^{2(n-2)}(M^n) \to HH^{2(n-2)}\left(\widetilde{\mathcal{A}}_{nov}\right)$$

is non-zero.

Proof. We recall from Lemma 2.5.18 that there is an action of $\tilde{\Gamma}_n^*$ on $HH^*(\widetilde{\mathcal{A}}_{nov})$, and the $\tilde{\Gamma}_n^*$-invariant part is

$$HH^*_{\boldsymbol{G}_{\mathbb{Z}}}\left(\widetilde{\mathcal{A}}_{nov}\right)^{\tilde{\Gamma}_n^*} \cong \Lambda[\alpha]/\alpha^{n-1}$$

as a $\mathbb{Z}$-graded Λ-algebra, where α has degree 2.

It follows from Lemma 8.2.2 and Lemma 2.5.19 that the image of $\mathcal{CO}([\omega])$ under this isomorphism is $g \cdot \alpha$, for some invertible $g \in \Lambda^*$. Therefore, because $\mathcal{CO}$ is a Λ-algebra homomorphism, the image of $\mathcal{CO}([\omega]^{n-2})$ under this isomorphism is $g^{n-2}\alpha^{n-2}$, which does not vanish in $\Lambda[\alpha]/\alpha^{n-1}$. Because it has degree $2(n-2)$, this completes the proof. □

Corollary 8.2.5. *The full subcategory*

$$\widetilde{\mathcal{A}}_{nov} \subset \mathcal{F}(M^n)$$

split-generates the Fukaya category.

Proof. Follows from Proposition 8.2.4 and Theorem 5. □

Theorem 1 now follows from Corollary 7.5.5, Assumption 8.1.1, and Corollary 8.2.5.

Appendix A. Strict group actions on Fukaya categories

This section is based on the argument of [2, Section 8b]. Suppose that we have a finite group Γ, which acts on a Kähler pair $(M, \boldsymbol{D})$, permuting the divisors $\boldsymbol{D}$, and preserving the Liouville one-form α. We consider the relative Fukaya category $\mathcal{F}(M, \boldsymbol{D})$ defined in Section 5.1. There is an obvious action of Γ on the objects of the relative Fukaya category. We would like to say that this action extends to an action on the relative Fukaya category, in a suitable sense.

Recall the notion of a strictly Γ-equivariant A_∞ structure from Definition 2.4.14. Naïvely, one might try to argue that Γ acts on the moduli spaces of pseudoholomorphic disks used to define the structure maps of the Fukaya category, and hence the structure maps are strictly Γ-equivariant. However, this does not work: for Γ to act on the moduli spaces of pseudoholomorphic disks, we would have to make a Γ-equivariant choice of perturbation data, which would destroy our chances of achieving transversality. Instead, we have the following:

Proposition A.0.6. *In the situation described above, there is a fully faithful embedding of $\mathcal{F}(M, \boldsymbol{D})$ into a strictly Γ-equivariant A_∞ category. The order-0 part of this embedding is a quasi-equivalence, and respects the action of Γ on objects, in the sense that $F(\gamma \cdot L)$ is quasi-isomorphic to $\gamma \cdot F(L)$.*

Proof. We consider a category $\widetilde{\mathcal{F}}(M, \boldsymbol{D})$ with objects (γ, L), where L is an object of $\mathcal{F}(M, \boldsymbol{D})$ and $\gamma \in \Gamma$. Think of (γ, L) as representing the object $\gamma \cdot L$ of $\mathcal{F}(M, \boldsymbol{D})$, but we now have $|\Gamma|$ copies of each object.

We define an action of Γ on these objects, via

$$\gamma_1 \cdot (\gamma_2, L) := (\gamma_1 \cdot \gamma_2, L).$$

Now for each pair of objects $((1, L_0), (\gamma, L_1))$ of $\widetilde{\mathcal{F}}(M, \boldsymbol{D})$, we choose a regular Floer datum for the objects $(L_0, \gamma \cdot L_1)$ of $\mathcal{F}(M, \boldsymbol{D})$. We then define Floer data for pairs of objects $((\gamma_0, L_0), (\gamma_1, L_1))$ by acting with γ_0 on the Floer data for $((1, L_0), (\gamma_0^{-1} \cdot \gamma_1, L_1))$. We thus define morphism spaces

$$CF^*((\gamma_0, L_0), (\gamma_1, L_1))$$

for all pairs of objects. We define the Floer differential μ_0^1 as before, and note that it is now strictly Γ-equivariant.

Now for each tuple of objects $\boldsymbol{L} := ((1, L_0), (\gamma_1, L_1), \ldots, (\gamma_k, L_k))$, with associated generators $\boldsymbol{y}$, we choose regular, consistent perturbation data on the moduli spaces $\mathcal{R}(\boldsymbol{p}, \ell)$. We then define perturbation data for tuples $\boldsymbol{L} = ((\gamma_0, L_0), \ldots, (\gamma_k, L_k))$ by acting with γ_0 on the perturbation data chosen for $((1, L_0), (\gamma_0^{-1} \cdot \gamma_1, L_1), \ldots, (\gamma_0^{-1} \cdot \gamma_k, L_k))$. This allows us to define the rest of the Floer products μ^k. Note that they are strictly Γ-equivariant.

Observe now that the full subcategory with objects $(1, L)$ is equivalent to $\mathcal{F}(M, \boldsymbol{D})$ (making the corresponding choice of perturbation data). Thus, we have an inclusion of $\mathcal{F}(M, \boldsymbol{D})$ as a full subcategory of $\widetilde{\mathcal{F}}(M, \boldsymbol{D})$. Furthermore, if we restrict to the affine Fukaya category, then this inclusion is a quasi-equivalence, because each object (γ, L) is quasi-isomorphic to the element $(1, \gamma \cdot L)$ of the subcategory. This concludes the proof. □

References

[1] P. Seidel, "Fukaya categories and deformations," *Proceedings of the International Congress of Mathematicians (Beijing)* **2** (2002) 351–360, `arXiv:math/0206155`.

[2] P. Seidel, "Homological mirror symmetry for the quartic surface," `arXiv:0310414`.

[3] P. Seidel, *Fukaya categories and Picard-Lefschetz Theory.* European Mathematical Society, 2008.

[4] P. Seidel, "Homological mirror symmetry for the genus two curve," *Journal of Algebraic Geometry* **20** no. 4, (Dec., 2011) 727–769, `arXiv:0812.1171`.

[5] M. Abouzaid, K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, "Quantum cohomology and split generation in Lagrangian Floer theory," *in preparation* .

[6] P. Candelas, X. de la Ossa, P. Green, and L. Parkes, "A pair of Calabi-Yau manifolds as an exactly soluble superconformal theory," *Nuclear Physics B* **359** no. 1, (July, 1991) 21–74.

[7] A. Givental, "Equivariant Gromov-Witten invariants," *International Mathematics Research Notices* **1996** no. 13, (1996) 613–663, `arXiv:alg-geom/9603021`.

[8] A. Givental, "A mirror theorem for toric complete intersections," *Topological field theory, primitive forms and related topics (Kyoto, 1996), Progr. Math.* **160** (1998) 141–175.

[9] M. Kontsevich, "Homological algebra of mirror symmetry," *Proceedings of the International Congress of Mathematicians (Zürich, 1994)* (1994) 120–139, `arXiv:9411018`.

[10] A. Polishchuk and E. Zaslow, "Categorical mirror symmetry: the elliptic curve," *Adv. Theor. Math. Phys.* **2** (1998) 443–470, `arXiv:math/9801119`.

[11] A. Polishchuk, "Massey and Fukaya products on elliptic curves," *Adv. Theor. Math. Phys.* **4** (2000) 1187–1207, `arXiv:math.AG/9803017`.

[12] K. Fukaya, "Mirror symmetry of abelian varieties and multi-theta functions," *Journal of Algebraic Geometry* **11** (2002) 393–512.

[13] M. Abouzaid and I. Smith, "Homological mirror symmetry for the four-torus," *Duke Math. J.* **152** no. 3, (2010) 373–440, `arXiv:0903.3065`.

[14] M. Kontsevich and Y. Soibelman, "Homological mirror symmetry and torus fibrations," *Symplectic geometry and mirror symmetry, World Scientific* (2001) 203–263, `arXiv:math.SG/0011041`.

[15] Y. Nohara and K. Ueda, "A note on homological mirror symmetry for the quintic 3-fold," `arXiv:1103.4956`.

[16] N. Sheridan, "On the homological mirror symmetry conjecture for pairs of pants," *J. Diff. Geom* **89** no. 2, (2011) 271–367, `arXiv:1111.0632`.

[17] N. Sheridan, "Homological mirror symmetry for Fano hypersurfaces in projective space," *in progress* .

[18] K. Fukaya, Y.-g. Oh, H. Ohta, and K. Ono, "Anchored Lagrangian submanifolds and their Floer theory," *Mirror symmetry and tropical geometry, Contemp. Math.* **527** (2010) 15–54, `arXiv:0907.2122`.

[19] V. I. Arnold, "Characteristic class entering in quantization conditions," *Funkcional. Anal. i Priložen* **1** (1967) 1–14.

[20] K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, *Lagrangian intersection Floer theory – anomaly and obstruction.* American Mathematical Society, 2007.

[21] D. Orlov, "Derived categories of coherent sheaves and triangulated categories of singularities," *Algebra, Arithmetic, and Geometry* (2009) 503–531, `arXiv:math/0503632`.

[22] A. Polishchuk and A. Vaintrob, "Matrix factorizations and Cohomological Field Theories," `arXiv:1105.2903`.

[23] A. Căldăraru and J. Tu, "Curved A-infinity algebras and Landau-Ginzburg models," `arXiv:1007.2679`.

[24] A. Efimov, "Homological mirror symmetry for curves of higher genus," `arXiv:0907.3903`.

[25] G. Mikhalkin, "Decomposition into pairs-of-pants for complex algebraic hypersurfaces," *Topology* **43** no. 5, (2004) 1035–1065.

[26] O. Cornea and F. Lalonde, "Cluster Homology," `arXiv:0508345`.

[27] C. Weibel, *An introduction to homological algebra.* Cambridge University Press, Cambridge Studies in Advanced Mathematics, 38, 1994.

[28] G. Hochschild, B. Kostant, and A. Rosenberg, "Differential forms on regular affine algebras," *Trans. Amer. Math. Soc.* **102** (1962) 383–408.

[29] M. Gerstenhaber, "The cohomology structure of an associative ring," *Annals of Mathematics* **78** no. 2, (1963) 267–288.

[30] M. Kontsevich, "Deformation quantization of Poisson manifolds," *Letters in Mathematical Physics* **66** no. 3, (Dec., 2003) 157–216.

[31] D. McDuff and D. Salamon, *J-holomorphic Curves and Symplectic Topology.* American Mathematical Society Colloquium Publications. American Mathematical Society, 2004.

[32] E. Ionel and T. Parker, "Relative Gromov-Witten invariants," *Annals of Mathematics* **157** no. 1, (Jan., 2003) 45–96.

[33] U. Frauenfelder, "Gromov convergence of pseudoholomorphic disks," *J. Fixed Point Theory Appl.* **3** no. 2, (2008) 215–271.

[34] K. Cieliebak and K. Mohnke, "Symplectic hypersurfaces and transversality in Gromov-Witten theory," *Journal of Symplectic Geometry* **5** no. 3, (2007) 281–356, `arXiv:math/0702887`.

[35] A. Floer, H. Hofer, and D. Salamon, "Transversality in elliptic Morse theory for the symplectic action," *Duke Mathematical Journal* **80** no. 1, (1995) 251–292.

[36] Y.-G. Oh, "On the structure of pseudo-holomorphic discs with totally real boundary conditions," *J. Geom. Anal.* **7** no. 2, (1997) 305–327.

[37] M. Markl, "Transferring A-infinity (strongly homotopy associative) structures," *Rend. Circ. Mat. Palermo* **79** (2006) 139–151, `arXiv:math/0401007`.

[38] M. Markl, "Ideal perturbation lemma," *Communications in Algebra* **29** no. 11, (2001) 5209–5232, `arXiv:math/0002130`.

[39] D. Orlov, "Triangulated categories of singularities and D-branes in Landau-Ginzburg models," *Proc. Steklov Inst. Math.* **246** (2004) 227–248, `arXiv:math/0302304`.

[40] T. Dyckerhoff, "Compact generators in categories of matrix factorizations," *Duke Mathematical Journal* **159** no. 2, (Apr., 2011) 223–274, `arXiv:0904.4713`.

[41] M. Ballard, D. Favero, and L. Katzarkov, "Orlov spectra: bounds and gaps," `arXiv:1012.0864`.

[42] P. S. Aspinwall, "Landau-Ginzburg to Calabi-Yau dictionary for D-branes," *Journal of Mathematical Physics* **48** (2007) , `arXiv:math.AG/0503632`.

[43] A. A. Beilinson, “Coherent sheaves on $\mathbb{P}^n$ and problems of linear algebra,” *Functional Analysis and its Applications* **12** no. 3, (1978) 68–69.